

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/46407 A2(51) International Patent Classification⁷: C12N 15/12,
15/11, 15/63, 1/21, 5/10, C07K 14/705, 16/28, A01K
67/027, C12Q 1/68, G01N 33/68

(21) International Application Number: PCT/US01/45661

(22) International Filing Date: 5 December 2001 (05.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/729,094 5 December 2000 (05.12.2000) US(71) Applicant: PE CORPORATION (NY) [US/US]; 761
Main Avenue, Norwalk, CT 06859 (US).(72) Inventors: CHATURVEDI, Kabir; c/o Celera Genomics
Corporation, 45 West Gude Drive C2-4 #21, Rockville,MD 20850 (US). WEI, Ming-Hui; c/o Celera Genomics,
45 West Gude Drive C2-4#21, Rockville, MD 20850
(US). KETCHUM, Karen, A.; c/o Celera Genomics, 45
West Gude Drive C2-4#21, Rockville, MD 20850 (US).
DIFRANCESCO, Valentina; c/o Celera Genomics, 45
West Gude Drive C2-4#21, Rockville, MD 20850 (US).
BEASLEY, Ellen, M.; c/o Celera Genomics, 45 West
Gude Drive C2-4#21, Rockville, MD 20850 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZM, ZW.

[Continued on next page]

(54) Title: ISOLATED HUMAN TRANSPORTER PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN TRANS-
PORTER PROTEINS, AND USES THEREOF

1 GTCTCCCTCC CGCGCGATGG CCTCGGCGCT GAGCTATGTC TCCAAGTTCA
51 AGTCCTTCTGT GATCTTGTTC GTCCACCCGCT TCCTGCTGCT GCCACTCGTC
101 ATTCTGATGC CGGCCAAGTT TGTCAAGGTG GCCTACGTCA TCATCCTCAT
151 GGCCATTTC TGGTGACAG AAGTCATCCC TCTGGCTGTC ACCTCTCTCA
201 TGCTGTCTTT GCTTTTCCCA CTCCTCCAGA TTCTGGAATC CAGGCAGGTG
251 TGTGTCCAGT ACATGAAGGA CACCAACATG CTGTTCTCTG CGGCCTCAT
301 CGTGGCCGTG GCTGTGAGG GCTGGAACCT GCACAAGAGG ATCGCCCTGC
351 GCACGCTCTC CTGGGTGGGG GCCAAGCCTG CACGGCTGAT GCTGGGCTTC
401 ATGGGCGTCA CAGCCCTCCT GTCCATGTGG ATCAGTAACA TGGCAACCAC
451 GGCCATGATG GTGCCATCTG TGGAGGCCAT ATTGACGAG ATGGAAGCCA
501 CAAGCGCAGC CACCGAGGCC GGCCTGGAGC TGGTGACAA GGGCAAGGCC
551 AAGGAGCTGC CAGGAGGTCA AGTATTTTT GAAGGCCCA CTCTGGGGCA
601 GCAGGAAGAC CAAGAGCGGA AGAGTTTGTG TAAGGCCATG ACCCTGTGCA
651 TCTGTACGCG GGCCAGCATC GGGGGCACCG CCACCTGTAC CGGACGGGA
701 CCCAAGCTGG TGCTCTTGGG CCAGATGAAC GAGTTGTTT CTGACAGCAA
751 GGACCTCGTG AACTTTGCTT CTTGTTTGGC ATTTGCTTT CCCAATATGC
801 TGTGTATGCT GCTGTTCCGC TGGCTGTGGC TCCAGTTTGT TTACATGAGA
851 TTCAATTTTA AAAAGTCTGT GGGCTGCGGG CTAGAGAGCA AGAAAAACGA
901 GAAGGCTGCC CTCAGGTGCT TGCAGGAGGA GTACCGGAAG CTGGGGCCCT
951 TGTCCTTCGG GGAGATCAAC GTGCTGATCT GCTTCTTCTT GCTGGTATC
1001 CTGTGGTTCT CCCGAGACCC CGGCTTCATG CCGGCTGSCG TGAATGTTGC
1051 CTGGGTGGAG GGTGAGACAA AGTATGCTCT CGATGCCACT GTGGCCATCT
1101 TTGTGGCCAC CTTGCTATTC ATTGTGCTTT CACAGAAGCC CAAGTTTAAC
1151 TTCCGACGCG AGACTGAGGA AGAAAGGAAA ACTCCATTTT ATCCCTCTCC
1201 CCTGCTGGAT TGAAGGTAA CCCAGGAGAA AGTCCCTTGG GGCATCTGTC
1251 TGCTACTAGG GGGCGGATTT GCTCTGGCTA AAGGATCCGA GGCCTCGGGG
1301 CTGTCCGTGT GGATGGGGAA GCAGATGGAG CCCTTGACAG CAGTGCCCTC
1351 GGCAGCCATC ACCTTGATCT TGTCTTGTCT CGTTGCGGTG TTCAGTGAGT
1401 GCACAAGCAA CGTGGCCACC ACCACCTTGT TCCTGCCCAT CTTTGCCTCC
1451 ATGCTCTGCT CCATCGGCTC CAATCCGCTG TACATCATGC TGCCCTGTATC
1501 CCTGAGTGCC TCCTTTGCTT TCATGTTGCC TGTGGCCACC CTTCCAAATG
1551 CCATCGTGT CACCTATGGG CACCTCAAGG TTGCTGACAT GGTGAAAAAC
1601 GGAGTCATAA TGAACATAAT TGGAGTCTTC TGTGTGTTT TGGCTGTCAA
1651 CACCTGGGGA CGGGCCATAT TTGACTTGGA TCATTTCCCT GACTGGGCTA
1701 ATGTGACACA TATTGAGACT TAGGAAGAGC CACAAGACCA CACACACAG
1751 CTTACCCCTC CTCAGGACTA CGGAACCTTC TGGCACACCT TGTACAGAGT
1801 TTGGGGTTTC ACACCCCAAA ATGACCCAA CATTGTCACA CACCAACAAA
1851 ACCCAGCCAA TGGGCCACCT CTCTCTCCAA GCCCAGATGC ACAGATGGTC
1901 ATGGGAGCTT GGAGGGTAGG CTCAGAAATG AAGGGAAACC CTCAGTGGGG
1951 TGCTGGACCC ATCTTTCCCA AGCCTTGCCA TTATCTCTGT GAGGAGAGGC
2001 AGGTAGCCGA GGGATCAGGA TGCAGGCTGC TGTACCCGCT CTGCTCTAAG
2051 CATCCGCCAC ACAGGCTCTT GGTGTTTCACT CGCTTCGTCC TAGATAGTTT
2101 AATGGGAAT CAGATCCCTT GGTGAGAGC TAAGACAACC ACCTACCAAT
2151 GCCATGTCCT CTTCAGCTC ACCTTGAGCA GCCTCAGATC ATCTCTGTCA
2201 CTCTGGAGAG GACACCCGAG CCA (SEQ ID NO: 1)

FEATURES:

5'UTR: 1 - 16
Start Codon: 17
Stop Codon: 1721
3'UTR: 1724

(57) Abstract: The present invention provides amino acid
sequences of peptides that are encoded by genes within the human
genome, the transporter peptides of the present invention. The
present invention specifically provides isolated peptide and nucleic
acid molecules, methods of identifying orthologs and paralogs of
the transporter peptides, and methods of identifying modulators
of the transporter peptides.

WO 02/46407 A2



(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ISOLATED HUMAN TRANSPORTER PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS, AND USES THEREOF

RELATED APPLICATIONS

5 The present application claims priority to U.S. Application No. 09/729,094, filed December 5, 2000 (Atty. Docket CL000662).

FIELD OF THE INVENTION

10 The present invention is in the field of transporter proteins that are related to the sodium-dependent dicarboxylate transporter subfamily, recombinant DNA molecules, and protein production. The present invention specifically provides novel peptides and proteins that effect ligand transport and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

15

BACKGROUND OF THE INVENTION

Transporters

20 Transporter proteins regulate many different functions of a cell, including cell proliferation, differentiation, and signaling processes, by regulating the flow of molecules such as ions and macromolecules, into and out of cells. Transporters are found in the plasma membranes of virtually every cell in eukaryotic organisms. Transporters mediate a variety of cellular functions including regulation of membrane potentials and absorption and secretion of molecules and ion across cell membranes. When present in intracellular membranes of the Golgi apparatus and endocytic vesicles, transporters, such as chloride channels, also regulate organelle
25 pH. For a review, see Greger, R. (1988) Annu. Rev. Physiol. 50:111-122.

30 Transporters are generally classified by structure and the type of mode of action. In addition, transporters are sometimes classified by the molecule type that is transported, for example, sugar transporters, chlorine channels, potassium channels, etc. There may be many classes of channels for transporting a single type of molecule (a detailed review of channel types can be found at Alexander, S.P.H. and J.A. Peters: Receptor and transporter nomenclature supplement. Trends Pharmacol. Sci., Elsevier, pp. 65-68 (1997) and <http://www-biology.ucsd.edu/~msaier/transport/titlepage2.html>.

 The following general classification scheme is known in the art and is followed in the present discoveries.

Channel-type transporters. Transmembrane channel proteins of this class are ubiquitously found in the membranes of all types of organisms from bacteria to higher eukaryotes. Transport systems of this type catalyze facilitated diffusion (by an energy-independent process) by passage through a transmembrane aqueous pore or channel without evidence for a carrier-mediated mechanism. These channel proteins usually consist largely of α -helical spanners, although β -strands may also be present and may even comprise the channel. However, outer membrane porin-type channel proteins are excluded from this class and are instead included in class 9.

Carrier-type transporters. Transport systems are included in this class if they utilize a carrier-mediated process to catalyze uniport (a single species is transported by facilitated diffusion), antiport (two or more species are transported in opposite directions in a tightly coupled process, not coupled to a direct form of energy other than chemiosmotic energy) and/or symport (two or more species are transported together in the same direction in a tightly coupled process, not coupled to a direct form of energy other than chemiosmotic energy).

Carrier-type transporters include the Dicarboxylate/Amino Acid:Cation (Na^+ or H^+) Symporter ("DAACS") family, which catalyze Na^+ and/or H^+ symport together with (a) a Krebs cycle dicarboxylate (malate, succinate, or fumarate), (b) a dicarboxylic amino acid (glutamate or aspartate), (c) a small, semipolar, neutral amino acid (Ala, Ser, Cys, Thr), (d) both neutral and acidic amino acids or (e) most zwitterionic and dibasic amino acids. The bacterial members are of about 450 (420-491) amino acid residues while the mammalian proteins are of about 550 (503-574) residues in length. These proteins possess between ten and twelve putative transmembrane spanners (TMSs). A specific topological model in which 7 α -helical TMSs are followed by a reentrant loop-pore structure followed by one final TMS is presented in Slotboom et al., *Microbiol. Mol. Biol. Rev.* 63: 293-307 (1999). All of the bacterial proteins cluster together on the phylogenetic tree as do the mammalian proteins. The mammalian permeases that transport neutral amino acids cluster separately from those that are specific for the acidic amino acids. Among the mammalian proteins are neuronal excitatory amino acid neurotransmitter permeases.

Pyrophosphate bond hydrolysis-driven active transporters. Transport systems are included in this class if they hydrolyze pyrophosphate or the terminal pyrophosphate bond in ATP or another nucleoside triphosphate to drive the active uptake and/or extrusion of a solute or solutes. The transport protein may or may not be transiently phosphorylated, but the substrate is not phosphorylated.

PEP-dependent, phosphoryl transfer-driven group translocators. Transport systems of the bacterial phosphoenolpyruvate:sugar phosphotransferase system are included in this class. The product of the reaction, derived from extracellular sugar, is a cytoplasmic sugar-phosphate.

Decarboxylation-driven active transporters. Transport systems that drive solute (e.g., ion) uptake or extrusion by decarboxylation of a cytoplasmic substrate are included in this class.

Oxidoreduction-driven active transporters. Transport systems that drive transport of a solute (e.g., an ion) energized by the flow of electrons from a reduced substrate to an oxidized substrate are included in this class.

Light-driven active transporters. Transport systems that utilize light energy to drive transport of a solute (e.g., an ion) are included in this class.

Mechanically-driven active transporters. Transport systems are included in this class if they drive movement of a cell or organelle by allowing the flow of ions (or other solutes) through the membrane down their electrochemical gradients.

Outer-membrane porins (of b-structure). These proteins form transmembrane pores or channels that usually allow the energy independent passage of solutes across a membrane. The transmembrane portions of these proteins consist exclusively of b-strands that form a b-barrel. These porin-type proteins are found in the outer membranes of Gram-negative bacteria, mitochondria and eukaryotic plastids.

Methyltransferase-driven active transporters. A single characterized protein currently falls into this category, the Na⁺-transporting methyltetrahydromethanopterin:coenzyme M methyltransferase.

Non-ribosome-synthesized channel-forming peptides or peptide-like molecules. These molecules, usually chains of L- and D-amino acids as well as other small molecular building blocks such as lactate, form oligomeric transmembrane ion channels. Voltage may induce channel formation by promoting assembly of the transmembrane channel. These peptides are often made by bacteria and fungi as agents of biological warfare.

Non-Proteinaceous Transport Complexes. Ion conducting substances in biological membranes that do not consist of or are not derived from proteins or peptides fall into this category.

Functionally characterized transporters for which sequence data are lacking. Transporters of particular physiological significance will be included in this category even though a family assignment cannot be made.

Putative transporters in which no family member is an established transporter. Putative transport protein families are grouped under this number and will either be classified elsewhere

when the transport function of a member becomes established, or will be eliminated from the TC classification system if the proposed transport function is disproven. These families include a member or members for which a transport function has been suggested, but evidence for such a function is not yet compelling.

5 Auxiliary transport proteins. Proteins that in some way facilitate transport across one or more biological membranes but do not themselves participate directly in transport are included in this class. These proteins always function in conjunction with one or more transport proteins. They may provide a function connected with energy coupling to transport, play a structural role in complex formation or serve a regulatory function.

10 Transporters of unknown classification. Transport protein families of unknown classification are grouped under this number and will be classified elsewhere when the transport process and energy coupling mechanism are characterized. These families include at least one member for which a transport function has been established, but either the mode of transport or the energy coupling mechanism is not known.

15

Ion channels

An important type of transporter is the ion channel. Ion channels regulate many different cell proliferation, differentiation, and signaling processes by regulating the flow of ions into and out of cells. Ion channels are found in the plasma membranes of virtually every cell in eukaryotic organisms. Ion channels mediate a variety of cellular functions including regulation of membrane potentials and absorption and secretion of ion across epithelial membranes. When present in intracellular membranes of the Golgi apparatus and endocytic vesicles, ion channels, such as chloride channels, also regulate organelle pH. For a review, see Greger, R. (1988) Annu. Rev. Physiol. 50:111-122.

20 Ion channels are generally classified by structure and the type of mode of action. For example, extracellular ligand gated channels (ELGs) are comprised of five polypeptide subunits, with each subunit having 4 membrane spanning domains, and are activated by the binding of an extracellular ligand to the channel. In addition, channels are sometimes classified by the ion type that is transported, for example, chlorine channels, potassium channels, etc. There may be many classes of channels for transporting a single type of ion (a detailed review of channel types can be found at Alexander, S.P.H. and J.A. Peters (1997). Receptor and ion channel nomenclature supplement. Trends Pharmacol. Sci., Elsevier, pp. 65-68 and <http://www-biology.ucsd.edu/~msaier/transport/toc.html>.

There are many types of ion channels based on structure. For example, many ion channels fall within one of the following groups: extracellular ligand-gated channels (ELG), intracellular ligand-gated channels (ILG), inward rectifying channels (INR), intercellular (gap junction) channels, and voltage gated channels (VIC). There are additionally recognized other
 5 channel families based on ion-type transported, cellular location and drug sensitivity. Detailed information on each of these, their activity, ligand type, ion type, disease association, drugability, and other information pertinent to the present invention, is well known in the art.

Extracellular ligand-gated channels, ELGs, are generally comprised of five polypeptide subunits, Unwin, N. (1993), *Cell* 72: 31-41; Unwin, N. (1995), *Nature* 373: 37-43; Hucho, F., et al., (1996) *J. Neurochem.* 66: 1781-1792; Hucho, F., et al., (1996) *Eur. J. Biochem.* 239: 539-
 10 557; Alexander, S.P.H. and J.A. Peters (1997), *Trends Pharmacol. Sci.*, Elsevier, pp. 4-6; 36-40; 42-44; and Xue, H. (1998) *J. Mol. Evol.* 47: 323-333. Each subunit has 4 membrane spanning regions: this serves as a means of identifying other members of the ELG family of proteins. ELG bind a ligand and in response modulate the flow of ions. Examples of ELG include most
 15 members of the neurotransmitter-receptor family of proteins, e.g., GABAI receptors. Other members of this family of ion channels include glycine receptors, ryandyne receptors, and ligand gated calcium channels.

The Voltage-gated Ion Channel (VIC) Superfamily

Proteins of the VIC family are ion-selective channel proteins found in a wide range of
 20 bacteria, archaea and eukaryotes Hille, B. (1992), Chapter 9: Structure of channel proteins; Chapter 20: Evolution and diversity. In: *Ionic Channels of Excitable Membranes*, 2nd Ed., Sinaur Assoc. Inc., Pubs., Sunderland, Massachusetts; Sigworth, F.J. (1993), *Quart. Rev. Biophys.* 27: 1-40; Salkoff, L. and T. Jegla (1995), *Neuron* 15: 489-492; Alexander, S.P.H. et al., (1997), *Trends Pharmacol. Sci.*, Elsevier, pp. 76-84; Jan, L.Y. et al., (1997), *Annu. Rev. Neurosci.* 20: 91-123; Doyle, D.A, et al., (1998) *Science* 280: 69-77; Terlau, H. and W. Stühmer
 25 (1998), *Naturwissenschaften* 85: 437-444. They are often homo- or heterooligomeric structures with several dissimilar subunits (e.g., α_1 - α_2 - δ - β Ca^{2+} channels, $\alpha_1\beta_2$ Na^+ channels or $(\alpha)_4\beta$ K^+ channels), but the channel and the primary receptor is usually associated with the α (or α_1) subunit. Functionally characterized members are specific for K^+ , Na^+ or Ca^{2+} . The K^+ channels
 30 usually consist of homotetrameric structures with each α -subunit possessing six transmembrane spanners (TMSs). The α_1 and α subunits of the Ca^{2+} and Na^+ channels, respectively, are about four times as large and possess 4 units, each with 6 TMSs separated by a hydrophilic loop, for a total of 24 TMSs. These large channel proteins form heterotetra-unit structures equivalent to the homotetrameric structures of most K^+ channels. All four units of the Ca^{2+} and Na^+ channels are

homologous to the single unit in the homotetrameric K^+ channels. Ion flux via the eukaryotic channels is generally controlled by the transmembrane electrical potential (hence the designation, voltage-sensitive) although some are controlled by ligand or receptor binding.

Several putative K^+ -selective channel proteins of the VIC family have been identified in prokaryotes. The structure of one of them, the KcsA K^+ channel of *Streptomyces lividans*, has been solved to 3.2 Å resolution. The protein possesses four identical subunits, each with two transmembrane helices, arranged in the shape of an inverted teepee or cone. The cone cradles the "selectivity filter" P domain in its outer end. The narrow selectivity filter is only 12 Å long, whereas the remainder of the channel is wider and lined with hydrophobic residues. A large water-filled cavity and helix dipoles stabilize K^+ in the pore. The selectivity filter has two bound K^+ ions about 7.5 Å apart from each other. Ion conduction is proposed to result from a balance of electrostatic attractive and repulsive forces.

In eukaryotes, each VIC family channel type has several subtypes based on pharmacological and electrophysiological data. Thus, there are five types of Ca^{2+} channels (L, N, P, Q and T). There are at least ten types of K^+ channels, each responding in different ways to different stimuli: voltage-sensitive [K_a, K_v, K_{vr}, K_{vs} and K_{sr}], Ca^{2+} -sensitive [BK_{Ca}, IK_{Ca} and SK_{Ca}] and receptor-coupled [K_M and K_{ACH}]. There are at least six types of Na^+ channels (I, II, III, μ 1, H1 and PN3). Tetrameric channels from both prokaryotic and eukaryotic organisms are known in which each α -subunit possesses 2 TMSs rather than 6, and these two TMSs are homologous to TMSs 5 and 6 of the six TMS unit found in the voltage-sensitive channel proteins. KcsA of *S. lividans* is an example of such a 2 TMS channel protein. These channels may include the K_{Na} (Na^+ -activated) and K_{Vol} (cell volume-sensitive) K^+ channels, as well as distantly related channels such as the Tok1 K^+ channel of yeast, the TWIK-1 inward rectifier K^+ channel of the mouse and the TREK-1 K^+ channel of the mouse. Because of insufficient sequence similarity with proteins of the VIC family, inward rectifier K^+ IRK channels (ATP-regulated; G-protein-activated) which possess a P domain and two flanking TMSs are placed in a distinct family. However, substantial sequence similarity in the P region suggests that they are homologous. The b, g and d subunits of VIC family members, when present, frequently play regulatory roles in channel activation/deactivation.

30 The Epithelial Na^+ Channel (ENaC) Family

The ENaC family consists of over twenty-four sequenced proteins (Canessa, C.M., et al., (1994), Nature 367: 463-467, Le, T. and M.H. Saier, Jr. (1996), Mol. Membr. Biol. 13: 149-157; Garty, H. and L.G. Palmer (1997), Physiol. Rev. 77: 359-396; Waldmann, R., et al., (1997), Nature 386: 173-177; Darboux, I., et al., (1998), J. Biol. Chem. 273: 9424-9429; Firsov, D., et

al., (1998), EMBO J. 17: 344-352; Horisberger, J.-D. (1998). Curr. Opin. Struc. Biol. 10: 443-449). All are from animals with no recognizable homologues in other eukaryotes or bacteria. The vertebrate ENaC proteins from epithelial cells cluster tightly together on the phylogenetic tree: voltage-insensitive ENaC homologues are also found in the brain. Eleven sequenced *C. elegans* proteins, including the degenerins, are distantly related to the vertebrate proteins as well as to each other. At least some of these proteins form part of a mechano-transducing complex for touch sensitivity. The homologous *Helix aspersa* (FMRF-amide)-activated Na⁺ channel is the first peptide neurotransmitter-gated ionotropic receptor to be sequenced.

Protein members of this family all exhibit the same apparent topology, each with N- and C-termini on the inside of the cell, two amphipathic transmembrane spanning segments, and a large extracellular loop. The extracellular domains contain numerous highly conserved cysteine residues. They are proposed to serve a receptor function.

Mammalian ENaC is important for the maintenance of Na⁺ balance and the regulation of blood pressure. Three homologous ENaC subunits, alpha, beta, and gamma, have been shown to assemble to form the highly Na⁺-selective channel. The stoichiometry of the three subunits is alpha₂beta₁gamma₁ in a heterotetrameric architecture.

The Glutamate-gated Ion Channel (GIC) Family of Neurotransmitter Receptors

Members of the GIC family are heteropentameric complexes in which each of the 5 subunits is of 800-1000 amino acid residues in length (Nakanishi, N., et al, (1990), Neuron 5: 569-581; Unwin, N. (1993), Cell 72: 31-41; Alexander, S.P.H. and J.A. Peters (1997) Trends Pharmacol. Sci., Elsevier, pp. 36-40). These subunits may span the membrane three or five times as putative α -helices with the N-termini (the glutamate-binding domains) localized extracellularly and the C-termini localized cytoplasmically. They may be distantly related to the ligand-gated ion channels, and if so, they may possess substantial β -structure in their transmembrane regions. However, homology between these two families cannot be established on the basis of sequence comparisons alone. The subunits fall into six subfamilies: a, b, g, d, e and z.

The GIC channels are divided into three types: (1) α -amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA)-, (2) kainate- and (3) N-methyl-D-aspartate (NMDA)-selective glutamate receptors. Subunits of the AMPA and kainate classes exhibit 35-40% identity with each other while subunits of the NMDA receptors exhibit 22-24% identity with the former subunits. They possess large N-terminal, extracellular glutamate-binding domains that are homologous to the periplasmic glutamine and glutamate receptors of ABC-type uptake permeases of Gram-negative bacteria. All known members of the GIC family are from animals.

The different channel (receptor) types exhibit distinct ion selectivities and conductance properties. The NMDA-selective large conductance channels are highly permeable to monovalent cations and Ca^{2+} . The AMPA- and kainate-selective ion channels are permeable primarily to monovalent cations with only low permeability to Ca^{2+} .

5 The Chloride Channel (ClC) Family

The ClC family is a large family consisting of dozens of sequenced proteins derived from Gram-negative and Gram-positive bacteria, cyanobacteria, archaea, yeast, plants and animals (Steinmeyer, K., et al., (1991), Nature 354: 301-304; Uchida, S., et al., (1993), J. Biol. Chem. 268: 3821-3824; Huang, M.-E., et al., (1994), J. Mol. Biol. 242: 595-598; Kawasaki, M., et al, 10 (1994), Neuron 12: 597-604; Fisher, W.E., et al., (1995), Genomics. 29:598-606; and Foskett, J.K. (1998), Annu. Rev. Physiol. 60: 689-717). These proteins are essentially ubiquitous, although they are not encoded within genomes of *Haemophilus influenzae*, *Mycoplasma genitalium*, and *Mycoplasma pneumoniae*. Sequenced proteins vary in size from 395 amino acyl residues (*M. jannaschii*) to 988 residues (man). Several organisms contain multiple ClC family 15 paralogues. For example, *Synechocystis* has two paralogues, one of 451 residues in length and the other of 899 residues. *Arabidopsis thaliana* has at least four sequenced paralogues, (775-792 residues), humans also have at least five paralogues (820-988 residues), and *C. elegans* also has at least five (810-950 residues). There are nine known members in mammals, and mutations in three of the corresponding genes cause human diseases. *E. coli*, *Methanococcus jannaschii* and 20 *Saccharomyces cerevisiae* only have one ClC family member each. With the exception of the larger *Synechocystis* paralogue, all bacterial proteins are small (395-492 residues) while all eukaryotic proteins are larger (687-988 residues). These proteins exhibit 10-12 putative transmembrane α -helical spanners (TMSs) and appear to be present in the membrane as homodimers. While one member of the family, *Torpedo* ClC-O, has been reported to have two 25 channels, one per subunit, others are believed to have just one.

All functionally characterized members of the ClC family transport chloride, some in a voltage-regulated process. These channels serve a variety physiological functions (cell volume regulation; membrane potential stabilization; signal transduction; transepithelial transport, etc.). Different homologues in humans exhibit differing anion selectivities, i.e., ClC4 and ClC5 share a 30 $\text{NO}_3^- > \text{Cl}^- > \text{Br}^- > \text{I}^-$ conductance sequence, while ClC3 has an $\text{I}^- > \text{Cl}^-$ selectivity. The ClC4 and ClC5 channels and others exhibit outward rectifying currents with currents only at voltages more positive than +20mV.

Animal Inward Rectifier K^+ Channel (IRK-C) Family

IRK channels possess the "minimal channel-forming structure" with only a P domain, characteristic of the channel proteins of the VIC family, and two flanking transmembrane spanners (Shuck, M.E., et al., (1994), J. Biol. Chem. 269: 24261-24270; Ashen, M.D., et al., (1995), Am. J. Physiol. 268: H506-H511; Salkoff, L. and T. Jegla (1995), Neuron 15: 489-492; Aguilar-Bryan, L., et al., (1998), Physiol. Rev. 78: 227-245; Ruknudin, A., et al., (1998), J. Biol. Chem. 273: 14165-14171). They may exist in the membrane as homo- or heterooligomers. They have a greater tendency to let K^+ flow into the cell than out. Voltage-dependence may be regulated by external K^+ , by internal Mg^{2+} , by internal ATP and/or by G-proteins. The P domains of IRK channels exhibit limited sequence similarity to those of the VIC family, but this sequence similarity is insufficient to establish homology. Inward rectifiers play a role in setting cellular membrane potentials, and the closing of these channels upon depolarization permits the occurrence of long duration action potentials with a plateau phase. Inward rectifiers lack the intrinsic voltage sensing helices found in VIC family channels. In a few cases, those of Kir1.1a and Kir6.2, for example, direct interaction with a member of the ABC superfamily has been proposed to confer unique functional and regulatory properties to the heteromeric complex, including sensitivity to ATP. The SUR1 sulfonylurea receptor (spQ09428) is the ABC protein that regulates the Kir6.2 channel in response to ATP, and CFTR may regulate Kir1.1a. Mutations in SUR1 are the cause of familial persistent hyperinsulinemic hypoglycemia in infancy (PHHI), an autosomal recessive disorder characterized by unregulated insulin secretion in the pancreas.

20 ATP-gated Cation Channel (ACC) Family

Members of the ACC family (also called P2X receptors) respond to ATP, a functional neurotransmitter released by exocytosis from many types of neurons (North, R.A. (1996), Curr. Opin. Cell Biol. 8: 474-483; Soto, F., M. Garcia-Guzman and W. Stühmer (1997), J. Membr. Biol. 160: 91-100). They have been placed into seven groups ($P2X_1$ - $P2X_7$) based on their pharmacological properties. These channels, which function at neuron-neuron and neuron-smooth muscle junctions, may play roles in the control of blood pressure and pain sensation. They may also function in lymphocyte and platelet physiology. They are found only in animals.

The proteins of the ACC family are quite similar in sequence (>35% identity), but they possess 380-1000 amino acid residues per subunit with variability in length localized primarily to the C-terminal domains. They possess two transmembrane spanners, one about 30-50 residues from their N-termini, the other near residues 320-340. The extracellular receptor domains between these two spanners (of about 270 residues) are well conserved with numerous conserved glycyl and cysteyl residues. The hydrophilic C-termini vary in length from 25 to 240 residues. They resemble the topologically similar epithelial Na^+ channel (ENaC) proteins in possessing (a)

N- and C-termini localized intracellularly, (b) two putative transmembrane spanners, (c) a large extracellular loop domain, and (d) many conserved extracellular cysteal residues. ACC family members are, however, not demonstrably homologous with them. ACC channels are probably hetero- or homomultimers and transport small monovalent cations (Me^+). Some also transport Ca^{2+} ; a few also transport small metabolites.

The Ryanodine-Inositol 1,4,5-triphosphate Receptor Ca^{2+} Channel (RIR-CaC) Family

Ryanodine (Ry)-sensitive and inositol 1,4,5-triphosphate (IP_3)-sensitive Ca^{2+} -release channels function in the release of Ca^{2+} from intracellular storage sites in animal cells and thereby regulate various Ca^{2+} -dependent physiological processes (Hasan, G. et al., (1992) Development 116: 967-975; Michikawa, T., et al., (1994), J. Biol. Chem. 269: 9184-9189; Tunwell, R.E.A., (1996), Biochem. J. 318: 477-487; Lee, A.G. (1996) *Biomembranes*, Vol. 6, Transmembrane Receptors and Channels (A.G. Lee, ed.), JAI Press, Denver, CO., pp 291-326; Mikoshiba, K., et al., (1996) J. Biochem. Biomem. 6: 273-289). Ry receptors occur primarily in muscle cell sarcoplasmic reticular (SR) membranes, and IP_3 receptors occur primarily in brain cell endoplasmic reticular (ER) membranes where they effect release of Ca^{2+} into the cytoplasm upon activation (opening) of the channel.

The Ry receptors are activated as a result of the activity of dihydropyridine-sensitive Ca^{2+} channels. The latter are members of the voltage-sensitive ion channel (VIC) family. Dihydropyridine-sensitive channels are present in the T-tubular systems of muscle tissues.

Ry receptors are homotetrameric complexes with each subunit exhibiting a molecular size of over 500,000 daltons (about 5,000 amino acid residues). They possess C-terminal domains with six putative transmembrane α -helical spanners (TMSs). Putative pore-forming sequences occur between the fifth and sixth TMSs as suggested for members of the VIC family. The large N-terminal hydrophilic domains and the small C-terminal hydrophilic domains are localized to the cytoplasm. Low resolution 3-dimensional structural data are available. Mammals possess at least three isoforms which probably arose by gene duplication and divergence before divergence of the mammalian species. Homologues are present in humans and *Caenorabditis elegans*.

IP_3 receptors resemble Ry receptors in many respects. (1) They are homotetrameric complexes with each subunit exhibiting a molecular size of over 300,000 daltons (about 2,700 amino acid residues). (2) They possess C-terminal channel domains that are homologous to those of the Ry receptors. (3) The channel domains possess six putative TMSs and a putative channel lining region between TMSs 5 and 6. (4) Both the large N-terminal domains and the smaller C-terminal tails face the cytoplasm. (5) They possess covalently linked carbohydrate on

extracytoplasmic loops of the channel domains. (6) They have three currently recognized isoforms (types 1, 2, and 3) in mammals which are subject to differential regulation and have different tissue distributions.

IP₃ receptors possess three domains: N-terminal IP₃-binding domains, central coupling or regulatory domains and C-terminal channel domains. Channels are activated by IP₃ binding, and like the Ry receptors, the activities of the IP₃ receptor channels are regulated by phosphorylation of the regulatory domains, catalyzed by various protein kinases. They predominate in the endoplasmic reticular membranes of various cell types in the brain but have also been found in the plasma membranes of some nerve cells derived from a variety of tissues.

The channel domains of the Ry and IP₃ receptors comprise a coherent family that in spite of apparent structural similarities, do not show appreciable sequence similarity of the proteins of the VIC family. The Ry receptors and the IP₃ receptors cluster separately on the RIR-CaC family tree. They both have homologues in *Drosophila*. Based on the phylogenetic tree for the family, the family probably evolved in the following sequence: (1) A gene duplication event occurred that gave rise to Ry and IP₃ receptors in invertebrates. (2) Vertebrates evolved from invertebrates. (3) The three isoforms of each receptor arose as a result of two distinct gene duplication events. (4) These isoforms were transmitted to mammals before divergence of the mammalian species.

The Organellar Chloride Channel (O-ClC) Family

Proteins of the O-ClC family are voltage-sensitive chloride channels found in intracellular membranes but not the plasma membranes of animal cells (Landry, D, et al., (1993), J. Biol. Chem. 268: 14948-14955; Valenzuela, Set al., (1997), J. Biol. Chem. 272: 12575-12582; and Duncan, R.R., et al., (1997), J. Biol. Chem. 272: 23880-23886).

They are found in human nuclear membranes, and the bovine protein targets to the microsomes, but not the plasma membrane, when expressed in *Xenopus laevis* oocytes. These proteins are thought to function in the regulation of the membrane potential and in transepithelial ion absorption and secretion in the kidney. They possess two putative transmembrane α -helical spanners (TMSs) with cytoplasmic N- and C-termini and a large luminal loop that may be glycosylated. The bovine protein is 437 amino acid residues in length and has the two putative TMSs at positions 223-239 and 367-385. The human nuclear protein is much smaller (241 residues). A *C. elegans* homologue is 260 residues long.

The protein of the present invention is very similar to the dicarboxylate transporters. They bind a variety of divalent organic anions. Some of these carriers import acetylaspartate into the glial cells and play an important role in myelination. Others maintain succinate levels in

placenta and kidneys. Those expressed in the renal brush border may be relevant to pharmacological research. This sequence is also homologous to the family of sodium-sulfate transporters, which carry divalent inorganic anions across the cell membrane. Like its homologues, this transporter has 12 transmembrane helices.

5 Mitochondria and perhaps other organelles contain dicarboxylate transporters, which pump organic acids in and out of these compartments. Spatial distribution of divalent acids may affect the rates of the Krebs cycle, amino acid synthesis and other ergogenic and metabolic pathways. Sometimes, their local concentration exceeds physiological levels, which leads to formation of calcium stones.

10 The sequence presented here can be used to search for the specific interactors using affinity chromatography and the yeast two-hybrid system. Synthetic peptides and cytrate-derived compounds can be designed and used as inhibitors for these transporters.

For a review related to the dicarboxylate transporters, see references by Huang *et al.*, *J Pharmacol Exp Ther* 2000 Oct;295(1):392-403, Chen *et al.*, *J Biol Chem* 1998 Aug 14;273(33):20972-81, Pajor, *J Biol Chem* 1995 Mar 17;270(11):5779-85, Wang *et al.*, *Am J Physiol Cell Physiol* 2000 May;278(5):C1019-30, Chen *et al.*, *Arch Biochem Biophys* 2000 Jan 1;373(1):193-202.

Transporter proteins, particularly members of the sodium-dependent dicarboxylate transporter subfamily, are a major target for drug action and development. Accordingly, it is 20 valuable to the field of pharmaceutical development to identify and characterize previously unknown transport proteins. The present invention advances the state of the art by providing a previously unidentified human transport proteins.

SUMMARY OF THE INVENTION

25 The present invention is based in part on the identification of amino acid sequences of human transporter peptides and proteins that are related to the sodium-dependent dicarboxylate transporter subfamily, as well as allelic variants and other mammalian orthologs thereof. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic 30 proteins, and serve as targets for the development of human therapeutic agents that modulate transporter activity in cells and tissues that express the transporter. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen.

DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides the nucleotide sequence of a cDNA molecule or transcript sequence that encodes the transporter protein of the present invention. In addition structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence.

5 Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen.

FIGURE 2 provides the predicted amino acid sequence of the transporter of the present invention. In addition structure and functional information such as protein family, function, and modification sites is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

10 FIGURE 3 provides genomic sequences that span the gene encoding the transporter protein of the present invention. In addition structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence. 55 SNPs, including 4 indels, have been identified in the gene encoding the transporter protein provided by

15 the present invention and are given in Figure 3.

DETAILED DESCRIPTION OF THE INVENTION

General Description

The present invention is based on the sequencing of the human genome. During the

20 sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a transporter protein or part of a transporter protein and are related to the sodium-dependent dicarboxylate transporter subfamily. Utilizing these sequences, additional

25 genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human transporter peptides and proteins that are related to the sodium-dependent dicarboxylate transporter subfamily, nucleic acid sequences in the form of transcript sequences, cDNA

30 sequences and/or genomic sequences that encode these transporter peptides and proteins, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the transporter of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially

important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known transporter proteins of the sodium-dependent dicarboxylate transporter subfamily and the expression pattern observed. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known sodium-dependent dicarboxylate transporter family or subfamily of transporter proteins.

Specific Embodiments

Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the transporter family of proteins and are related to the sodium-dependent dicarboxylate transporter subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figures 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the transporter peptides of the present invention, transporter peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprising the amino acid sequences of the transporter peptides disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the transporter peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated transporter peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. For example, a nucleic acid molecule encoding the transporter peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the transporter peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The transporter peptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a transporter peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the transporter peptide. "Operatively linked" indicates that the transporter peptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the transporter peptide.

In some uses, the fusion protein does not affect the activity of the transporter peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant transporter peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A transporter peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the transporter peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the transporter peptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is

determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the transporter peptides of the present invention as well as being encoded by the same genetic locus as the transporter peptide provided herein. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel transporter maps to public BAC AC034305, which is known to be located on human chromosome 17.

Allelic variants of a transporter peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the transporter peptide as well as being encoded by the same genetic locus as the transporter peptide provided herein. Genetic locus can readily be determined based on the genomic information

provided in Figure 3, such as the genomic sequence mapped to the reference human. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel transporter maps to public BAC AC034305, which is known to be located on human chromosome 17. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Paralogs of a transporter peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the transporter peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a transporter peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the transporter peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a transporter peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the transporter peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the transporter peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a transporter peptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic

residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant transporter peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind ligand, ability to transport ligand, ability to mediate signaling, etc.

5 Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis and can be used to identify critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

10 Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 15 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as transporter activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or 20 photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the transporter peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to 25 be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a transporter peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the transporter peptide or could be chosen for the 30 ability to perform a function, e.g. bind a substrate or act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the transporter peptide, e.g., active site, a transmembrane domain or a substrate-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide

fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in transporter peptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the transporter peptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature transporter peptide is fused with another compound, such as a compound to increase the half-life of the transporter peptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature transporter peptide, such as a leader or secretory sequence or a sequence for purification of the mature transporter peptide or a pro-protein sequence.

Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding partner or ligand) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein or ligand (such as, for example, in a transporter-effector protein interaction or transporter-ligand interaction), the protein can be used to identify the binding partner/ligand so as to develop a system to identify inhibitors of the binding interaction. Any or all of these uses are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

Substantial chemical and structural homology exists between the dicarboxylate transporter protein described herein and dicarboxylate transporters (see Figure 1). As discussed in the background, dicarboxylate transporters are known in the art to be involved in the major determinant of urinary excretion of citrate, the potent inhibitor of calcium salt crystallization urinary excretion of citrate, the potent inhibitor of calcium salt crystallization. Accordingly, the dicarboxylate transporter, and the encoding gene, provided by the present invention is useful for treating, preventing, and/or diagnosing dicarboxylate transporter related diseases such as kidney disorder.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, transporters isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the transporter. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual Northern blot shows expression in fetal liver and spleen. A large percentage of pharmaceutical agents are being developed that modulate the activity of transporter proteins,

particularly members of the sodium-dependent dicarboxylate transporter subfamily (see Background of the Invention). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1.

5 Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Such uses can readily be determined using the information provided herein, that known in the art and routine experimentation.

The transporter polypeptides (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to transporters that

10 are related to members of the sodium-dependent dicarboxylate transporter subfamily. Such assays involve any of the known transporter functions or activities or properties useful for diagnosis and treatment of transporter-related conditions that are specific for the subfamily of transporters that the one of the present invention belongs to, particularly in cells and tissues that express the transporter. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen.

15 Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver.

The transporter polypeptides are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the transporter, as a biopsy or expanded in cell culture. Experimental data as provided in FIGURE 1 indicates

20 expression in the fetal liver and spleen. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the transporter protein.

The polypeptides can be used to identify compounds that modulate transporter activity of the protein in its natural state or an altered form that causes a specific disease or pathology associated with the transporter. Both the transporters of the present invention and appropriate

25 variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the transporter. These compounds can be further screened against a functional transporter to determine the effect of the compound on the transporter activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the transporter to a

30 desired degree.

Further, the transporter polypeptides can be used to screen a compound for the ability to stimulate or inhibit interaction between the transporter protein and a molecule that normally interacts with the transporter protein, e.g. a substrate or a component of the signal pathway that the transporter protein normally interacts (for example, another transporter). Such assays typically

include the steps of combining the transporter protein with a candidate compound under conditions that allow the transporter protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the transporter protein and the target, such as any of the associated effects of signal transduction such as changes in membrane potential, protein phosphorylation, cAMP turnover, and adenylate cyclase activation, etc.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for ligand binding. Other candidate compounds include mutant transporters or appropriate fragments containing mutations that affect transporter function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity, or a fragment that binds ligand but does not allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) transporter activity. The assays typically involve an assay of events in the signal transduction pathway that indicate transporter activity. Thus, the transport of a ligand, change in cell membrane potential, activation of a protein, a change in the expression of genes that are up- or down-regulated in response to the transporter protein dependent signal cascade can be assayed.

Any of the biological or biochemical functions mediated by the transporter can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the transporter can be assayed. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot

shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver.

Binding and/or activating compounds can also be screened by using chimeric transporter proteins in which the amino terminal extracellular domain, or parts thereof, the entire
5 transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a ligand-binding region can be used that interacts with a different ligand than that which is recognized by the native transporter. Accordingly, a different set of signal transduction components is available as an end-
10 point assay for activation. This allows for assays to be performed in other than the specific host cell from which the transporter is derived.

The transporter polypeptides are also useful in competition binding assays in methods designed to discover compounds that interact with the transporter (e.g. binding partners and/or ligands). Thus, a compound is exposed to a transporter polypeptide under conditions that allow the
15 compound to bind or to otherwise interact with the polypeptide. Soluble transporter polypeptide is also added to the mixture. If the test compound interacts with the soluble transporter polypeptide, it decreases the amount of complex formed or activity from the transporter target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the transporter. Thus, the soluble polypeptide that competes with the target transporter region is
20 designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the transporter protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

25 Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ³⁵S-labeled) and the candidate
30 compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of transporter-binding protein found in the

bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a transporter-binding protein and a candidate compound are incubated in the transporter protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the transporter protein target molecule, or which are reactive with transporter protein and compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the transporters of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of transporter protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the transporter pathway, by treating cells or tissues that express the transporter. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. These methods of treatment include the steps of administering a modulator of transporter activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the transporter proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the transporter and are involved in transporter activity. Such transporter-binding proteins are also likely to be involved in the propagation of signals by the transporter proteins or transporter targets as, for example, downstream elements of a transporter-mediated signaling pathway. Alternatively, such transporter-binding proteins are likely to be transporter inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a transporter protein is fused

to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a transporter-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the transporter protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a transporter-modulating agent, an antisense transporter nucleic acid molecule, a transporter-specific antibody, or a transporter-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The transporter proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. The method involves contacting a biological sample with a compound capable of interacting with the transporter protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the

present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered transporter activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

10 *In vitro* techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the transporter protein in which one or more of the transporter functions in one population is different from those in another population. The peptides thus allow a

target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other ligand-binding regions that are more or less active in ligand binding, and transporter activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect
5 within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Accordingly, methods for treatment include the
10 use of the transporter protein or fragments.

Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof.
15 As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still
20 selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')₂, and
25 Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is
30 administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the transporter proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or transporter/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or

abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the transporter peptide to a binding partner such as a ligand or protein binding partner. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array.

Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a transporter peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the transporter peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide
5 sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence when such a
10 nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID
15 NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprises
20 several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic
25 sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity
30 modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a

protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

5 As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the transporter peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3'
10 sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form
15 DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the peptides
20 of the present invention as well as nucleic acid molecules that encode obvious variants of the transporter proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis
25 techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules
30 provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel transporter maps to public BAC AC034305, which is known to be located on human chromosome 17.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridation conditions are well known in the art.

Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. 55 SNPs, including 4 indels, have been identified in the gene encoding the transporter protein provided by the present invention and are given in Figure 3.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel transporter maps to public BAC AC034305, which is known to be located on human chromosome 17.

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Experimental data as provided in
5 FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver.

Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is
10 determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in transporter protein expression relative to normal results.

In vitro techniques for detection of mRNA include Northern hybridizations and *in situ*
15 hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a transporter protein, such as by measuring a level of a transporter-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a transporter gene
20 has been mutated. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate transporter nucleic acid expression.

25 The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the transporter gene, particularly biological and pathological processes that are mediated by the transporter in cells and tissues that express it. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. The method typically includes assaying the ability of the compound to modulate the expression of the
30 transporter nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired transporter nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the transporter nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for transporter nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the transporter protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of transporter gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of transporter mRNA in the presence of the candidate compound is compared to the level of expression of transporter mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate transporter nucleic acid expression in cells and tissues that express the transporter. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for transporter nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the transporter nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the transporter gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative

compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in transporter nucleic acid expression, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in transporter genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the transporter gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the transporter gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a transporter protein.

Individuals carrying mutations in the transporter gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides information on SNPs that have been identified in a gene encoding the transporter protein of the present invention.. 55 SNP variants were found, including 4 indels (indicated by a "-"). As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel transporter maps to public BAC AC034305, which is known to be located on human chromosome 17. Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point

mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a transporter gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

5 Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays
10 such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant transporter gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv.*
15 *Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and
20 Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective
25 amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship).
30 Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the transporter gene in an individual in order to select an appropriate compound or dosage regimen for treatment.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production

of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control transporter gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of transporter protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into transporter protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of transporter nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired transporter nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the transporter protein, such as ligand binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in transporter gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired transporter protein to treat the individual.

The invention also encompasses kits for detecting the presence of a transporter nucleic acid in a biological sample. Experimental data as provided in FIGURE 1 indicates expression in the fetal liver and spleen. Specifically, a virtual northern blot shows expression in fetal liver and spleen. In addition, PCR-based tissue screening panel indicates expression in human fetal liver. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting transporter nucleic acid in a biological sample; means for determining the amount of transporter nucleic acid in the sample; and means for comparing the amount of transporter nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect transporter protein mRNA or DNA.

Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee *et al.*, PCT application W095/11995 (Chee *et al.*), Lockhart, D. J. *et al.* (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. *et al.* (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown *et al.*, US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al.*) which is incorporated herein in its entirety by

reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the transporter proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the transporter gene of the present invention.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed

herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and*
5 *Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed.
10 Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close
15 confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in
20 separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will
25 include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified transporter gene of the present invention can be routinely identified using the sequence information disclosed herein can be
30 readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a
5 plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules
10 when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the
15 vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may
20 be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not
25 limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers.
30 Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include

initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 5 (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, 10 papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold 15 Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are 20 well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme 25 digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and 30 CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for

affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterotransporter. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

5 Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as transporters, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

10 Where the peptide is not secreted into the medium, which is typically the case with transporters, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity
15 chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may
20 include an initial modified methionine in some cases as a result of a host-mediated process.

Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a transporter protein or peptide that can be further purified to
25 produce desired amounts of transporter protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the transporter protein or transporter protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native transporter protein is useful for assaying
30 compounds that stimulate or inhibit transporter protein function.

Host cells are also useful for identifying transporter protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant transporter protein

(for example, stimulating or inhibiting function) which may not be indicated by their effect on the native transporter protein.

Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a transporter protein and identifying and evaluating modulators of transporter protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the transporter protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the transporter protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al.* *PNAS* 89:6232-6236 (1992). Another example of a

recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 251:1351-1355 (1991). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect ligand binding, transporter protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* transporter protein function, including ligand interaction, the effect of specific mutant transporter proteins on transporter protein function and ligand interaction, and the effect of chimeric transporter proteins. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more transporter protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:
 - (a) an amino acid sequence shown in SEQ ID NO:2;
 - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
 - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3; and
 - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:
 - (a) an amino acid sequence shown in SEQ ID NO:2;
 - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
 - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3; and
 - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
3. An isolated antibody that selectively binds to a peptide of claim 2.

4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

6. A gene chip comprising a nucleic acid molecule of claim 5.

7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.

8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.
14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.
15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.
16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.

17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.

18. A method for treating a disease or condition mediated by a human transporter protein, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.

19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.

20. An isolated human transporter peptide having an amino acid sequence that shares at least 70% homology with an amino acid sequence shown in SEQ ID NO:2.

21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence shown in SEQ ID NO:2.

22. An isolated nucleic acid molecule encoding a human transporter peptide, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 or 3.

23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 or 3.

```

1  GTCTCCCTCC CGCGCGATGG CCTCGGCGCT GAGCTATGTC TCCAAGTTCA
51 AGTCCCTTCGT GATCTTGTTT GTCACCCCGC TCCTGCTGCT GCCACTCGTC
101 ATTCTGATGC CCGCCAAGTT TGTCAGGTGT GCCTACGTCA TCATCCTCAT
151 GGCCATTTC TGGTGCACAG AAGTCATCCC TCTGGCTGTC ACCTCTCTCA
201 TGCCTGTCTT GCTTTTCCCA CTCTTCCAGA TTCTGGACTC CAGGCAGGTG
251 TGTGTCCAGT ACATGAAGGA CACCAACATG CTGTTCTTGG GCGGCCTCAT
301 CGTGGCCGTG GCTGTGGAGC GCTGGAACCT GCACAAGAGG ATCGCCCTGC
351 GCACGCTCCT CTGGGTGGGG GCCAAGCCTG CACGGCTGAT GCTGGGCTTC
401 ATGGGCGTCA CAGCCCTCCT GTCCATGTGG ATCAGTAACA TGGCAACCAC
451 GGCCATGATG GTGCCCATCG TGGAGGCCAT ATTGCAGCAG ATGGAAGCCA
501 CAAGCGCAGC CACCGAGGCC GGCCTGGAGC TGGTGGACAA GGGCAAGGCC
551 AAGGAGCTGC CAGGGAGTCA AGTGATTTT GAAGGCCCA CTCTGGGGCA
601 GCAGGAAGAC CAAGAGCGGA AGAGGTTGTG TAAGGCCATG ACCCTGTGCA
651 TCTGTACGCC GGCCAGCATC GGGGGCACCG CCACCTGAC CGGGACGGGA
701 CCCAAGCTGG TGCTCCTGGG CCAGATGAAC GAGTTGTTTC CTGACAGCAA
751 GGAACCTCGT AACTTTGCTT CCTGGTTTGC ATTTGCTTTT CCAACATGC
801 TGGTGATGCT GCTGTTCGCC TGGCTGTGGC TCCAGTTTGT TTACATGAGA
851 TTCAATTTTA AAAAGTCCTG GGGTGCGGG CTAGAGAGCA AGAAAAACGA
901 GAAGGCTGCC CTCAAGGTGC TGCAGGAGGA GTACCGGAAG CTGGGGCCCT
951 TGTCTTTCGC GGAGATCAAC GTGCTGATCT GCTTCTTCTT GCTGGTTCATC
1001 CTGTGGTTCT CCGGAGACCC CGGCTTCATG CCCGGCTGGC TGACTGTGTG
1051 CTGGGTGGAG GGTGAGACAA AGTATGTCTC CGATGCCACT GTGGCCATCT
1101 TTGTGGCCAC CTGCTATTTC ATTGTGCTTT CACAGAAGCC CAAGTTTAAC
1151 TTCCCGAGCC AGACTGAGGA AGAAAGGAAA ACTCCATTTT ATCCCCCTCC
1201 CCTGTGAGT TGGAAAGTAA CCCAGGAGAA AGTGCCCTGG GGCATCGTGC
1251 TGCTACTAGG GGGCGGATT TCTCTGGCTA AAGGATCCGA GGCTCGGGG
1301 CTGTCCGTGT GGATGGGGAA GCAGATGGAG CCCTTGCACG CAGTGCCCCC
1351 GGCAGCCATC ACCTTGATCT TGTCTTGTCT CGTTGCCGTG TTCACTGAGT
1401 GCACAAGCAA CGTGGCCACC ACCACCTTGT TCCTGCCCAT CTTTGCCTCC
1451 ATGTCTCGCT CCATCGGCCT CAATCCGCTG TACATCATGC TGCCCTGTAC
1501 CCTGAGTGCC TCCTTTGCC T CATGTTGCC TGTGGCCACC CCTCCAAATG
1551 CCATCGTGT CACCTATGGG CACCTCAAGG TTGCTGACAT GGTGAAAACA
1601 GGAGTCATAA TGAACATAAT TGGAGTCTTC TGTGTGTTT TGGCTGTCAA
1651 CACCTGGGGA CGGGCCATAT TTGACTTGGA TCATTTCCTT GACTGGGCTA
1701 ATGTGACACA TATTGAGACT TAGGAAGAGC CACAAGACCA CACACACAGC
1751 CCTTACCCTC CTCAGGACTA CCGAACCTTC TGGCACACCT TGTACAGAGT
1801 TTTGGGGTTC ACACCCCAA ATGACCCAAC GATGTCCACA CACCACCAA
1851 ACCAGCCAA TGGGCCACCT CTTCTCCAA GCCCAGATGC AGAGATGGTC
1901 ATGGGCAGCT GGAGGGTAGG CTCAGAAATG AAGGGAACCC CTCAGTGGGC
1951 TGCTGGACCC ATCTTTCCCA AGCCTTGCCA TTATCTCTGT GAGGGAGGCC
2001 AGGTAGCCGA GGGATCAGGA TGCAGGCTGC TGTACCCGCT CTGCCTCAAG
2051 CATCCCCAC ACAGGGCTCT GGTTTTCACT CGCTTCGTCC TAGATAGTTT
2101 AAATGGGAAT CAGATCCCTT GGTGAGAGC TAAGACAACC ACCTACCAGT
2151 GCCCATGTCC CTCCAGCTC ACCTTGAGCA GCCTCAGATC ATCTCTGTCA
2201 CTCTGGAAGG GACACCCAG CCA (SEQ ID NO:1)

```

FEATURES:

```

5'UTR:      1 - 16
Start Codon: 17
Stop Codon: 1721
3'UTR:      1724

```

FIGURE 1A

HOMOLOGOUS PROTEINS:

Top 10 BLAST Hits:

gi 2811122 gb AAB97879.1 (U87318) NaDC-2 [Xenopus laevis]	682	0.0
gi 4506979 ref NP_003975.1 solute carrier family 13 (sodium-de...	629	e-179
gi 3065814 gb AAC31165.1 (AF058714) sodium-dicarboxylate cotra...	626	e-178
gi 10280599 gb AAG15426.1 AF201903_1 (AF201903) Na/dicarboxylat...	624	e-178
gi 2499524 sp Q28615 NDC1_RABBIT RENAL SODIUM/DICARBOXYLATE COTR...	602	e-171
gi 8132324 gb AAF73251.1 AF154121_1 (AF154121) sodium-dependent...	543	e-153
gi 4322346 gb AAD16019.1 (AF081825) sodium-dependent high-affi...	531	e-149
gi 5531902 gb AAD44522.1 AF102261_1 (AF102261) sodium-dicarboxy...	515	e-145
gi 10439272 dbj BAB15477.1 (AK026413) unnamed protein product ...	490	e-137
gi 2499526 sp Q07782 NASU_RAT SODIUM/SULFATE COTRANSPORTER (NA(...	489	e-137
gi 9507109 ref NP_062354.1 solute carrier family 13 (sodium/su...	486	e-136
gi 6912690 ref NP_036582.1 sulfate transporter 1 >gi 6224691 g...	483	e-135
gi 2499525 sp P70545 NDC2_RAT INTESTINAL SODIUM/DICARBOXYLATE C...	437	e-121
gi 6226757 sp Q93655 YV06_CAEEL HYPOTHETICAL 66.2 KDA PROTEIN F...	400	e-110
gi 630683 pir S43561 YCR37C homolog K08E5.2 - Caenorhabditis e...	390	e-107

EST:

gi 751038 /dataset=dbest /taxon=9606 /...	519	e-145
gi 2658836 /dataset=dbest /taxon=9606 ...	416	e-114

EXPRESSION INFORMATION FOR MODULATORY USE

gi|751038 Human fetal liver spleen
gi|2658836 Human fetal liver spleen

Tissue Expression: Human fetal liver

FIGURE 1B

```

1 MASALSYVSK FKSEVILEVT PLLLLPLVIL MPAKFVRCAY VIILMAIYWC
51 TEVIPLAVTS LMPVLLFPLF QILDSRQVCV QYMKDTNMLF LGGLIVAVAV
101 ERWNLHKRIA LRTLLWVGAK PARMLLGFMG VTALLSMWIS NMATTAMMVP
151 IVEAILQOME ATSAATEAGL ELVDKKGAK LPQSQVIFEG PTLGQQEDQE
201 RKRLCKAMTL CICYAASIGG TATLTGTGPN VVLLGQMNEL FPDSDKLVNF
251 ASWFAFAPFN MLVMLLFAWL WLQFVYMRFN FKKSWSGCGLE SKKNEKAALK
301 VLQEEYRKLK PLSFAEINVL ICFFLLVILW FSRDPGFMPG WLTVAWVEGE
351 TKYVSDATVA IFVATLLFIV LSQKPKFNFR SQTEERKTP FYPPPLLDWK
401 VTQEKVPWGI VLLGGGFAL AKGSEASGLS VWMGKQMEPL HAVEPPAAIL
451 ILSLLVAVFT ECTSNVATTT LFLPIFASMS RSIGLNPLYI MLPCTLSASF
501 AFMLPVATPP NAIVFTYGHK KVADMVKTGV IMNIIGVFCV FLAVNTWGRA
551 IFDLDFPDW ANVTHIET (SEQ ID NO:2)

```

FEATURES:**Functional domains and key regions:**

[1] PDOC00001 PS00001 ASN_GLYCOSYLATIONN-glycosylation site

Number of matches: 2

```

1 194-197 NSSL
2 607-610 NVTH

```

-----[2] PDOC00005
PS00005 PKC_PHOSPHO_SITEProtein kinase C phosphorylation site

Number of matches: 3

```

1 222-224 THR
2 336-338 SKK
3 417-419 SQK

```

-----[3] PDOC00006
PS00006 CK2_PHOSPHO_SITECasein kinase II phosphorylation site

Number of matches: 5

```

1 222-225 THRE
2 358-361 SFAE
3 426-429 SQTE
4 428-431 TEEE
5 609-612 THIE

```

-----[4] PDOC00008
PS00008 MYRISTYLN-myristoylation site

Number of matches: 7

```

1 93-98 GLIVAV
2 118-123 GAKPAR
3 264-269 GGTATL
4 271-276 GTGPNV
5 460-465 GGGFAL
6 468-473 GSEASG
7 574-579 GVIMNI

```

-----[5] PDOC00978
PS01271 NA_SULFATESodium:sulfate symporter family signature

543-559 ASFAFMLPVATPPNAIV

FIGURE 2A

Membrane spanning structure and domains:

Helix	Begin	End	Score	Certainty
1	19	39	2.318	Certain
2	59	79	1.932	Certain
3	89	109	0.776	Putative
4	130	150	1.930	Certain
5	213	233	1.646	Certain
6	258	278	2.102	Certain
7	318	338	1.743	Certain
8	359	379	1.781	Certain
9	411	431	1.337	Certain
10	448	468	1.879	Certain
11	493	513	1.996	Certain
12	534	554	1.781	Certain

BLAST Alignment to Top Hit:

Alignment to top blast hit:

>gi|2811122|gb|AAB97879.1| (U87318) NaDC-2 [Xenopus laevis]
Length = 622

Score = 682 bits (1741), Expect = 0.0

Identities = 332/619 (53%), Positives = 439/619 (70%), Gaps = 55/619 (8%)

Query: 1 MASALSYVSKFKSFVILFVTPLLLLPLVILMPAKFVRCAYVIILMAIYWCTEVIPLAVTS 60
M S ++ +++ I+F+ PL LLPL +++P K C +VII+MA++WCTE +PLAVT+
Sbjct: 1 MVSIGKWILANRNYFIIFLVPLFLLPLPLVVPTKEASC GFVVIIMALFWCTEALPLAVTA 60

Query: 61 LMFVLLFPLFQILD SRQVCVQYMKDTNMLFGLGLIVAVAVERNLHKRIALRTLWVGAK 120
L PVLLFP+ I+DS VC QY+KDTNMLF+GGL+VA++VE+WNLHKRIALR LL VG K
Sbjct: 61 LFPVLLFPMMGIMDSTAVCSQYLKDTNMLFIGGLLVAISVEKWNLHKRIALRVLLIVGVK 120

Query: 121 PARMLGFMGVTALLSMWISNMATTAMMVPIVEAILQOMEA----- 161
PA L+LGFM VTA LSMWISN ATTAMM+PI +A+++Q+ +
Sbjct: 121 PALLLLGFMVVTAFLSMWISNTATTAMMPIAQAVMEQLHSSEKVDERVEGNSNTQKNV 180

Query: 162 -----TSAATEAGLELV DKGKAKELPGSQ-----VIFEGP 191
T A G E+ +K P Q ++ E
Sbjct: 181 NGMENDMYESVMPSGKMALADNTYATENEGFEIQEKSTKDPEPSKQEKQSIGPIVIEPE 240

Query: 192 TLGQQEDQERKR---LCKAMTLCICYAASIGGTATLTGTGPNVLLGQMNELFPDSKDLV 248
Q E++++++ +CK M+LC+CY+ASIGG ATLTGT PN+V+ GQM+ELFP++ +++
Sbjct: 241 DEKQTEEKQKEKHLKICKGMSLCVCYSASIGGIATLTGTPNLVMKGQMDLFPENNNII 300

Query: 249 NFASWFAFAFPNMLVMLLFAWLWLFQVYMRFNFKKSWGCG--LESKKNEKAALKVLQEEY 306
NFASWF FAFP MLV+L +WLWLQF+Y+ NFKK++GCG E K+ EK A +V+ E+
Sbjct: 301 NFASWFGFAFPMTMLVLLALSWLWLFQFIYLGVNFKKNFGCGGNAEQKEKEKRAFRVISGEH 360

Query: 307 RKLGPLSF AEINVLICFFLLVILWFSRDPGFMPGWLTVAVVEGETKYVSDATVAIFVATL 366
+KLG ++FAEI+VL+ F LLV+LWF+R+PGFM PGW T+++ +G + V+DATVAIFV+ +
Sbjct: 361 KKLGSMTFAEISVLVLFILVLLWFTREPGFM PGWATISFNKGGKEMVTDATVAIFVSLM 420

Query: 367 LFIVLSQKPKFNFRSQTEEEERKTPF-YPPPLLDWKVTQEKVPWGIVLLGGGFALAKGSE 425
+F S+ P F ++ + K PP LLDWK EK+PW IV+LLGGGFALAKGSE
Sbjct: 421 MFFFPSELPSEKYQDTPKPGMKPKLRVPPALLDWKT VNEKMPWNIVILLGGGFALAKGSE 480

Query: 426 ASGLSVMMGKQMEPLHAVPPAAITLILSLLVAVFTECTSNVATTTFLPIFASMSRSIGL 485
SGLS+W+G+++ PL ++PPAAI LIL LLVA FTECTSNVATTTFLPI ASM+++I L
Sbjct: 481 ESGLSLWLGEKLTPLQSIPPAAIALILCLLVATFTECTSNVATTTFLPIILASMAKAIQL 540

Query: 486 NPLYIMLPCTLSASFAMLPVATPPNAIVFTYGHKLVADMVKTGVIMNIIGVFCVFLAVN 545
NPLYIMLPCTLSAS AFMLPVATPPNAI F+YG LKV DM K G+++NI+GV + LA+N
Sbjct: 541 NPLYIMLPCTLSASLAFMLPVATPPNAIAFSYGLKVIDMAKAGLLNILGVLITITLAIN 600

Query: 546 TWGRAIFDL DHPDWANVT 564
+WG +F+L FP WAN T
Sbjct: 601 SWGFYMFNLGTFPSWANAT 619 (SEQ ID NO:4)

Hammer search results (Pfam):

No match

FIGURE 2B

```

1  TTCAACCATT GTGGAAGACA CTGTGGCGAT TCCTCAAGGA TCTAGAACCA
51  GAAATATCAT TTGACCCAGC AATTTTATTA CTGGGTATAT ACCCAAAGGA
101 TTATAAATCA TGCTGCTATA AAGACACATG CACACTATTT ACAATAGCAA
151 AGACTTAAAA CCAACCCAAA TGTCCATCAA TGATAGACTG GATAAAGAAA
201 ATGTGGCACA TACATACCAT GGAATACTAT GCAGCCATTA AAAATAATGA
251 GGTCTATGTC TTTGCAGGGA CATGGATGAA GCTGGAAGCC ATCATTCTCA
301 GCAAACATAAC ACAGGAACAG AAAACCAAAC ACCACATGTT CTCAGTCATA
351 AGTGGGAGTT GAACAGTGAG AACGCATTGA CACAGGGAGG GGAACATCAC
401 ACACGGGGGC CTGTCAGGGG GTTGGAGGGC AAGGGGAGGG AGAGCATTAG
451 GACAAATACC TAATGCATGT GGGTCTTAAA ACCTAAATGT CCGGTTGATA
501 GCTGCAGCAA ACCACCATGG CACATGTATA CCTATGTAAC AAACCTGCAC
551 ATTCTGCACA TGTATCCCAG AACTTAAAGT AAAATTAAAA AAAAAGAAAA
601 GAAAAAGAA CTGAAGTTGT TTAATTGCTC TCATTTCATG ATCCCGGAGA
651 AAAAGGTTTG AGTGCACATC CTGGATTAGG CACTGAGAAA GGCACTAGCT
701 GGACAGGTGG TGATGAATAA AACAGACAGT AAATAGAAAT TACATCATAA
751 TAATGTGTCA TATATTTTAA AATAGCTACA AGATATTTTA AATGTTCTCA
801 CCACAAAGAA ATGACAAATA TTTGGGCCAG ACGCGGTGGC TCACGCCTGT
851 AATCCAGCA CTTTGGGAGA CCGAGGTGGG CGGATCACCT GAGGTGAGGA
901 GTTCGAGGAC AGCCTGGCTA ACATGGTGAA ACCCCATTTC TACTAAAAAT
951 GCAAAAAATT AGCCGGGCGT GGTGGTGCAC ACCTGTAATC CCAGCTACTT
1001 GGGAGGCTGA AGCAGGAGAT TTGCTTGAA CTAGGTGGCA GAGGTTGCAG
1051 TGAGCCGAGA TCGTGCCACT GCACTCCAGC CTGGGTGACA GGAGCAACAAC
1101 TCTGTCTCAA ACAAACAAAC AAAAAACAAA AACAGAGAA ATGATAAATA
1151 TTCGAGTGAT AAATATGCTC ATTAGCCTGA TTTGAACACA CCACAATTAT
1201 ACACACATTG AAAAATCACA TGGTACCCCG TAAATATAGA CAATGATTTG
1251 TCAATTAAAA ATGAAATAAC ACTTAAAAAA TAAAAAAGTA AAAAGTAAAA
1301 ATTACACCAA TAAATATAAG AGGTACAAAT TGTGCTAAGT GCCCTGGGGA
1351 CACAGGAAG GCGGGAAGAC CCAGGGCTAT ATGCATGAGA GTTACAAAGG
1401 GAAAAGGACA GGAGGGAGGC AATTGCAGGA GGGGCTTGGG AGAATGCATG
1451 TCCTTGGGTG CAGGTTCACAG GAAGGAATC ATGAGCTTGA TTCAGGATGT
1501 GTTGAAATTT CGGGCCGAGA CACGTCCAGT CTGCGGAAGG CTGGACATCT
1551 GGGACTCTGG CATCATGGCT GGGTTGAAGG CAGAGGATGG TAATCACTAA
1601 GGAGCGCGCT GTGGTTAGGC CACCAGCATG GATGAGACTC CCCAAAGGA
1651 AGCTCGAGAA TGAGAGGCAG GCAGAGGAGA GGAAGAAGA AAATCACAGA
1701 GGTGGGGATG TCTTTGCATC CGTGTGTCTC CAGTGCCCAA AACAGGGCCT
1751 CCGGAGAAAG AGGTGCTCGG CACCTGTCTG TTGCTGGGCG GGCTGAATGA
1801 ATACATGGGC GACTGTCTCA GTGTGCGCTT AGTTGTGTCC CTTCTCTCT
1851 AGAGCTCCGT TTCCCTCTGA CCTGGGTCGG GCGGGCAGCT GCGGCTGCTG
1901 AGGCTCGGTG GGGCCCTCC AAGACGCGTG TCCGCATCTG CCCGCCGGG
1951 GTCTGCGGGG TGACAGCTCC ACTGGAGCGC GACAGCCCTT GGGACAGAGG
2001 AGGACAGTGG CTTGCTTCC CTGTGCGATC GCCCAGGAGC TCCGGGCCGG
2051 AGAGTGCGAG CGGGGAAAAG GGGTCTCTGA CCTAGAGTGG GCGGACGCTG
2101 GCGAGGAAGC CAGGGGGGAC CGGGAAGCGA GGCCCGCGGT GCGGAGGGCG
2151 CGGGGCGTGG GGGGACACCT CTCGGAGAGA CACCGGAGGG GCGGAAGTAA
2201 GGAGATGGAA AGGAGAGGGA GATCGGGGAG ATAGACCTGA GAGACCCAGA
2251 GGCCTGCAGA GAGTTTCATC CGGACCCCTT CAGAGCCCAG GAAAGAGCAG
2301 ATGCGGAGCG GGGAGGGCGC CTTACGCCAA AGCGGGCAGC ACCAGTGACC
2351 AAAACACGCC CCGCTTGCCA GCCCGGGGAC GCACCTCTGC CTCGGCAGCG
2401 CAGGAGAGGC TTGGACAGCG CGAGATGCTA GGGCCAGGCT TGCCCCTAGA
2451 GGGCTGGCCC GAAGCGTTGG AGTCCAAAGA CGCCTCCAC CGCCGCCGGG
2501 TGGCAGAATT GGGGGCAGGC GCGTCCACA GACCCCGAGG GGTGGCCCCG
2551 CCCCAGGGCC GCGGGGAGGC GCCCCGCTGC GGGGCGGAGT GTGCACCGCC
2601 CCCTGCCCAA TCCCCGGGGA CTGTGGCCCC TTCTTAAGCC CGCGGCGCCT
2651 CTAGCTGCCC CTCACTCGTC TCGCCGCCCA GTCTCCCTCC CGCGCGATGG
2701 CCTCGGCGCT GAGCTATGTC TCCAAGTTCA AGTCCTTCGT GATCTTGTTC
2751 GTCACCCGCG TCCTGCTGCT GCCACTCGTC ATTCTGATGC CCGCCAAGGT
2801 CAGTTGCATC TCCAGGCAGC CCTTCGGACA CCCGCGCTCC TGTGCCACT
2851 AACGGGCACC GATCCCGGGA GCCCTGAGCT GGAGCGCACG GATTTCGCGG
2901 GGAGCACAGC TCTCCCGGGG CGCGCGCACT CAGGAGCTCC AGGTGCGGGA
2951 TGGGAGGTGC CCTGTAAAGA ATCTGAGGGG CATGGCGACC CCAGGGCGCA
3001 CCACCTTTGG GGTTTACAGA TCCCAGGGCG CAAGAGCCGT CCAGGCAAGC
3051 ACGGAAACCT CGAAGTGAGC ACAGATCTCA GCCACACAGA TCCCAGCCTT
3101 AGGCTCAGCC CCTGGCTCCG AATCGAATCT CCCACAGTGC ATAACCCTGT
3151 TTCCCCCCAA AATGCCACCT GCGCCAACAG GGAACCTGGG AGCTTGCCCT
3201 TCCCTCTCTC TCTCTGTCT TTTCCCTTCG CCAAAGAAGA CTTCAGCTG
3251 TAGGTGGCTT CTGCCGTCAG GAGGGACCTA CAGGAAAAAA ATCATCACCC
3301 ACGTGGATCC TGCGCTGTCT TTGCCACTCT CTGGCCCTTC CTGGGCGCTT
3351 AGTGTCTCTA TCTATGATCC ACATTCTCTC CAACCTGGAG AGCCACATCT
3401 GATTTCATAAT CCTGCTCCAG GCTGCAGGCA GGTGGGGT GTGCTGCTTT
3451 GCCTCTGCCC TGCTGGGCT GTAGCAGGAG GCGGACACA TTCCCAGAGC
3501 TCGGACGCTT GGGTGGCAGG ACCTGGAGTT GCAGGGAAGC TTCTCCAG
3551 GCCCTAGTCT CTAATGCTT CTGTGAGGGA GAGAGAGAAT GAATGCGCTT
3601 GGCCGAGGG TGGGCGCAGG CTCCACTGGG CTGTGCACAG CCAGTTTGGC
3651 GGAGGCCCAA GCCCTTTGAA GCCTTTGTG GCTGCTGGCT GCTCCTCTCT
3701 CGTTCTCTCT TTCAGCCCTT TCACTCTCAG CCCAGACAGG AAACCTCCAG

```

FIGURE 3A

```

3751 CTCCCCACCT CCCCTCCCCA GGCAGGTTTG GGAAACAGAG GAGCTCTTCA
3801 GGGGATGCTC TGGGGGGGAG CTCTAGAGGA AGGGAGTGCA CTGGGGTGTC
3851 AGGAAGCCAA CCTGCAAGAG AACTGGACTT CCACCATTCT GTCATCTGTG
3901 TGACTTTAGC CAAGTGCCTG TGCTCTCTGG GCCTTGGTGT TCTCATCTGT
3951 ACAATGGGGC TAGGTGGTTT GGTCTGACA CTTTAAGGCT TTGGGAATCA
4001 AAAGGGATTA ACGTTAGTTC CTAGGATGGG GGAGGGGAG ACTGGGAGGA
4051 GCCCTTGGGT GGGCCTAGCA CAGGCCCTGG ATGGGTCAAG GACAGCAGAT
4101 CCAACTGTGG AGGCTGTGCA GTTGCCACAC CAACTGTGGG CAGCCACACT
4151 TCCTCTGTAG CAAGAAGTCT GGGGTGTGTA TTGTCCAGGG GAAGTAGCCA
4201 GGAGGAGAG CTGCGATTTC AGCTCTGCTG GTAGGGAGTG ATGTTCCCTG
4251 GAATGATTAT AGTAGCTTGG CTGACCTTCC TGCCACAGGA GACCCCACTC
4301 ACACACACAC ACATACACAC TCTCTCTCTC ACACACACAC ACTCACAAC
4351 ACACACAGAC ACACACACAA ACACACAGAC ACACACAAAA CACACGCACA
4401 AACACAAACA CACACAAAAA CACACGCACA AACACACACC CAAACACACA
4451 AACACAGACA CACAAACACA CACACAAACA CACACACAAA ACACACAAAA
4501 AACACAAACA CACACATACA CATACACACA CACACACACA CACACCCTGA
4551 ACTGGAAGAC CTAACCTCAGT GTGTGTGTAT GTGTGTGTAT GTGTGTGTGT
4601 GTAAGAGAGA GAGAGAGAGA GATTAAGCTG TCCTTTGAGT GAGGACCAGG
4651 GAGGGGAAGA AGAGAACCCA GGGAGAGTCC TTCCAAAGGC TGCCTTCACG
4701 AGCTTTCCTT CTGGCGGGGT TGGGTGAGGA CCCTGGACCT TGTCTTCTTG
4751 TTTTTCCTCT TTCTGCCTGT TTTGGTCACC CTGCCCCAC CCTCCATGGC
4801 CGCCCCATTG TGCAAGGAAA CCCAGAGGGT ACACAGCAGG GGCAGGGCAG
4851 CTGGGAAGCT GGTGAGAAGC TGGGAGGACC TTGGCAGCCT GAGCAACACA
4901 CTCTTTGCCA GGAGGTGACT CCCAGGGCAC GCCACCCTCT GCCAACACCC
4951 AGGCCTCTCT CCTCACCAGC TGTCTCCAGT TTTCTGTCTT CCACCTGGAT
5001 TCCCTCCTGG CCTCATCTCT GCTCCACTCT CTCTATCCTT CCTCTGGGTC
5051 TTTTTTTAAT TGAIAAAAAA TTTAATGAAA TAAATGATAG ATTTCTTGTA
5101 TCACCTTATT TATTAAATG TAAAGGTTT CTTTTTTGCA AATCTGTAAG
5151 ATATAAAGTA AAAATAAAAG TACACTCAA TCCCATAGT TATTCACATT
5201 TTGATGAACA TCTTCCAGA TGAATCTCTC TCTCTCTTCC CAGACACACA
5251 CACACACACA CACACACAGT AGGTTTTGCC TGCATTTTTT CATTAAGTGG
5301 TGTGTCAGGA CACCCTGCCT TGTAAATGTT GAACTTTTCT AACATCCGCT
5351 TCCCATCCTG CCTCTCCTCT TGACACTGTG GAGGCATTCT AGACTAGGGG
5401 GGTCAAGCCT GTTGACTTCA GGGATGAGGC ACCTCCTGGG CTCTAAATA
5451 GTGGCGCGGA GGTGAGGGGG CAGTTAACCT TGTGTCTCGT CCTCTTCTCT
5501 AGTGGGTCTG CTTGACTCCT CCAGGAACGC ACAGTGATCA TTGGTGACGC
5551 AGCCACACCTA CTGCTTCTAA GTTTAGAGAA TCAAAAGTTA CCGAGGACTT
5601 TGTGCGCCAT ATGGGAAGAA TGAGCACTCT TAAATCCACG ATTTGCAGAT
5651 GAAGACATGA AACAAGAGGG GACAGGGACC AGGATTGGGA GCAGGAGGAG
5701 TAATTTATGA GCGACATTGT TTAGAATTGC TATCACTTGA TGATAGTAAG
5751 AAGCAAACCTA ATTTTTCAGT AATATTATTG TTTTAAATTT CTCTCCAATG
5801 CGCCCTCTCA TTGTCTGCCC CTGGAGGCAT CATTCTGATG GCCTGCCCAG
5851 GGTACACCCC GACACCAAGC CCCAAGGAAG TTAGTGGCTG CCAAAGGCCA
5901 GACAGTGGCT GACAGTGGGC GCCAATCATA TCTGTCTGGT GTCAAAGCCT
5951 GGGCTTCCAG TCACGCTGCT GTTCCGCTTT TAGTTACAGC GCTGTCAACC
6001 AGCATTGACC CTTTCTCCTG CCCTCACCTT GCCCCACAAC AGGGGAACTT
6051 TGGCAAAATA CAGAGACATT TTTTGCTGTC CAACCTGGAG ACAGTCTTAC
6101 TGGCATCTCA TAGGTGGAGG CCAGCGGTGC TCTAAACACC CTGCAGTGCA
6151 CAGCTCCAC AACAAAGCAT CATTAGCCCC AAAATGTGAG TGTGCCGAGG
6201 CTGAGGGACC CTGCCCTCCA GTAGGGAGGT GCCCTGGTTT GCTCGTGGGA
6251 TGCTGAAAAA AGATTTATTT TTTTGTGGCT GATAACACAA CCCTGACAAA
6301 GAATTTCCAA GTCTTCTGCT ACTGTTTGTG GCAATAATAA CATACGCTCT
6351 TCTGGGTGAT GAGAAGCAGG GATTGTGTAC AGGTGCATCT GTTCTTCAGC
6401 AGCATTGTCA GAGTTAAACT CAGATGAATG CTATTGATTC CTTTAAATAA
6451 CATTGTCAAA GATGGCCGGG CACAGTGGCT CATGCCCTGA ATCCAGCAC
6501 TTTGGGAGGC CGAGACAGGT GGATCACGAG GCCAGGAGAT CAAGACCATC
6551 CTGGCCAACA TGGTGAAATC CCCTCTCTAC TAAAAATACA AAAATTAGCC
6601 GGGCGTGGTG GCGCTTGTCT GTAGTCCAG CTACTCAGGA GACTGAGGCA
6651 GGGAATCCG TTGAACCTGG GAGGTAGAGG CTGCAGTGAG CCAAGATTGC
6701 ACCACTGCAC TCCAGCCTGG GGACAGAGCA AGGCTCTGTC TCAAAAATAA
6751 GTAAGTAAGT AAATAAATAA ATAAATAAAT AAATAAGCAA GAATTTGCAA
6801 AGATATCCTA AGTGTTGGGC CTGTTCTGGA TGCTGAGGAC GGTGATCTAC
6851 AAATACAGCA GGTCTTGAA TAAATGTTGAT TCATTCAATA TCATTTTATT
6901 ATAATGTTGA TGAGGGAGGG AAAAAAAGG AAGGATCCCT TGAGCCCAGG
6951 AGATGGAGGT TACAGTGAGC TGTGACCGTG CCACTTCACT CCCACCTGGG
7001 CAACAGAGCC AGACCTGTCT TCAAAAAAAA AAAAAAGAG AATAAGAGAGC
7051 GAGAGAGAAA GAAAAGAAAA TGATTACTGG CTGGGGCCAC TGTCTGTGTG
7101 GAGCGTGAC ATTACCTCG TGTCCACATG GCTTTTCTTT GGCTAGTATG
7151 GTTTCCTTCC ACATCCCAA CCCGTGCACG TTAGTGAAAT TGGAGTGTCT
7201 GTATGGTCCC TGTCTGAGTG AGCGTGGGCG TCGGTGTGAG TGTGCTTCT
7251 GCAATGGGAT GGCATCTTGT CCAGGGCTGG TTTCCACCTT GTACCTGAG
7301 CTGCCCCGAG AGGATCTGGT CACCAAGAC CCTGACCTGC TGTAACCTGG
7351 TAAATAATTA TCTAATTTGT TTTCAATGTT TCTTAAGTAT ATGTATAGCT
7401 CACATTCCCT TCAGTGTGTA ATATTGGAAG TGTTTGTGTC TTTATTTAAG
7451 ATCTCCGTGA TGTTTTTGTG ACCAGAAATA TGCTGTAGAA ATTTAACTGT
7501 TGTTCCTAGC AATTTGCCTA TGGGAATATT GGCTTATGTT GTTTCGCTTA

```

FIGURE 3B

7551 CGCATTGCAA TTTCCAAAA CCAATCAATG ATGTTAAGTG AGGACTCACT
 7601 GTACTGTTTG TGCTTTCGAG TCACGCACTG GTTGTGGTGG TAGAAGGACA
 7651 GTTGAGGAAA CAGTGACAAC TCCATATGCT AATGGCTGGG GAGGGTACTC
 7701 AGAGGAAGGG CACAAACCAG ACTATAGAAG AGGCGCAGGG AGACATCTAA
 7751 GAAGGAATCT TGAGGTGGG CGCGGTGGCT CACGCCGTGA ATCCCAGCAC
 7801 TTTAGGGTGC TGAGGTGGGC GGATCATGAG GTCAGGAGTT CGAGATCAGC
 7851 CTGGCCAAATG TGGCAAAACC CCGTCTCTAC TAAAAATACA AAAATTAGCA
 7901 GGGCGTGGTG GCAGGTGCCT GCAATCCCAA CTCCGGAGGT TGAGGCAGGA
 7951 GAATCGCTTG AACCCGGGAG GTGGAGGTTG CAGTGAGCTG AGATTGTGCC
 8001 ATTGCACTCC AGCCTGGGCA ACAGGATCGA AACTCTGTCA CACACACACA
 8051 CACAAAAAAT ACTGATGAAA CATAAAACAA CCTAGGGAGG TGGCTAGTTT
 8101 TATCACATAA TTATTATTAC TTTTATTTC AATAGCTTTAG GGGTACACGT
 8151 AGTTTTCCGT TACATGAATG AATTGGATAG TGGTGAAGTC TGAGATTTTA
 8201 ATCCCTCCCT CCTATCCAC CCTGTCTGCT TCTAAGTCTC CAGTATCCAT
 8251 TCGACCACGC TATATACCTC TGGATACCCA TAGCTTAGCT CCCACTTATA
 8301 AGGGAGAACA TGCATAATT GGCTTTCCAT TGCTGAGTCA CTTCTCTTAG
 8351 AACAAATGCC TCCTAGGCGG CAAGAGCGAC ACTCCATCTC AAAAATAAAA
 8401 TAATAATAAA ACCAAAAAAA CCAGGTATTT TATTTCTCTT CTCCTTCTCC
 8451 TCTTCTCTCT TCTTTTCTT CTTTCTCCAT CCCCTTCTCT CTCCTTCTTC
 8501 TATCCCTCTC TCCTCTCTCT CCTCTTCTC CTTCTTCTCT CTTTCTTCTC
 8551 TTGTCTCTAT CTTGATCTTT TCTTTTGAGA GGCAGCTAAT CCAAGGTTTG
 8601 AGAAGATGAA AGAAGCTGCC TAGAACCACA CAGCTGGGAA GGAGGGAGGC
 8651 AGGGAGGAGG GGTGGGAATG GGGCAGGAGT CCTTTCGCAA TAGATCCCTG
 8701 GCCTGACCCG GGAAGCTGT GCTGACGAG GCTGGGGAAC AAGATGACTT
 8751 TGAGGGGAAT CCTCTGAGA TCAGCACTGT GTCTTGACAA TCCATGCCAG
 8801 CCGCCGTCCG GAGTGTCTG GGGGTGGGGA GAGGGAGGCG GCAACACGCT
 8851 GAGGCCTCAG GACTGTCTCT TCAGTTTGTG AGGTGTGCTT ACGCATCAT
 8901 CCTCATGGCC ATTACTGGT GCACAGAAGT CATCCCTCTG GCTGTACCTT
 8951 CTCTCATGCC TGTCTTGCTT TTCCCACTCT TCCAGATTCT GGACTCCAGG
 9001 CAGGTGAGCA GACCCAAGGG ATCCTGGTGA CTTTCTGGTT CTCCTCTTCT
 9051 CTCTTTCTCT AGTCCCACT GTGAGTCGCA CAGGCCTGGG GGTGACCGGA
 9101 AAACCTCAT TTGTGGATTC TCCCTGGCAG GGAGACACCA CTCGAGCCTG
 9151 CATCCCACT CCAAGCTGTC CTTGAAGTCA GCATCTGGGG ACTGGGTGGC
 9201 TCTAGTGTGT GGCAAGGGAC AGTCCTGATG AGGCCTTCGT GCCACGCTCC
 9251 AGGTGTGTGT CCAGTACATG AAGGACACCA ACATGCTGTT CCTGGGCGGC
 9301 CTCATCGTGG CCGTGGCTGT GGAGCGCTGG AACCTGCACA AGAGGATCGC
 9351 CTTGGGCAGG CTCTCTGGG TGGGGGCCAA GCCTGCACGG TAATTACGCC
 9401 TTCTCTCTCT TGCCACGTGG CTCTGCATGA GCCCCAGGGC TGGGAAGGGG
 9451 TGGAGGATGG CACAGACCAG GCCATCCACT GGTGAGGGCT GGCCATGGGC
 9501 TTACCTGGAC TTGGCTGGGT GGGGTGCACT TATAGCTTTA GTGGGAGAGA
 9551 CCAGATGCAT GCGTGGTGGT GGCACATGGT GAGCAGCAGT AAGTAAGGGT
 9601 CCTCGAATCC AGAGGAGGTG GGTGAGCAAG AGTCCTTGCA GGCTTGGAAG
 9651 GCTTTCTGGG GGAGGCACT AGCTGCAGGG TTCCACCGGG AACAAATTGG
 9701 ATGAGGGCTG GATCAAGCTG TGTCTGATAG GATAAGGGA GCAGGGCAGA
 9751 AGTGGCTCAA CTACCCAGCT CATGGGGAAG CAGAAAGGTC CTCTCTCCAA
 9801 GCTGGAGCAT CTATTTCCAC TGCAAAAGAG CTCTTATCT TCCCCGATAT
 9851 CACTCAGTAC CCCAGCTTCT CTCTCCATT CCAGGATCTC TCCTGCCAAT
 9901 CTAGCTAGCC ATTTCCAGCT AAGCCATGGA GTCAATATA TCATAATCAT
 9951 AACCAATAATC AATCATGATC ATAATGGGTA TATTGAGTGT CTACAAGGCC
 10001 CCAGGCATGA TACCAGGAGC TTATGAATTG CCTCATTTAA TTCTTACCAC
 10051 AACTCTGAGA GCTGAGTATT CTTACTGCC ACATTCTGTG GATGAGGGAT
 10101 TGGAGGCAGA GAGGGATAAA GTGATTGCT CATGGACACA CGGGGATTGG
 10151 ACCAGCTTC TGTGATAAGG CCTGTGCTCT CTCTAATCAG AAACCTCAGG
 10201 CATATCTTCC TTTTGAGACA ATGTGTCCCT TCAATGATGG CAGCTCCTTG
 10251 GCCAGCCAT CAGGAGTCAG CTGCTGGTCA GTTTACGGTA AATTCTCTCT
 10301 GAGGCGCCCC TGTGTAGGG GCTGTGCTAG ACCCTGAGCA CACACAGACG
 10351 CTAGCTGCC CTGCAAGCAA CCCATGCCCG GGGCTCCCTT GGTGCCTCAG
 10401 ATTTCCCTGA GCAGGGGACT TTGTGGCTCT CTGATCTGTT ACACATCCTG
 10451 GCATTGATTC CTTCAGTAA TTGCTTTAGC ATCAAATCAA AAGCCATCAT
 10501 ATTTTCTAGA AATGAGAGAC CCCAGGAAAG TGGACCTCAG GGCCCTCAGA
 10551 ATTTCTCTGC TTGGCTCCCT TGAGTGGCCA GCTTGGGTGG GAGGCCACTC
 10601 CAGTGGGTTT CATTCTGCAG CATGCTGGAG AGCTTCCACT TCCAAACCCA
 10651 AGTTCACACA TGCTTCTGTA TCCTTCTGCT CACCTTGCTC CTCTGAGTAT
 10701 GGTCTCCGGT TGTCCAAGGC ACTGCCTGTC CTGGGAGTCA CCTGTATGTG
 10751 AGGCACCTTG GGTGCCTTGA GATATCATGT AGAAGCCTTG GTTCTTCTCA
 10801 GACAACTCCA TTCATGCAAA CTCTCCCCCT CCTCCTAGCC TGGGTCCCGG
 10851 GCTTTGTTTT TTTTGGGTC CATAATGTCT GCCTGTGTGG ACAGCAGCTT
 10901 GGGCCCTGGT GCAGAACAGC TCCTAGGTCC CTTCTCAGG CTCTACCCC
 10951 TGCCCTGTCT CCTACCCCA GGTGAATTAG GAGCCCTGAG GAGGAGCCTG
 11001 GCTGCAGCGA GGCCACAGA CTGAGAGTAG CTGAGCTCCT TCTGTCCCTA
 11051 GCCTTGGACA GCTGGGCGAT GTAGAGCCAC AGAGCAGAGT CAGGCCCTGC
 11101 CCTGCTCACA GCGCAAGGAG AGAGCAGACA TGGAAACAGG TGCTTTGAAC
 11151 CCAGCACAGC GATGATTAGA GTAGGGGGAA GGATTGAGAA GGGTCAGGCC
 11201 AGCCCCACCT GGTGCACACA CTGAGAGCGT GGTCCAGAG GAGGGATGTT
 11251 GTTTGAGCAG GCTCTGAAG ACCATGAGGA GTCTTCTCTA TAGACAGCAG
 11301 AGAAGGGAGC AGGGGTACAA AGCAAAGGGA GTGTTTCTTC TGAATACTGT

FIGURE 3C

11351 TTGTGTGATA GCTCCACTGC AGCATGGAGG GGTCAAAGTG TATGTGCGGG
11401 GCGGAGGGGA GATGGCAGGG TTGGAGTGGC AGCCGGGAGA ATAGTCACAC
11451 TTTCCCAAGC TCCCTCCCCA GCTCACCCTA CCCCTACTCT GCTTAGCCCT
11501 TCTGAATTC TGAGAGGTGC AACAGAGTTT GGGGGTGGGT GGGAAATTTCC
11551 TAGCCAGAAG TGGGAAGCTG GGGCTGCCTG CACATAGGGG TATTCCAGCA
11601 CACCCTAGGG CAAGCTCATA TTGAGTTGGC ACCATCTGGA TGCCTGGGCT
11651 TCCCTTGCTA GATGGTGGGG CAGGGGTGCT CCTTAGAACC ACGACTGGAT
11701 CTGAGGCCTC TTGGTAACCC CAGAAGCAAG CAGAGTAGAC ATCAGTCATG
11751 GGTGTGGGAG AGGCAGGAGG GAGAGAGGAA TGGAGGAAGC AAAGAAGGGA
11801 AGGAGGGAGG GAGGGGAGGC TCTAAAACCG TCATCCCTAT TCCAATATCT
11851 GATCTTGAAT TGGCCTCAAC ACCTGTGCAT CCCTGCAGGG GTGGACCCAG
11901 TCCCCAGTTG CTTCCCAGGG AGTACGGGGG TGGGGTGGGG ATTCTCTGGC
11951 TTTCTTCCCT GCCCTCCTC TGCAGGCTGA TGCTGGGCTT CATGGGCGTC
12001 ACAGCCCTCC TGTCCATGTG GATCAGTAAC ACGGCAACCA CGGCCATGAT
12051 GGTGCCCATC GTGGAGGCCA TATTGCAGCA GATGGAAGCC ACAAGCGCAG
12101 CCACCAGGC CGGCTGGAG CTGGTGGACA AGGGCAAGGC CAAGGAGCTG
12151 CCAGGTGAGC CCCTGGCCAG GGCAC TGCCA GGGCACAACA GCAGCCTTCC
12201 CCTCCCTCTG CTGGCAAATG CTTTGGCCAC CTCCTTCTCC CTGTCTGCTT
12251 CCCGGAGCCC TCCTTTAAAC ACGCATAGAG AAAAAAAT AGAAAATACT
12301 GTTGTCTTAA GTTTTAGGAG GGGATTATTG CACACAATT AGATCCTTTA
12351 ATAGAGCTTT GAACAAAGTC TCACCCTCAG TTCCCATCAG TTGCAGAAAT
12401 CAGTGTGTTT ACCTGATTAT TCATTGGGGC ATCTTTCGAG CACTTAGGGA
12451 TGCCCTCTAC TCCTTGCTAC TCCTGCTCAT CCTCAAGGAG GCCTTTTCTG
12501 ACCTCCTCGA GCAGCTCAAA TCCTTCCACT CTCTGCTCCC ATAGTCTGCG
12551 GGCTTGCGGT CCCATGCTTG CTTCCCTGCT AGGTGCGAAG CTCAGGGAAG
12601 ACGAGTCAGC ATCTACCTTG CCGTCTGCGG TGTTCCTTA CCATCCCCAG
12651 CCGAGTCAG TAGAGTCAGG GTCTGTGGCT GACGGCCTGA TTGCCAGACC
12701 CTGGGCAAGG TCCTGGGGCT TACAGAGAGG AATCGGGCAC ATCCCTGCCA
12751 GCAACTCTTA TGGAGCCAG TGGGGCAGCT AAATCAGCAG AGCTGGGATT
12801 TCCCAATCCT CAGGTCAGCA GCAGAGTCAG GACCTGGGGC TGGGTGGGCA
12851 GCCCCATGA CTGGCTCAGC TAACAGCGCT GTGCCCCCA CAGGGAGTCA
12901 AGTGATTTTT GAAGGCCCA CTCTGGGGCA GCAGGAAGAC CAAGAGCGGA
12951 AGAGGTTGTG TAAGGCCATG ACCCTGTGCA TCTGTACGC GGCCAGCATC
13001 GGGGGCACCG CCACCCTGAC CGGGACGGGA CCCAACGTGG TGCTCTGGG
13051 CCAGATGAAC GAGTGAGTCC TTGGTGCAC CTCTGGGGA CAACGAAGTG
13101 GGTACCGGGG CTGGAGGGAC CTGCCACCT CTCTCTGCTC CTCTGCAGAG
13151 TCCTGGAAAG CCTCGGGGCA GCCAGACCTG GCCTGGGAGC CTGGCAGGGG
13201 TGGAAAGATG TGGCCCCATC TAGCCTCTGT GTCTTGGCAC CCCTGTGCCC
13251 ACACAGAGAG CTTAGAGAGG ATAGGGAGCT GATGTACAGG GAGCTAACGT
13301 CCCAGTCTG TTTCTGCTAT GATGCAAGAC CCACCACCTC CCCTGGGGTC
13351 AGGGACTCTG GCTCAGAGAG GGAGTGTGGA TTGAACCTCT AGCTAAAGTC
13401 ATGGCAGATG ACAATGTACT TCCAGACGCT GGTCTCTGG TTGAAACTTG
13451 TAGAAATAG ACACCTCTAA AAGACTCCCC AGCACTCCCT TTGCTCACTG
13501 CTTTGGTGG CTAATGGTGA TGGCCCCATG GCATCCGAGG TCTACAGATG
13551 GTATGAAGGG CTGGGGTTGG GTCATTCACT GCTTCACTGC TTCGTATAG
13601 TCCCTTGTG AGGTATCAGG TGAACCATGG GATGGTTTGG AACTTTCTAG
13651 CCTTGGCCAC AAAGGGATGC AGGCCATGAG GACCCCAAGA GGGAGAGAAA
13701 CCTTGGGCCCT GCCGCGGGT AGTCATGGTC TGTGAGGGT GGCAGATGC
13751 CTGGGGCTTC CAGGCATGTC TGGTACATAA ATGTACTAAT TGAGGTATGT
13801 ACTAATTGCA GTGGGCAGGC AAAAAATAA GGTGATGCCA TCCTTTGCAG
13851 ACAGGAGCCT GGACAGGGGT GGGGAGGGCA GTGGGCGCAG GAGCTGGGAG
13901 GTGGAAGGA CAGGTCTGGA GCCTGGCTGG GCAGAAACGT GAGGTTC AAC
13951 AACCCTGTTG TTTTAATTTT GGGAGTGTGT TCTGTAATGA TATCCTTACA
14001 GTTCTCCAGT AACTTTCTTT GGGAGAGCA GCCCGTCTGG GCTGAGTGGG
14051 GAAAGCTCTG CGCCTGCTTT GACACTCTTG AGCTAAAGGG GCGCCCTCTG
14101 GGGCTAGCAG AGCCCCGGGG ATGGGAGGCG GGGCCTGTGG TGGAGTGCAC
14151 CCTCTCCAG CCTCCGCTCT GGAAGCTTT TGAGATTTC TTTGCTAAGT
14201 GGGGGGACCG TTCTTTGCAG AAACCCACAG AGCGAGATTG CTGAGGTCTC
14251 TGCAGATCCC CAAAGATGTC AGCCAAATTA CATGCATGTG TATAAAGGT
14301 GTATTTTTCT TTTTTTCTT TTTGAGACAA GTCTCGCTCT GTCCGCCAGG
14351 CTGGAGTGCA GTGGCGCGAT GTTGGCTCAC TGCAACCTCT GCCTCCTGGG
14401 TTCAAGCGAT TCTCCCGCTT CAGCCTCCCT ATTAGCTGGG ATTACAGGCG
14451 CCGGCCACCA TGCTTGTTA ATTTTGTAT TTTTAGTGGA GACGGGGTTT
14501 CACCATGTTG GCCAGGCCAG TCTTAAGCTC CTGACCTGT GCGCCACCTG
14551 CCTCGGCTC CCAGAGTCTT GGAATTACAG GCGTGAGCCC CTGCGCCCGG
14601 CCACAAAGTT GTATTTTTCT GGAGGGATGG GCCATAACTT CCATGAGACT
14651 CTAGCAAGG CCTGGACACA CAGAAGAGTC AGTGGGTCAT TTCTCGGCCT
14701 TGTCTTGTG TGTGGCCATG TTCTGAGGCT CCCACTCGAT TAGGGGACAA
14751 TGCTTGGCAA TGGACTTGGT GGCTAGACCT CAGGAGGATG TGGCCTCCAC
14801 ACAGGCGCGC CTCTCAGGGC CCAGCTGCTG CTCCGTCCCC ACGCACAGGG
14851 CCAGGCTGGC TCCACAGCT CAGCATCTGA GGTGGGGGCC GGTGTCTTCT
14901 TGAGGTTGT TTTCTGACAG CAAGGACCTC GTGAACCTTG CTTCTGTGTT
14951 TGCATTTGCC TTTCCCAACA TGCTGGTGAT GCTGTGTGTT GCCTGGCTGT
15001 GGCTCCAGTT TGTTTACATG AGATTCAAGT AAGTTTGAGC TGCTCACAGC
15051 CTAATTATGC CTAAGAGCTG CAGAAGAGCC CTCAGACTCA
15101 ATAGGCAGGT TTACAAAGTC CTTCTGTCTT GGCCCTGATC TTTCTCCAGC

FIGURE 3D

```

15151 CCTGTCTCCT GCTAGTCTGC CCTCCTGTTT CTTGGAACCC AGGCTGCTCA
15201 CTGAGCTTTG TGCACACGTG GTCCCTTTTC CCTGGAATGC CATTCTCTAC
15251 CTTCCACCTT CCTCAGCCTT CAAGGCTAGT TCAAAATGCTG CTTCCCTGAC
15301 TTTTCCCCAC CCCCATTCCA TCTCTGAGCG GCCCCTGGGC ATATCACAGG
15351 CTTGTCTCTT AGTATCTGCA TTTGGCTTCC GGTGACTTTG AATTCTCTCA
15401 GAACCACTCT GATGCTGGGC ACCCCGCACA GCTCCAGCA CAGGGAGGAA
15451 GAGCAGGCAG GTTAAAGCAA TTAAGATAA GCTGGTCCCC ACGTGCCAGT
15501 TCGACATTGC TGGACAAGCT TCCTCTTTGC CGTGTGGGTC CATCAGGCCA
15551 GGTCAACGCA AACCTGTGAC TTAGCTCTGA GCTGAGCGCA TACGCTCTGT
15601 GCCTCAATGC ACGGGGAGTT TAAGTCGAGT AAAACCAGCA GTGATTATGA
15651 CCAATCCAT CCAAACCCAG ACATTTACTG AATACCTCTG GTGTTCCAG
15701 CAGTGACAG GTCCTAGAAA GTTTACCTTC CTGTTCTAG CACACAGGCA
15751 AGTTCATCAG GGGTCACCTT TGATGGCAGC CAGACTTTGG ACAGAAACCA
15801 TGACCTGTGG CTGACAAATA GCTAAAAAAA AGTTATTTGTT TTTCTAAAAAC
15851 ACACAAATTT ATCTGTGGTG CAAAGGTGAT CAGGCCACAC CAGGATAGAA
15901 AGTACTCAGC TCTGAGTTAA GTGCCTGTGC TCTGTGCCTC CATCCACAGG
15951 AAGTTCGAGC CAAGTCAAAC CAGGGGAATT TGTACCAGA GGGAGAGAC
16001 TGCAGAGCTC AGAGGCAAAA GTGCCACCGG AAACCTGTGA TTTTGTGGGG
16051 AAAATAGGGA ATTTCTCTAA GTTTCTTCT GAAGGAGGAA CTGTTTGA
16101 AACTCCCATT AAAAAGTTGC TATACAGGCC GGGCGCGATG GCTCACACCT
16151 GTAATCCCAA CATTTTTGGA GGCCGAGGTG GGCAGATCGC CTGAGGTGAG
16201 GAGTTTGTA CCAGCCTGGC CAACATGGTG AAACCCCGTC TCTACTAAAA
16251 ATACAAAAAT TAGCCGGGCG TGGTAGCCCA CGCCTGAAAT CCCAGCACTT
16301 TGGGAGGCCA AGGAGGGGCG ATCGCCTGAG GTCAGGAGCT CGAGACCAGC
16351 CTGGCCAACA TGGTGAAACC CCATCTCTAC TAAAAATACA AAAGTTAGCT
16401 GGGCATGGTG GCACATGCCT GTAACCCAG CTACTGGGA GGCTGAGGCA
16451 GGAGAATTGC TTGAAGCCGG GAGGTAGAGG TTGCAGTAAG CCAAGATCAT
16501 GCCACTGCAC TCCAGCCTGG GCGACAGAGC AAGACTCTGT CTCAAAACAA
16551 AAAAAAAGT TGCTATACAT ATTCAAAACA ATCATAATAA TGATAGTAAG
16601 AATGACAAAT TTAATGATCA TTGCCCAAAC CCCACTCTGT CTTGCCCATG
16651 GACGGGCGAG GGGAAACTGT TTGCATGGCT GCCTGGCCAC CCAGCCTGGC
16701 TTTGACAGTA GCTCTCTTTG CCCTGCCTCT TGAATCTGCA CCAGGGCCAA
16751 AGTCCCTGTT ATTGTTCAC ATCCGTGCGA CAGGTCTCTC AGGAGATGGT
16801 CCTGAACCTG CTGCAGGTGA GCATCTGTGT CTCCCTCATG GGCACAGGGA
16851 ATAATAATGA CCAACATTTA TTGAGTGCTC ATCATGTGCC AGACATGATT
16901 TCGAGCGCTC TTTTCTTTT TTTATTTTAT TTTATTTTAT TTTATTTTAT
16951 TATTTATTTA TTTATTTTAT TATTTATTTA TTTATTTTGG AGACAGTGTC
17001 TTGCTCTGTC ACCCATGCTG GAGTGCAGTG GTATGATCTC GGCTCGCTGC
17051 AACCTCCACC ACCTGGGTTT AAGCAATTCC CCCTGCCTCA GCCTCCCAAG
17101 TAGCTGGAAT TACAGGCACC CACCACCACC ATGCCTGGCT AATTTTGTGA
17151 TTTTTTAGTA GAGATGGGGT TTTGCCATGT TGGCCAGGCT GGTCTTGAAC
17201 TCCTAACCTC CGGTGATCCG CCCTCCTTGG CCTCCCAAAG TGCTGGCGTT
17251 ACAGATGTGA GCCACCTCGC CTGGCCCAAG CACTCTTAAA CTTAATTAAT
17301 TATTTATTTA ACCTGCGAGG TCAGCACTAT TATTATTATT CCAATTTTAC
17351 AGACAAGGCA ACTGAGGCAT GGAGAGGTGA TGTGGTCAAC ACAGAGCTTT
17401 GTAACAGGGA AGTAGGGGGA CTGAGACTTG AACCCAGGCC CTTTGGCTCC
17451 CACTGCATGG CATCCCTCTT TGGGAGGCT GAGGGTTGCT GTCCTTAGTT
17501 GCCTCCAGAC CTAAGCATGA CCAGGTGTCA GAAACACTAG TTGGGGCCGG
17551 GGCTGCCCTA GAACCCCAAG GCCTACTGAG AAAGAGGAGG GAGATAGCAT
17601 GGCGCCGAGG CCGCAAGGGC ACCATCAGCT TCTTGTCTGG CCAGAGGCAG
17651 ATGTGAGGCC CCTGGAGACT CACAGCCAGA ACCTGAAGCT GAGTCCACCC
17701 AGCCTGGCAC GGCCTTCATC AGCTTTTGTG GACTGGCGGG GGAGCCTGAG
17751 AGTGCTGCA GCAGGGGGCT TCTGAGCATG CTCGTGGTGG GGTGCTGGC
17801 TGCAGTCCAG TCCCACCCCT TCCCCTTCCC GACGGGCCAC CTTAGTTTGG
17851 ACGCATGCAG TGTGGCTGGC CGGGGTAGCT CACGGCAGCT TTGTTTTGGC
17901 TCCAGATCTG GAAGGTAGAG GACAGCTTTT ACATTCGGTT TGAGTGGTGG
17951 GAACAGTGCT CTGGCCAGG CCACGTCTCT CCACAACTA AGACCTGGTG
18001 GTCCCTGCCCT GCCTTTGTGG CCTCATGGAC CTCCTCCACT GAGGCCAGGG
18051 AGCACCTGTC TCAGCGGCAG GAGGCAGCTC CACTGTCAGC TGTGCTCTC
18101 ACTAGAGTTC CTCATCTGAA CGATCCTGGA GAACGAGGTT AAGTCTCTGG
18151 CCTCTAGCCT AATCCAGAAC AACTATCTTG CTGAAGAGCC TAGTGCAGCC
18201 TCCTAGGCTA TATCTAGCCA AAGGGGCCAG ACCCCACCCC AGGACCACCA
18251 AGAACTACAT GGGATATTAT TACTGGTTAT ACCTAACTGT CCCAACCAGG
18301 CTTACCTCCT GTAATAGCCA TGAGGGTTCT TTGGGACCCC TGCCAGGGCA
18351 GAGGCATGCA AAGCTCAAGA ATCTCTCCCC TCTTGTGTGG TCTGCAACAT
18401 ATTCAGTCCA AGTTCACCAT GGTGCATCAT GGTGAAGGCT GTTCTGCTGC
18451 AGGAGGACTC TGTGGTCCCC ACCCTGACC CTGACCTAGG CCCCTCACAG
18501 GCCAACTGGA TCCATTACT TGCATCTCAT GCCAGCCTGG TCATCACCAG
18551 ATGAAATTA CCCAGAGATG AGAGCAAAGC TGCTCAGCAC GAGAGACTCT
18601 GAAGGCTTGG CGGTACCACT GTGGGGCACT GGCATTGGAA GACTGCATAC
18651 TCCATGCAGC CCCAGAGTCT GCAGCTACTG TGGTGTGGG GATGAGCTGC
18701 CAGCACCAAA TGCAGGCTCT GGCTCCTGGG CCAGTAGTAA TACCAAGGTC
18751 ACCCTTATG CTGGAACCT GAAGCCCTG GCTGAGCCCC AGGGTCTCTA
18801 GGACGACAGT TGGCAGCAGA GAGGTGCTTG GTAGAGCACA AACTTTACTA
18851 AGCCAGGGT GTGGCAGCAG AGAGGCCCTG TCTTACACCA GCAGGCCAT
18901 CCCTGTGCGG GATGTCTAGA GAGTGTCCCT AGCGGGTGAC CCTCAGGACA

```

FIGURE 3E

```

18951 CACGGCCTTG CCCAGCAGGG AGATCCTAGC CAGCCGTGTA GACCTGAGGT
19001 CCCATCAGTG TTGCCCTCCTT TTCTGACCCC TGAGCACCCC AGAAAGCTGT
19051 GACCTGATGT CCTGGTGTCC CCATGTTCCA GGCCAAGCCA CCATCACACC
19101 AACACTTGGC CCTCACACTC TCCAAGGCTG TTCACATCCA GCACTGGCTT
19151 CAGGAATGAG CTCCTATTCC ATCAACCCCT TCCCTCCTAT GATTATGTCT
19201 CATGGCCCCC GGAAGGGCT CTCACGAGGG AGGGCTCTCC AGGACAATAC
19251 TCTTGGCCTT GCCCACCCTT TCAAACCAAC AGTGGCTGGA ACTGGAATGT
19301 GTGAATGGAA TATTCAGCAT ACCTTGAGGC CTTAGTCCTA TGCACAGTGG
19351 CCCCAGTTAT CCCCCTCCA CAGCTGAGCT CCCCTTTACA CCTCCTCAA
19401 GAACCTCCTC TCCTCCCTGC CTCCTCATGC CAACGCCACC TTAGGGGAGG
19451 CCGTGCAGGA CACCCTGGAC AATGGACACT GGTCCCAAGG GGGCCCATCC
19501 AGGATGGGGG TGCCATCCTG GGTGTCTTC CTTCTTGCCC TAGCCATGCT
19551 TGCTGCTAAC CCCAGGGTCT CCTGGATCCC TAATCCTGCA CCTCCAACCT
19601 CAGGGAACAC AAGGACCCAT TCTGCCCTG ACTAGCCCTG TCTGCCAGGG
19651 TTCATACFCA CTCCTGCGAT CTCCTGAGC CACCTTGGTG ATGGGGGTTG
19701 GCATCCCAAC ACCATCGAAG GCAGCTCCAG GCTGAGGTGG AAGGAGGAAG
19751 ACTTGGGAAG CATGTGAGGG AGCCCTGTTT CCACCTTGCG CAGGCTCCGA
19801 AGCTCCTTAT GGCTTCCCC CAGGTGACCC TGGAGCAGCC AGTCTCCAGG
19851 TGCTTGGGCA CCTGCCGAGA CCCTCTAGCC TCTCTACAGA GACTTTTTTC
19901 CTAGTACATT CTGGGATGGA AGAACAGGAG AGGGAAGAG GCAGGAAGGG
19951 CCTTTCTCCA GGCCCCATAG CAGGCGAGGA CAGCATTATG TGTCTTTTGT
20001 CTACATTCTG CTGTAGAACA TTTAGGCTCC ATCTGACCAG CACCTGAGCC
20051 AACCAGTCTG CCCTGCCCTT CTCTCATCTT TGCATTCTCC AGTTTTAAAA
20101 AGTCTTGGGG CTGCGGGCTA GAGAGCAAGA AAAACGAGAA GGCTGCCCTC
20151 AAGTGTCTGC AGGAGGAGTA CCGGAAGCTG GGGCCCTTGT CCTTCGCGGA
20201 GATCAACGTG CTGATCTGCT TCTTCTGCT GGTCTATCTG TGGTTCTCCC
20251 GAGACCCCGG CTTCTATGCC GGCTGGCTGA CTGTTGCCGT GGTGGAGGTT
20301 GAGACAAAGT AAGTCTTGGG TTCAATAGAA ATCGCTGGCT TAGGGCCAGG
20351 CGCTTGGCT CACACATGTA ATCCAGCAC TTTGGGAGGC TGAGGTGGGT
20401 GGGTCACCTG AGGTCAGGAG ATCGAGACCA TCCTGGCCAA CATGGTGAAA
20451 CCCTGTCTCT ACTAAAAATA CAGAAAATTA GCGAGGCATG GTGGCACATG
20501 CCTGTAGTCC CAGCTACTTG GGAGACTGAG GCAGGAGAAT CACTTGAACC
20551 CAGGAGGCG AGGTTGCAGT GAGCCAGAT CGTGCCACTG CACTCCAGCC
20601 TGGCAACAG AGAGAGACTC CGTCTCAAAA AAAGAGAAAG AAAGACACCA
20651 CTGGCTTAGT GCACTAGTGC CTAAATGCTG CTGGTCTCGG CTACAGGTGG
20701 CAAGAGGAAT GTGGGCCAGG CACTCATGCT TGGTCAAGAC TTTTCTCTTT
20751 TTGGGAGCTG GGTTCAGAG AGCACTCTGT TGGTTTCATG ACTCATTTTT
20801 GTTTTCTGAC CAAGCTCCAC AATAAGACCC TAATGTGTTT CTGTGGTATC
20851 CTCTCCTCCC TGAGTAGGCT GAGCAGAAAA TCCTTGGCCA GGCAGGGTGG
20901 CCAGAGCTGT GATGAGAGAG ATTCTTGGG CTAGGAGTAG GGTTCCCAGA
20951 GCTCTAGTTT CCAAATCTCT GCTCTGCCAT CTTCCCTTTC TCATCTTCAC
21001 ATCTGGTCAA ATCCCTCCAA AGGCACACAT CTAGGGAGCT TCATAGACAG
21051 AGACTTGGCA AAGGGGGTAC ATGTAGTTTC TCTCCTGGCT AAGACGTTGT
21101 CAGAATGGAA GAAAGGATGA GAAACATGTA CATCTAGAA AAGGCAGAAG
21151 ATGTGGGCAG GGAGATGCTG GTATGATGGC CATTTCTGTT TGAAGGTGCG
21201 CTTAGGTGAG CACCAAAGTC TTCTATGGTCA CCTTGGTGAA CCCAGACAGA
21251 ATTCTAGAGA ACCTGGTCAA GAAGAGGTCC TGAATACAC TTATGGAGAA
21301 TGCACGCTGA GAGGGGGAAG TAAACTGCTT AGGATCACCC AAAGTTGGTG
21351 GTCAAGAGTG TGGGCATCTT GATTTCTAGC CAGGATTGAG TCTCCCATAC
21401 CACTCTTATT TTTTATTATT TTGAGACAG AGTCTCACAC TGTACCCAG
21451 GCTGGAGTGC AATGGCATGA TCTCAACTCA CTGCAACCTC CACCTCCAG
21501 CAATTCTPCT GCCTCAGCCT GCCGAGTAGC TGGGATTACA GGCGCCCGCC
21551 AGCATGTCTG GCTAATTTTT TGTATTTTGA GTAGAGACGG GGTTCACATA
21601 TGTGGCCAG GCTGGTCTTG AACTCCTGAC CTCGTGATCC GCCCGCCTCA
21651 GCCTTCCAAA GTGCTGGGAT TACAAGTGTG AGCCACTGCA CCTGGCCACC
21701 ACTCTTGACC TTGACTTTTA AGGCTGTGAG CCTGTTTCTT TGCATAGAAG
21751 CATTTGGACA CAGAACTGCC GGAGTTGTGA TGGGTTTGTG GAGTGACTGT
21801 CTCTGTGCGA GATGAGCTGT GCTTTTCCCC ACCTAGGTAT GTCTCCGATG
21851 CCACTGTGGC CATCTTTGTG GCCACCCTGC TATTCATTGT GCCTTCACAG
21901 AAGCCCCAAG TTAACCTCCG CAGCCAGACT GAGGAAGGTA AGTCTCCTGT
21951 TCTGATCGCC CAGTCATCAG GACTGGAGCC CTGGAACCAA AGGGTCACTA
22001 TGGGATGCCT TGGGCCCTAG AGGGAGAAAA TCCCATCATA TCCAAGAGGA
22051 TTGGCTACAA AAGCCTGGGA AACAGTGGCT TTCAAGCCAC CGGTGGTATT
22101 ATTTAGTGCA AAATATCTTT TTTGCTTTTT AACATTTGAA TTTAACATT
22151 GAAATTTTAT TTATATTACA ACAGGAACAG AAAATGTTTC AAATTTTCCA
22201 TAACACTGAT TTCCATTGAG CACAATTTTT TGTTTTCTCT CTTCTCCCA
22251 GTCTTTGCTA ATATGCCTGT ATATTACATT ATAATCAACA CACACAGTTT
22301 GAATCCTATT TGTTTGTTGT TTCTTCTACC ACTTTTGATT GATATTACAC
22351 TATAAACATT TCCCACTATT GCTACAGTCT TCAATATAT TTTCTCTAAT
22401 AATGGCATTAT TATTGCGTTG AGGGGTTGTA ATCATTCTCC TGTTATAGAA
22451 CATTTTGGCT GTTTTGAATT TTTTATTTTC ATAAATTAAT GTTTTCTTGC
22501 ATATAGCTTT TCCTTTGAGG GTATTTTTC TTTAGGATAA ACTTCTAGGA
22551 GTAATATTGC TGGGGTGATA GAATACAAAG TCTTAATGGC CCTTAAATG
22601 TATGGCCAAA TTGCTTTTCA AAAAGGTCAT ACCAATTATC GATGCTATTG
22651 GCAGTGTGTG TAATAGTTTG ATCATATCCT CACCAGCAAT GTATATATTA
22701 TGTGAAACTT TAGCTAATTT ATAAGTAGGA GATGGTACCT CATTGCTCCT

```

FIGURE 3F

22751 ATTAGCTTTA TTCCCCCTTG ATTAGATTTT TTTTGTCTTC TAATGCTGCT
 22801 CTGGTGTGTG TGTGTGTGTG TGTGTGTGTG TGTGTGTGTG TGTGTGTCTT
 22851 TTTCTTCCCA GAAAGGAAAA CTCCATTTTA TCCCCCTCCC CTGCTGGATT
 22901 GGAAGGTAAC CCAGGAGAAA GTGCCCTGGG GCATCGTGCT GCTACTAGGG
 22951 GGCGGATTGG CTCTGGCTAA AGGATCCGAG GTAACCTCTC CAGCCACAGG
 23001 CTGCCAGAG CCCTCTTCTT CGTCAAGAGG GTGGCGTTTC TCCACCCCTC
 23051 CATCCCTGGG CTTGTGTGTT TCTGTGCTCG CATCCTTCGT ATAACCGCAC
 23101 ATTCCCTGAG GACATGGACT CTGTCTTGTC ATCTAGGAAC TCTACACCAC
 23151 ACACAGGGCC TGGAGACAG AAAGTAACCT TTGAGCGATT GCAGGAATGA
 23201 GTGAATGAGT GACCGTGGTT AGCCAAGAGA GGCAGAGGAC ACTGTCAGTT
 23251 ACCCTCTGGG GCTTGATCAC AATAATCTCT GCTTTGATTT GTCTGAGGGA
 23301 AATCTTTCTT TCCAATCCTT GTCAATATTG TTTGCTACTA CTTTGGTCC
 23351 TTCTACTGGC TACTTAACAT GGTAGCTACT TCAAAATTTT TCTTTAGCTA
 23401 AGTATGTAGC AGCGTAGGAG GTGAGGAACA TGTGGAAAA CACACAAAAA
 23451 TATAACTTTC TTTACCTCCT TCTTTCCCTC CTGGGGAAGA AATGAGCCAG
 23501 AGGGAGGGAT GAGCTAGCTT GCTGTGCTG TCCTCCAACC AACCATCTAC
 23551 CTACCCAAGT ATCCAGGAGT GTAATAGACA GACTTGGTCT AGTTATTGCT
 23601 GTTCTTCAA TATCTAGGAC ACAGCCTGGT GCCTAGTGGG TGCTAAGTTT
 23651 TTGCGGAGGT GAACAAATCC ATCCATCTAG TCACCTCTCC ATCCATCATC
 23701 CATGCATCTA TTCTGTCTT CATCCATCCA TCCGTCCGTC CATCCGTCCA
 23751 TCCATCCATC CATCCATCCA TCCATCCATC CATCCATCCA TCTGTCCATC
 23801 CATCCACCCA TCTATCTATC CACCCATCTA TCTATCCATC CATCCATCCA
 23851 CCCATCTATC CATCCATCCA TCCATCCATC CATCCATCCA TCCATCCATC
 23901 TATCCATCCA TCCATCCAAC ATTCCTATTA TGTCCCTAGT GTTATGCCAG
 23951 GCACAGAGAT TACAGAGGAG ATTGAGATAC GGTCCTGTG CGTGGCAGAC
 24001 TTTACAGACT AGGGAGGGGC ACATATATGA AAGGGCATT CAGGAAGTAG
 24051 CACACGAGCA AGGGAAAAAT GTGAGGTATT TAGCTGAGGA GAAGTAGAAG
 24101 ATGAGGCTGG TAAGGCTACC AGAAGCCACT TCTCTGAGGG CCCCAAGATA
 24151 GAGGGGTGTG GACTTGATCG TGAATGCAGT AGACAGCCAC TGAAGGACTG
 24201 AGGCCAGGGG GTGAGTTGGT CAGATCTGCA CATGAGGAAA TCACTCTGAT
 24251 GTCTGGAGTG GGGCCTGGG CTGGGCAGGG CTTGGAGGAG AACTAGCTGA
 24301 GACTCTGAGC CCTTCCATCT CACTCAGGCT CAGAACTTTG GACTCTGTGG
 24351 ACATTCTCTC CTCCTTTGGC CCCCAGCTCA GCACAGTCTC CAGCTTTACT
 24401 TCGGACTCAG ACTATTCCTG CTCAGCCTTC GTTGCTGACT TCTCTGTTCT
 24451 CCCTGAAACA GGAGTGTCTG CCCAGGCTCT GTCCCTGGCC TCTCTCTTTT
 24501 TTACACTTCA TTCTCTCCCT GGACAATCTC TTCTCAGCCC AAAGCCCTAA
 24551 ATCTAAACCT TCAATTTCTG GTTGAAATCA TTCTCCTGAG CTTCCAAAAC
 24601 TGTGGAGCAC TGAAGAGGAG GAGATGGATG TGAGACATTT GGGTGACTTG
 24651 GTGACTGACT GGGTATAAGG AAGGAGGGGA ACAGAGACCG GCAGCATGAC
 24701 TCCCAGCCTG CTGGGCTGGA TGGCTGGTGG ATGGTGAGT CATTACCAA
 24751 ACTGGGAGGC CCAGAGAGAG AAGCAGATTC TGGGCTATGG AGGATGAATG
 24801 CAGGGTGGAG CATGTTGAGT CTGTTGTGCT CTTGGGACAT CTGGATGGAC
 24851 ATTTCCAGAA GGCATATGGG TATGTAAATC CACATAGTAG GCCAGCTGGC
 24901 TGGAAATACA GATTTAGGAG ACAGCAGAGT GAGGACGGGG ATGAAAATGG
 24951 TGGGAATGGA TGAGGTCACC TATGAGGTGT AGAGAGAGAG GGTGGGAGG
 25001 GGAGGATGGG CCAAGCTTGC CCTGGCCCTG AAGGAACTGC AAGCTGGGAG
 25051 CGCTGAGATG ACTGCCCTCC TGGTGCTTCC CAGGCCCTCG GGCTGTCCGT
 25101 GTGGATGGGG AAGCAGATGG AGCCCTTGCA CGCAGTGCCC CCGGCAGCCA
 25151 TCACCTTGAT CTGTCTCTTG CTCGTTGCCG TGTTCACTGA GTGCACAAGC
 25201 AACGTGGCCA CCACCACCTT GTTCTTGCCC ATCTTTGCCT CCATGGTAAG
 25251 TAACCTGACA GTGGGGAGGA GCCCTTCCAT TTCACAGGAA CACATGGCCA
 25301 TATTGTGGGT CCCTGACGAG GCAGCAATGT CCAGGCCAGA CTCAGACCAG
 25351 GCTTTGGAGA CCCAGGCTG ACTGTGACGT GGATTTGTGG ACCCTGGATG
 25401 CCTCTGCCCT TGAGGCCCTC ACTGCTTTGC CACTCCTCTT TGTACCCTC
 25451 CTGCTGACCA AAGCACCAAC CATGGACCAA GTGCTCAAAT TTATTTTATA
 25501 AATCTAATTG GATTATTTT CAAAGCTGGG AGACAGGACT TGGGCTAAGG
 25551 AGGAGCAGGC CAGTGCCCTG GTCTCTGAGC ATGTAGCACA GGTGTGCAGG
 25601 AGGACTGCAG ACTGGGAGCA CCACTGGCTG GAAACCCAG GAAGAGGCTC
 25651 TGGAGGAGTG GGGACTTGGG AGTAGGTAGG AAGGGAGAGA GAATCTTGGG
 25701 AAGATGGAGC AGCACAAGGA AAGGCAATGG TGCACATGAC TGAGGACTCC
 25751 TGGAAAGCCTG GCTTGGTGAG CACAGGGATA AGGGATCCTG GGGAGTGGAG
 25801 AGAGGTAGCT GTCGGTTGTG GGAAGAGCTG CTGAGTGCCA GGCTAAGGCA
 25851 TTCTGTTCTA TGGACTAGCA TGTTTTTTAG TTGGGAGTTA GAAGAAAGCA
 25901 GAGCTTATAG GAAAATCAGT GGCTATGGTT TTTTTTTTTT TTTTTTTTTT
 25951 TTTTTTTTTT ATGCATTCC TTCTGTCTC CATTGCAAG ACGTACCAGC
 26001 TTCAGGGTAG TATGGAAAGA TCCCTGGTCT CGCAGTCAGA AGACCCGAGT
 26051 TCAAGATGTG GGATCTCTGA ACATGGCCCT TCAGTTCTTT CTTCGAGAG
 26101 CTGTCTGAT GGCCAAGTAA GATGAGGGCT ATGAAAAGCC TCTGTAGACT
 26151 GCAAAATGAG CATGGGAGAG GCTGTCATTA TTCTGGAATT GGGAGACAGA
 26201 TTTACAGAGG GCCTGAACAC AGGATTGAAG GTGGTGAATT TCCATTCGGC
 26251 TGCCTGGGCG TCTGCATGTA TAAAAAGCAA ACCTAAGTGG TTTTTTCTC
 26301 CTCCAAGTGA AGATGAAAGT GTTAAAAATA GCAAGGAGGT GAAAGTGTTC
 26351 AAAATAGCAA AGTGGCCTGT CTCTCTTCT CCTAAGCAGA CTGTCCAAAC
 26401 AGACGCCCCA TAGAAGGAGC ACCTTTTGAT ACTGGGCACG TGGTGGTGAT
 26451 GCCTCCTCTC TCCTAGCACA GGCCGTGGCT TTGTCTCTC CAGCCCTAAC
 26501 TGGGAGCACC GAGGGTTCCA ACCAGGCAAA TGCAGGCCCT AACGGGCTCT

FIGURE 3G

```

26551 TTGAAAACCG GCTTTTCTAG AACCAGGAAC CTCAAGTAAA AACTCCCCCA
26601 GCTACCTCTA AGGCCCATCA CACTCCTGTC TCACGCCAC CTATGAGAAA
26651 GGAAGAGGTG ATGGTCATTG AGCTGGGCTG CAGAGGAGTG TGAGGTGCAG
26701 ACACCATGAG GTACCCACAG CCAGGAAAAC GAGGATGGTC GGGGAGACGC
26751 GCCAGCGAAG AGCTGGGCCC CTGCGTGGGA CCCCTCAGTG GTTCCCAGGG
26801 GCGGTGGGAC TTGCGCAGTC CTTTCAGAGG GCTGTTTACC AACAGGAACC
26851 GTAACATTAA ACCTGCTCAG ACCCCTTGAC TCAGCAATTT CATGTCTGGG
26901 AATATATCTT AGGAAAATAA TCAGAGATGC CTACCAACAT ATGTGATGAT
26951 GATGTATGAC AGAATTATTA TACAAATATA TCCATAGTAA CAGGGGGTTT
27001 CTGAAAATAA ATTATCATAT ATTCATATAA TATGACATTA TCAGGCCATT
27051 AAAAATCACA GTTTCAAAGA GTAATAAAT GGAACATGC TCATAGTATA
27101 GTTTTTTAAA ATTGCAGATG GTATATGGCT AAAAATGTCT AATAATGCAA
27151 AGATGTATAC AGACCTTAAT CCTCTAGCCT CCTCCCTAGA GATGACCTCT
27201 GTTAATTTCT CAAATATTTT TCTGGATATT TTACACACTC ACACACTTTT
27251 TTTGAGACAG AGTTTCACTC TTGTCACCCA GGCTGGAGTG CAATGGTGTG
27301 ATCTTGGCTC ACTGCAACCT CCACCTCCCG GGTTCAGAG ATTCCTCTGC
27351 CTCAGCCTCC CGAGTAGCTG GGATTACAGG TGCCTGCCAC CTGCGCTGGC
27401 TAATTTTTTG TATTTTTAGT AGAGACGGGG TTTCACCACA TTGGTCAGGC
27451 TGGTCTCAAA CTCCTGACCT CAGGTGATCC GCCTGCCTTG GCCTCCCAAA
27501 GTGCTGGGAT TACAGGCGTG AGCCACTGCG CCCGGCCATT CATCTTAATT
27551 TTTAAAAAAT CTAACCATGA AGCCTTGGTT ATCTTGGAGA GCTTTCCTGA
27601 TTAGCACAAA AAGAAAAAAA AATCCAATTC TTTACAGCTG CATACTATTC
27651 CATATTTTGT ATGTGTCATA TTTTATTTAA CCATCCTGCT ATTAGTGACC
27701 ATTGAGTTGG CTTCCTGTGT TTTGCCGTTA CATGGTTGCA ACAAACATGT
27751 TTGCATGTGT CTGCCCTCAT GTGCATGATA CATGATTGAT TTGATAGATT
27801 TTAGGAATTA CATCATTCAT TCATACACTC AGCAAATATT TAATGAGTGC
27851 CTACTCTCTG ATAGGTGCTG TTGGATGTGG CTAATTTTAA AAGTGTAGAA
27901 TTTAAAAGGT GGCTACCAAA TTCCATGTGC AAAATGACCC CACGCATGTA
27951 TAAAAACACA CACATCCACA GATTATATAT CGGGAGAGAA GATGTGGTCC
28001 CTGGCCTCTA GGCTCTCTCA GTCTGTGGCA AGACAGACAG ACATGTGCAC
28051 CGCGCACTGT AAGGTTGAGC ACAGTCTAAG TACTCAGCAT GGTCTCTGGC
28101 ACATAGTAGG TGCCCAAGAA ATACATGTCG AATGAATTGA GGGGTAAGG
28151 CCTTCTAGGG CAGGTGGCCT CTGACCTCAG CCTTCAGTGT TCCGTAGGTG
28201 GAATTATCTG CCAGAGACGT GGCAAAAGGG AGAGGAACCA AGACTGAGGC
28251 ACAGAGGTTT AAACGTACCC GGCACATTCA GAGAATCCTT TTCAGAATCA
28301 CGTCCCCAAG AGCTTCTGTG TTCTGTACGG TGATGTTGCA GTGCTGTTTT
28351 TCCGCAGTCT CGCTCCATCG GCCTCAATCC GCTGTACATC ATGCTGCCCT
28401 GTACCCCTAG TGCCCTCCTT GCCTTCATGT TGCTGTGGC CACCCCTCCA
28451 AATGCCATCG TGTTACCTA TGGGCACCTC AAGGTTGCTG ACATGGTAAC
28501 ACAGCTGTTT TTATTTACTC CCGTCGGACT ATAACGCTGT TGTCATAAGG
28551 GATGCCCAT TTATGAATGA CAGAGTTTCA AAACGATGTC ATGTGACTTG
28601 GGAATGCCAC GGAACATCCA GACCTGTAGC CATTGTTGAC ATTTATAATG
28651 CAGCTTTTCT TCTTTTCTG AGATGATCTC AAGCCTCACA CACTGTCTTT
28701 TCTCTGAGGT GGGTTATAGA CTCTCCACC TGGAGAAGCC TGTGCAGGCA
28751 CCAGGGGAGT CTTTGAAGG GGTGAAGGTG GGGCTGAGGG ACTCATATGG
28801 CCAAGGATGA ACTTGACAAA TTAGCAAGAA CCATGAAGAT AGGCAGGGCA
28851 GGCTTAGGCA GCAGGGGGAT GCTAATGACA GTCACAGAGA TTTGTAGGGG
28901 TGCCTGAAGA GGTAGAAGCA GGGAGAGGGA GAGAGAGAGC ACTGCTGGG
28951 AGTAGATGAT GCCTTGAAA CAAATGTAGT CAGAGGAAGA ACTCTTCATT
29001 AGCTCTGTCA CTTTGTCTGG GAGAAGGGCA GCTTTCAGAC TCTGGGCTGG
29051 GAAAGAGGCA AGTGTTTGAG CCCAAGAGGC CAGAAATGTA CCTGGGACCA
29101 ATCGGGGTGT CGTTATCTCA GAGCCTCTGC TGGGTATCTC AGGGACTCCA
29151 TGAGCATTTT CAAAAAATAA GGTGGGTCCC AGAAACCATG GACTGCAAAC
29201 TTGACTCCAA TCCCCAGTA AAATATCTAC AACAGGGTAG TGAAGCGATG
29251 GTTAGTGACC ATGAGGGAAG CTTGCAGAGC AGGCATCAGA AAGAGCCTGA
29301 GGAGGTCCAC AGGGAAGCTG GCACGTCTCT GTAGGATAGT TAAGGCACTG
29351 GGGTGAGCAA TGAACCTGGA CTCACGGAAC ACTGGGCTCT GTGACCGTTT
29401 CCCTGAATGG CCTAAGCTGT TGCCCTCTGT CACTTCTCTG AGGTCATTTT
29451 CCAATGCGC ACGGGCATAG AGAACCCTAT CACTCTGCCT ACTTCCAGG
29501 GATGCCTTGA GCACTGAGGA TACCTGGGGG ACATGAAGTC GCACTGTCCT
29551 GGGGTCGGG ACACCCACAG CAGGGACAGA GCATGGCACA GGGACATCGA
29601 GGGCCAGTGA GCCGACCTTT TGTCTCTCTC TCTGAGAGCA CTAGTCCCCA
29651 GCAGGCTCA GGGTGCTGAC TCTGTCTCTT TTCCAGGTGA AAACAGGAGT
29701 CATAATGAAC ATAATTGGAG TCTTCTGTGT GTTTTGGGCT GTCAACACCT
29751 GGGGACGGGC CATATTTGAC TTGGATCATT TCCCTGACTG GGCTAATGTG
29801 ACACATATTG AGACTTAGGA AGAGCCACAA GACCACACAC ACAGCCCTTA
29851 CCTCCTCAG GACTACCGAA CCTTCTGGCA CACCTGTGAC AGAGTTTTGG
29901 GGTTCACACC CCAAAATGAC CCAACGATGT CCACACACCA CCAAAACCCA
29951 GCCAATGGGC CACCTCTTCC TCCAAGCCCA GATGCAGAGA TGGTCATGGG
30001 CAGCTGGAGG GTAGGCTCAG AAATGAAGGG AACCCTCAG TGGGCTGCTG
30051 GACCCATCTT TCCCAAGCCT TGCCATTATC TCTGTGAGGG AGGCCAGGTA
30101 CCGGAGGAT CAGGATGCAG GCTGCTGTAC CCGTCTGACC TCAAGCATCC
30151 CCCACACAGG GCTCTGGTTT TCACTCGCTT CGTCTAGAT AGTTTAAATG
30201 GGAATCGGAT CCCCTGGTTG AGAGCTAAGA CAACCACCTA CCAGTGCCCA
30251 TGTCCTTCC AGCTCACCTT GAGCAGCCTC AGATCATCTC TGTCACCTCTG
30301 GAAGGGACAC CCCAGCCAGG GACGGAATGC CTGGTCTTGA GCAACCTCCC

```

FIGURE 3H

```

30351 ACTGCTGGAG TGCAGTGGG AATCAGAGCC TCCTGAAGCC TCTGGGAAC
30401 CCTCCTGTGG CCACCACCAA AGGATGAGGA ATCTGAGTTG CCAACTTCAG
30451 GACGACACCTT GGCTTGCCAC CCACAGTGCA CCACAGGCCA ACCTACGCC
30501 TTCATCACTT GGTTCGTGTT TAATCGACTG GCCCCCTGTC CCACCTCTCC
30551 AGTGAGGCTC CTTCAACTCC TTGGTCCCCT GTTGTCTGGG TCAACATTG
30601 CCGAGAGGCC TTGGCTGGCA CCTCTGCGG TCCCCCTTTT CTCCCAGGCA
30651 GGTCACTCTT TCTGGGAGAT GCTTCCCCTG CCATCCCCAA ATAGCTAGGA
30701 TCACACTCCA AGTATGGGCA GTGATGGCGC TCTGGGGGCC ACAGTGGGCT
30751 ATCTAGGTCC TCCCTCACCT GAGGCCCAGA GTGGACACAG CTGTTAATTT
30801 CCACTGGCTA TGCCACTTCA GAGTCTTTCA TGCCAGCGTT TGAGCTCCTC
30851 TGGGTAATAA CTTCCTTTTG TTGACTGGCC TTCACAGCCA TGGCTGGTGA
30901 CAACAGAGGA TCGTTGAGAT TGAGCAGCGC TTGGTGATCT CTCAGCAAAC
30951 AACCCCTGCC CGTGGGCCAA TCTACTTGAA GTTACTCGGA CAAAGACCCC
31001 AAAGTGGGGC AACAACTCCA GAGAGGCTGT GGAATCTTC AGAAGCCCCC
31051 CTGTAAGAGA CAGACATGAG AGACAAGCAT CTCTTTTCCC CCGCAAGTCC
31101 ATTTTATTTT CTTCTGTGTC TGCTCTGGAA GAGAGGCAGT AGCAAAGAGA
31151 TGAGCTCCTG GATGGCATT TCCAGGCGAG GAGAAAGTAT GAGAGCCTCA
31201 GGAAACCCCA TCAAGGACCG AGTATGTGTC TGTTCTCTTG GGTGGGACGA
31251 TTCCTGACCA CACTGTCCAG CTCTTGCTCT CATTAATGCT TCTGTCTCCC
31301 GCGGAAAGCT CCACTGTGCT GCTGACTTGT CTCTGGTTT CTGCAGTGTG
31351 GGGAGCCGAG GGAGGTGGAT GAATGAACAG TTAGTTACGC CCTGCCACCC
31401 TGCTGGGTGC CAGGCCTTCC TGTCCCTGTT GAATCCACTA GTTATCTCGT
31451 TCGATCTTTG CAGCAACCC CTGAGATAGG AAGGTGTAT TATCTTGCTT
31501 TGTCTTTCAA AAAAAAGCG AGGCTCAGGG AGGCCAAGGG AAGTGTCCAA
31551 AGTCACACAT CAAGTTACTG GCAGTTACAG TTCCAACCAA GAGCTTCCAA
31601 CTCCATACCC CCTGCTCCTT CTGCTAGCCA TGAAGGGCTT TGGCCTTATA
31651 GGGCTTGTAG GGAAAGGTGA GTGGCCAAGA GCAAGTCCAT GCCAAGGGAA
31701 GATCTCCAAA CATGAGTCCC TGTCTGTTGC CTCCCCTGAG ATAGGCACAG
31751 GACAAAGTAT CAATGAGACA GGGTGGTCTT TGCCCTAAGA AGCAAAGTGT
31801 TTGGTTGGGG AGGGAAGTAG GGAAAAGGCT GCCACCTCCC CCCACCAAGG
31851 TACAACGTCT GACTTCCTTC CTCCCAGGCC CTCTATCACT GCCCTCTGTG
31901 CCGCTGCCGT TGAATGGCCT GCGCCACCCAG ACTGAGGGCT CTGACTGCCC
31951 ACCGAGTCTA GTGTCAGCAT TATGGCTGAC CCAGAGCAGG CTATACAGTT
32001 AGTATGATGG ATAAATAAAT GATTGGTCAG TGCAGTCAAT TAGGTGCAAG
32051 CTGTTGGTAG TAGGCAAGGT CAATGAAGGT CATCCAAGGT GGGCATTGAA
32101 GGATGAGTAG AATGGCCAGG GGTAATGGGG GAGGAAGTGG TGGGTGGGTG
32151 GAGGACTCTT CCAGACACCA TGTGGTTGAG GGCTGACAAA AAGCTGGGTG
32201 GAGGGCTTCC AGAGTGCCAA GCTCCACCT GAAGAGGCTG ACCAGAGGCC
32251 AATCCTAAAC AACTCTAGGT GTTGCTGGA GTTGCACTAA AGTGTATGGC
32301 CTCCCACACC AAACCCCTTG CTCTTAGGG CAAGGACCAC CCTGTCTCAT
32351 TGATCACTGT CCTGAGCCTA TCTCAGGGGA GGCAAAAGAG AGGGACCTGT
32401 ATTCAGAGAT CTTCCCTTGG CATGACTTGC TTTTGCCAC TTACCTTTCC
32451 CTACAAGCTC TATGAGGCCA AGGCCCTTCA TGGTTAGTGT AAGGAGCAGT
32501 GGGCATGGAG TTGGAAGATC TGGGTTGGAA CGGTAAGTGC CACTAACTCG
32551 ATGTGTGATT CTGAACACTT AACTTAGCCA TACATGCTCT CTTATTTGCT
32601 TTTGATGGCA AATAAGAGAA GGCCAGCAA ACAGTGGCTT AAACCAGAAG
32651 GTCAAATTAAT GTTACTTTT CAGGAAGTCT GTAGGTAGAT GGTGTCTGGC
32701 ATTGGCCCAA CAGCTCATT CAGCCTCCAA GGACTTGCAG TCCATAGTCC
32751 ACTCTGTCAT CTTAAAGCCT TCACACTTTT ACCCCCATGC TTGACCCCAA
32801 GGCTACATAC ACAGCT (SEQ ID NO:3)

```

FEATURES:

```

Start: 2697
Exon: 2697-2798
Intron: 2799-8874
Exon: 8875-9003
Intron: 9004-9252
Exon: 9253-9389
Intron: 9390-11975
Exon: 11976-12154
Intron: 12155-12893
Exon: 12894-13062
Intron: 13063-14905
Exon: 14906-15028
Intron: 15029-20092
Exon: 20093-20308
Intron: 20309-21836
Exon: 21837-21937
Intron: 21938-22861
Exon: 22862-22980
Intron: 22981-25083
Exon: 25084-25245
Intron: 25246-28357
Exon: 28358-28495
Intron: 28496-29686
Exon: 29687-29815

```

FIGURE 3I

Stop: 29816

SNPs:

DNA Position	Major	Minor	Domain	Protein Position	Major	Minor
609	T	G A	Beyond ORF(5')			
752	G	A	Beyond ORF(5')			
4623	A	- T	Intron			
4623	A	G T	Intron			
4699	C	T	Intron			
5062	A	G	Intron			
6158	T	C	Intron			
6573	C	A	Intron			
7120	A	G	Intron			
8411	A	C	Intron			
10035	A	G	Intron			
10849	G	A	Intron			
11916	T	C	Intron			
11962	C	T	Intron			
12333	C	T	Intron			
12375	A	C	Intron			
12418	T	C	Intron			
12603	G	A	Intron			
14225	G	C	Intron			
14416	C	T	Intron			
14643	A	C	Intron			
15612	C	T	Intron			
15685	C	T	Intron			
15971	C	T	Intron			
16175	G	A	Intron			
16589	C	A	Intron			
16980	A	-	Intron			
16993	T	- A	Intron			
17267	A	G T	Intron			
18804	C	T	Intron			
19084	C	T	Intron			
19669	G	A C T	Intron			
20397	G	C	Intron			
21575	T	C	Intron			
23363	C	T	Intron			
23413	T	C	Intron			
23945	T	G	Intron			
24483	C	A	Intron			
24643	A	G	Intron			
25329	G	A	Intron			
25421	G	A	Intron			
25797	T	G	Intron			
25926	C	T	Intron			
27289	T	C	Intron			
27591	G	T	Intron			
28245	T	A	Intron			
29337	G	A T	Intron			
29460	G	A C	Intron			
29994	A	T	Beyond ORF(3')			
30207	G	A	Beyond ORF(3')			
30497	T	G	Beyond ORF(3')			
30738	G	A	Beyond ORF(3')			
30758	T	C	Beyond ORF(3')			
31045	G	-	Beyond ORF(3')			
32591	C	T	Beyond ORF(3')			

Context:

DNA
Position

609 ACACAGGAACAGAAAACCAACACCACATGTTCTCAGTCATAAGTGGGAGTTGAACAGTG
AGAACGCATTGACACAGGGAGGGGAACATCACACACGGGGGCCTGTCAGGGGGTTGGAGG
GCAAGGGGAGGGAGAGCATTAGGACAAATACCTAATGCATGTGGGTCTTAAACCTAAAT
GTCCGGTTGATAGCTGCAGCAAACCCATGGCACATGTATACCTATGTAACAAACCTGC
ACATTCTGCACATGTATCCAGAAGCTTAAAGTAAATTAAGAAAAAGAAAAAGAAAAAG
[T, G, A]
ACTGAAGTTGTTTACTTGTCTCATTTCATGCATCCCGGAGAAAAAGGTTTGAGTGCACAT

FIGURE 3J

4623

4623

4699

5062

6158

BNSDOCID: <WO 0246407A2 | >

BNSDOCID: <WO 0246407A2 | >

- CCAATATCTGATCTTGAATTGGCCTCAACACCTGTGCATCCCTGCAGGGGTGGACCCAGT
CCCCAGTTGCTTCCCAGGGAGTACGGGGGTGGGGTGGGGATTCTCTGGCTTTCCCTCCCTG
[C, T]
CCCTCCTCTGCAGGCTGATGCTGGGCTTCATGGGCGTCACAGCCCTCCTGTCCATGTGGA
TCAGTAACACGGCAACCACGGCCATGATGGTGCCCATCGTGGAGGCCATATTGCAGCAGA
TGGAAGCCACAAGCGCAGCCACCAGGCGCGGCTGGAGCTGGTGGACAAGGGCAAGGCCA
AGGAGCTGCCAGGTGAGCCCTGGCCAGGGCACTGCCAGGCCACAACAGCAGCCTTCCCC
TCCCTCTGCTGGCAAATGCTTTGGCCACCTCCTTCTCCCTGTCTGCTTCCCGAGCCCTCCTTTAAACAC
GCATAGAGAAAAAAATAGAAAATACTGTTGTCTAAGTTTTAGGAGGGGATTATTGCA
[C, T]
ACAACTTAGATCCTTTAATAGAGCTTTGAACAAAGTCTCACCCCTCAGTTCCTCATAGTTG
CAGAAATCAGTGTGTTACCTGATTATTCATTGGGCATCTTTGAGCACTTAGGGATGC
CCCTCACTCCTTGCTACTCCTGCTCATCTCAAGGAGGCTTTTCTGACCTCCTCGAGCA
GCTCAAATCCTTCCACTCTCTGCTCCCATAGTCTGGGGCTTGGCGTCCCATGCTTGCTT
CCCTGCTAGGTGCGAAGCTCAGGGAAGACGAGTCAGCATCTACCTTGGCGTCTGCCGTGT
- 12333 GGCAACCACGGCCATGATGGTGCCCATCGTGGAGGCCATATTGCAGCAGATGGAAGCCAC
AAGCGCAGCCACCAGGCGCGGCTGGAGCTGGTGGACAAGGGCAAGGAGCTGCC
AGGTGAGCCCTGGCCAGGGCACTGCCAGGCCACAACAGCAGCCTTCCCTCCTCTGCT
GGCAAATGCTTTGGCCACCTCCTTCTCCCTGTCTGCTTCCCGAGCCCTCCTTTAAACAC
GCATAGAGAAAAAAATAGAAAATACTGTTGTCTAAGTTTTAGGAGGGGATTATTGCA
[C, T]
ACAACTTAGATCCTTTAATAGAGCTTTGAACAAAGTCTCACCCCTCAGTTCCTCATAGTTG
CAGAAATCAGTGTGTTACCTGATTATTCATTGGGCATCTTTGAGCACTTAGGGATGC
CCCTCACTCCTTGCTACTCCTGCTCATCTCAAGGAGGCTTTTCTGACCTCCTCGAGCA
GCTCAAATCCTTCCACTCTCTGCTCCCATAGTCTGGGGCTTGGCGTCCCATGCTTGCTT
CCCTGCTAGGTGCGAAGCTCAGGGAAGACGAGTCAGCATCTACCTTGGCGTCTGCCGTGT
- 12375 GCAGCAGATGGAAGCCACAAGCGCAGCCACCAGGCGCGGCTGGAGCTGGTGGACAAGGG
CAAGGCCAAGGAGCTGCCAGGTGAGCCCTGGCCAGGGCACTGCCAGGCCACAACAGCAG
CCTTCCCTCCTCTGCTGGCAAATGCTTTGGCCACCTCCTTCTCCCTGTCTGCTTCCCG
GAGCCCTCCTTTAAACACGCATAGAGAAAAAAATAGAAAATACTGTTGTCTAAGTTT
TAGGAGGGGATTATTGCACACAACCTTAGATCCTTTAATAGAGCTTTGAACAAAGTCTAC
[A, C]
CTCAGTTCCCATCAGTTGCAGAAATCAGTGTGTTACCTGATTATTCAATTGGGCATCTT
TCGAGCACTTAGGGATGCCCCCTCACTCCTTGCTACTCCTGCTCATCTCAAGGAGGCTT
TTCTGACCTCCTCGAGCAGCTCAAATCCTTCCACTCTCTGCTCCCATAGGTCTGGGGCTT
GGCGTCCCATGCTTCTTCCCTGCTAGGTGCGAAGCTCAGGGAAGACGAGTCAGCATCTA
CCTTGCCGTCTGCCGTGTTCCCTTACCATCCCCAGCCAGTGCAGTAGAGTCAGGGTCTG
- 12418 GAGCTGGTGGACAAGGGCAAGGCCAAGGAGCTGCCAGGTGAGCCCTGGCCAGGGCACTG
CCAGGCCACAACAGCAGCCTTCCCTCCTCTGCTGGCAAATGCTTTGGCCACCTCCTTC
TCCCTGTCTGCTTCCCGAGCCCTCCTTTAAACACGCATAGAGAAAAAAATAGAAAAT
ACTGTTGTCTAAGTTTTAGGAGGGGATTATTGCACACAACCTAGATCCTTTAATAGAGC
TTTGAACAAAGTCTCACCCCTCAGTTCCCATCAGTTGCAGAAATCAGTGTGTTACCTGAT
[T, C]
ATTCATTTGGGCATCTTTTCGAGCACTTAGGGATGCCCCCTCACTCCTTGCTACTCCTGCTC
ATCCTCAAGGAGGCTTTTCTGACCTCCTCGAGCAGCTCAAATCCTTCCACTCTCTGCTC
CCATAGGTCTGGGGCTTGGCGTCCCATGCTTGCTTCCCTGCTAGGTGCGAAGCTCAGGGA
AGACGAGTCAGCATCTACCTTGCCGTCTGCCGTGTTCCCTTACCATCCCCAGCCAGTGC
AGTAGAGTCAGGGTCTGTGGCTGACGGCCTGATTGCCAGACCCCTGGGCAAGGTCTGGGG
- 12603 TGTCTTAAGTTTTAGGAGGGGATTATTGCACACAACCTTAGATCCTTTAATAGAGCTTTGA
ACAAAGTCTCACCCCTCAGTTCCCATCAGTTGCAGAAATCAGTGTGTTACCTGATTATTC
ATTTGGGCATCTTTTCGAGCACTTAGGGATGCCCCCTCACTCCTTGCTACTCCTGCTCATCC
TCAAGGAGGCTTTTCTGACCTCCTCGAGCAGCTCAAATCCTTCCACTCTCTGCTCCCAT
AGGTCTGGGGCTTGGCGTCCCATGCTTGCTTCCCTGCTAGGTGCGAAGCTCAGGGAAGAC
[G, A]
AGTCAGCATCTACCTTGCCGTCTGCCGTGTTCCCTTACCATCCCCAGCCAGTGCAGTAG
AGTCAGGGTCTGTGGCTGACGGCCTGATTGCCAGACCCCTGGGCAAGGTCTGGGGCTTAC
AGAGAGGAATCGGGCACATCCCTGCCAGCACTCTTATGGAGCCCAAGTGGGGCAGCTAAA
TCAGCAGAGCTGGGATTTCCCAATCCTCAGGTGAGCAGCAGAGTCAGGACCTGGGGCTGG
GTGGGCAGCCCCATGACTGGCTCAGCTAACAGCGCTGTGCCACCACAGGGAGTCAAGT
- 14225 GGCTGGGCAGAAACGTGAGGTTCAACAACCCGTTTGTTTTAATTTTCGGGAGTGTTTTCTG
TAATGATATCCTTACAGTTCTCCAGTAACCTTCTTTGGGAAGAGCAGCCGCTCTGGGCTG
AGTGGGGAAAGCTCTGCGCTGCTTTGACACTCTTGAGCTAAAGGGGGCGCCCTGGGGC
TAGCAGAGCCCCGGGGATGGGAGGCGGGGCTGTGGTGGAAAGTGAACCTCCTCCAGCCTC
CGCTCTGGGAAGCTTTTGAATTTCTTTGCTAAGTGGGGGACCGTTCTTTGCAGAAAC
[G, C]
CACAGAGCGAGATTGCTGAGGTCTCTGCAGATCCCCAAAGATGTCAGCCAAATTACATGC
ATGTGTATAAAAGGTGTATTTTCTTTTCTTTTGGAGACAAGTCTCGCTCTGTGCG
CCAGGCTGGAGTGAGTGGCGCGATGTTGGCTCACTGCAACCTCTGCTCCTGGGTTCAA
GCGATTCTCCCGCTTACGCTCCCTATTAGCTGGGATTACAGGCGCCCGCCACCATGCCT
GGTTAATTTTGTATTTTGTAGTGAGACGGGGTTTACCATGTTGGCCAGGCCAGTCTTA
- 14416 CGGGATGGGAGGCGGGGCTGTGGTGGAAAGTGACCCCTCCTCCAGCCTCCGCTCTGGGAA
CGTTTGTAGATTTCTTTGCTAAGTGGGGGACCGTTCTTTGCAGAAACCCACAGAGCGA
GATTGCTGAGGTCTCTGCAGATCCCCAAAGATGTCAGCCAAATTACATGCATGTGTATAA
AAGGTGTATTTTCTTTTCTTTTGTAGACAAGTCTCGCTCTGTGCGCCAGGCTGGA
GTGCAGTGGCGCSATGTTGGCTCACTGCAACCTCTGCTCCTGGGTTCAAGCGATTCTCC
[C, T]
GCTTCAGCCTCCCTATTAGCTGGGATTACAGGCGCCCGCCACCATGCCTGGTTAATTTT

FIGURE 3M

GTATTTTGTAGTGGAGACGGGGTTTCACCATGTTGGCCAGGCCAGTCTTAAGCTCCTGACC
TTGTGCCCCACCTGCCTCGGCCTCCAGAGTCTTGGGAATTACAGGCGTGAGCCCTGCGC
CCGGCCACAAAGTTGTATTTTCTGGAGGGATGGGCCATAACTCCATGAGACTCTTAGC
AAGGCTTGACACACAGAAGAGTCAGTGGGTCAATTTCTCGGCCTTGCTTGTGCTGTGGC

14643 CGCCAGGCTGGAGTGCAGTGGCGCGATGTTGGCTCACTGCAACCTCTGCCTCCTGGGT
CAAGCGATTCTCCGCTTCAGCCTCCCTATTAGCTGGGATTACAGGCGCCCGCCACCATG
CCTGGTTAATTTTGTATTTTGTAGTGGAGACGGGGTTTCACCATGTTGGCCAGGCCAGTC
TTAAGCTCCTGACCTTGTGCCCCACCTGCCTCGGCCTCCAGAGTCTTGGGAATTACAGGC
TGAGCCCTTGCGCCCGGCCACAAAGTTGTATTTTCTGGAGGGATGGGCCATAACTTCC
[A, C]
TGAGACTCTTAGCAAGGCCTGGACACACAGAAGAGTCAGTGGGTCAATTTCTCGGCCTTGT
CTTGTGCTGTGGCCATGTTCTGAGGCTCCCACTCGATTAGGGGACAATGCTTGGCAATGG
ACTTGGTGGCTAGACCTCAGGAGGATGTGGCTCCACACAGGCGCGCTCTCAGGGCCCA
GCTGCTGCTCCGTCGCCACGACAGGGCCAGGCTGGCTCCACAGCTCAGCATCTGAGGT
GGGGCCCGGTGCTTCTTGTAGGTTGTTTCTGACAGCAAGGACCTCGTGAACCTTGTCT

15612 CCCATTCATCTCTGAGCGGCCCTGGGCATATCACAGGCTGTCTTTAGTATCTGCAT
TTGGCTTCCGGTGACTTTGAATTCCTCCAGAACCCTCTGATGCTGGGCACCCCGCACAG
CTCCAGCACAGGAGGAAGAGCAGGCAAGTTAAAGCAATTAAGATAAGCTGGTCCCGCA
CGTGCCAGTTCGACATTGTCTGGACAAGCTTCTCTTTGCCGTGTGGGTCCATCAGGCCAG
GTCACCGCAACCTGTGACTTAGCTCTGAGCTGAGCGCATACGCTCTGTGCCTCAATGCA
[C, T]
GGGGAGTTTAAAGTCAGTAAACCAGCAGTGATTATGACCAATCCATCCAAACCAGAC
ATTTACTGAATACCTCTGGTGTTCAGCAGTGTACAGTCTAGAAAGTTTACCTTCTCT
GTTCTAGCACACAGGCAAGTTCATCAGGGGTACCTTTGATGGCAGCCAGACTTTGGAG
AGAAACCATGACCTGTGGCTGACAAATAGCTAAAAAAAGTTATTGTTTTTCTAAAAAC
ACAAATTTATCTGTGGTGCAAGGTGATCAGGCCACACCAGGATAGAAAGTACTCAGCTC

15685 ACTTTGAATTCCTCCAGAACCCTCTGATGCTGGGCACCCCGCACAGCTCCAGCACAGG
GAGGAAGAGCAGGCAAGTTAAAGCAATTAAGATAAGCTGGTCCCGCACGTCGCAATTCGA
CATTGCTGGACAAGCTTCTCTTTGCCGTGTGGGTCCATCAGGCCAGGTCAACGCAACCC
TGTGACTTAGCTCTGAGCTGAGCGCATACGCTCTGTGCCTCAATGCACGGGGAGTTAAG
TCGAGTAAACACGACGTGATTATGACCAATCCATCCAAACCAGACATTTACTGAATA
[C, T]
CTCTGGTGTTCAGCAGTGTACAGGTCTTAGAAAGTTTACCTTCTGTTCTTAGCACAC
AGGCAAGTTCATCAGGGGTACCTTTGATGGCAGCCAGACTTTGGACAGAAACCATGACC
TGTGGCTGACAAATAGCTAAAAAAAGTTATTGTTTTTCTAAAAACACAAATTTATCTG
TGGTGCAAAAGGTGATCAGGCCACACCAGGATAGAAAGTACTCAGCTCTGAGTTAAGTGCC
TGTGCTCTGTGCCTCCATCCACAGGAAGTTCGAGCCAAGTCAAAACCAGGGGAATTTGTGA

15971 ACATTTACTGAATACCTCTGGTGTTCAGCAGTGTACAGGTCTTAGAAAGTTTACCTTC
CTGTTCTAGCACACAGGCAAGTTCATCAGGGGTACCTTTGATGGCAGCCAGACTTTGG
ACAGAAACCATGACCTGTGGCTGACAAATAGCTAAAAAAAGTTATTGTTTTTCTAAAAAC
ACACAAATTTATCTGTGGTGCAAGGTGATCAGGCCACACCAGGATAGAAAGTACTCAGC
TCTGAGTTAAGTGCCTGTGCTCTGTGCCTCCATCCACAGGAAGTTCGAGCCAAGTCAAAAC
[C, T]
AGGGGAATTTGTGACCAGAGGGAAGAGACTGCAGAGCTCAGAGGCAAAAGTGCCACGGA
AACCTGTGATTTTGTGGGAAATAGGGAATTTTCTAAGTTTCTTCTGAAGGAGGAAC
TGTTTTGAAACCTCCCATTAATAAGTTGCTATACAGGCCGGGCGGATGGCTCACACCTG
TAATCCCAACATTTTGGAGGCCGAGGTGGGCAGATCGCCTGAGGTCAGGAGTTGTAAAC
CAGCCTGGCCAACATGGTGAAACCCGCTCTCTACTAAAAATACAAAATTAGCCGGGCGT

16175 GGTGATCAGGCCACACCAGGATAGAAAGTACTCAGCTCTGAGTTAAGTGCCTGTGCTCTG
TGCTCCATCCACAGGAAGTTCGAGCCAAGTCAAAACCAGGGGAATTTGTGACCAGAGGGA
AGAGACTGCAGAGCTCAGAGGCAAAAGTGCCACGGAAACCTGTGATTTTGTGGGGAATA
TAGGGAATTTTCTAAGTTTCTTCTGAAGGAGGAAGTGTGTTGAAACCTCCCATTAATA
AGTTGCTATACAGGCCGGGCGGATGGCTCACACCTGTAATCCCAACATTTTGGAGGCC
[G, A]
AGGTGGGCAGATCGCCTGAGGTGAGGAGTTGTAAACAGCCTGGCCAACATGGTGAAACC
CCGTCTCTACTAAAAATACAAAATAGCCGGGCGTGGTAGCCACGCCTGAAATCCAG
CACTTTGGGAGGCCAAGGAGGCGGATCGCCTGAGGTGAGGAGCTCGAGACCAGCCTGGC
CAACATGGTGAAACCCATCTCTACTAAAAATACAAAAGTTAGCTGGGCATGGTGGCACA
TGCTGTAAACCCAGCTACTTGGGAGGCTGAGGCAGGAGAATTGCTTGAAGCCGGGAGGT

16589 ATCCAGCACTTTGGGAGGCCAAGGAGGCGGATCGCCTGAGGTGAGGAGCTCGAGACCA
GCCTGGCCAAACATGGTGAACCCCATCTCTACTAAAAATACAAAAGTTAGCTGGGCATGG
TGGCACATGCCTGTAAACCCAGCTACTTGGGAGGCTGAGGCAGGAGAATTGCTTGAAGCC
GGGAGGTAGAGGTTGCAATAGCCAAAGATCATGCCACTGCACTCCAGCTGGGCGACAGA
GCAAGACTCTGTCTCAAAACAAAAAAAGTTGCTATACATATTCAAAACAATCATAAT
[C, A]
ATGATAGTAAGATGACAAATATTAATGATCATTGCCCAAAACCCACTCTGTCTGCCCAT
GGACGGGCGAGGGAAACTGTTTGCATGGCTGCCTGGCCACCCAGCCTGGCTTTGACAGT
AGCTCTCTTTGCCCTGCTCTTGAATCTGCACAGGGCCAAAGTCTGTTCATTGTTTCA
AGCTCGTGAACAGGTCTCTCAGGAGATGGTCTGAACCTGCTGAGGTGAGCATCTGTG
TCTCTCATGGGCAACAGGAATAATAATGACCAACATTTATTAGTGTCTCATCATGTGC

FIGURE 3N

- 16980 TGCCTGGCCACCCAGCCTGGCTTTGACAGTAGCTCTCTTTGCCCTGCCTCTTGAATCTGC
ACCAGGGCCAAAGTCCTGTTTCATTGTTTCACATCCGTCGAACAGGTCTCTCAGGAGATGG
TCCTGAACCTGCTGCAGGTGAGCATCTGTCTCCTCATGGGGCAACAGGAATAATAATG
ACCAACATTTATTGAGTGCTCATCATGTGCCAGACATGATTTCGAGCGCTCTTTTCCTTT
CTTTATTTTATTTTATTTTATTTTATTTATTTATTTATTTATTTATTTATTTATTTATTT
[A, -]
TTTATTTTGGAGACAGTGTCTTGTCTGTCAACCATGCTGGAGTGCAGTGGTATGATCTC
GGCTCGCTGCAACCTCCACCACCTGGGTTCAAGCAATTCCTTCAGCTCAGCTCCCAAG
TAGCTGGAATTACAGGCACCCACCACCACCATGCCTGGCTAATTTTGTATTTTATTTAGTA
GAGATGGGGTTTGGCCATGTTGGCCAGGCTGGTCTTGAACCTCCTAACCTCCGGTGATCCG
CCCTCCTTGGCCTCCCAAGTGCTGGCGTTACAGATGTGAGCCACCTCGCTGGCCCAAG
- 16993 AGCCTGGCTTTGACAGTAGCTCTCTTTGCCCTGCCTCTTGAATCTGCACCAGGGCCAAAG
TCCTGTTTCATTGTTTCACATCCGTCGAACAGGTCTCTCAGGAGATGGTCTGAACCTGCT
GCAGGTGAGCATCTGTGTCTCCTCATGGGGCAACAGGAATAATAATGACCAACATTATT
GAGTGCTCATCATGTGCCAGACATGATTTCGAGCGCTCTTTTCCTTTCTTTATTTTATTT
TATTTTATTTTATTTTATTTTATTTTATTTTATTTTATTTTATTTTATTTTATTTTATTTT
[T, -, A]
CAGTGTCTTGTCTGTCAACCATGCTGGAGTGCAGTGGTATGATCTCGGCTCGCTGCAAC
CTCCACCACCTGGGTTCAAGCAATTCCTTCAGCTCAGCTCCCAAGTAGCTGGAATTAC
AGGCACCCACCACCACCATGCCTGGCTAATTTTGTATTTTATTTAGTAGAGATGGGGTTT
GCCATGTTGGCCAGGCTGGTCTTGAACCTCCTAACCTCCGGTGATCCGCCCTCCTTGGCCT
CCCAAGTGCTGGCGTTACAGATGTGAGCCACCTCGCTGGCCCAAGCACTCTTAACTT
- 17267 TATTTATTTATTTTATTTATTTTGGAGACAGTGTCTGTCTGTCAACCATGCTGGAGTGC
AGTGGTATGATCTCGGCTCGCTGCAACCTCCACCACCTGGGTTCAAGCAATTCCTTCAG
CTCAGCTCCCAAGTAGCTGGAATTACAGGCACCCACCACCACCATGCCTGGCTAATTTT
TGTATTTTATTTAGTAGAGATGGGGTTTGGCCATGTTGGCCAGGCTGGTCTTGAACCTCTAA
CCTCCGGTGATCCGCCCTCCTTGGCCTCCCAAGTGCTGGCGTTACAGATGTGAGCCAC
[A, G, T]
CGCTGGCCCAAGCACTCTTAACTTAAATTTTACACAACACCTGCGAGGTGAGCAC
TATTATTATTTATTTCCCAATTTACAGACAAGGCAACTGAGGCATGGAGAGGTGATGTGGTC
AACACAGAGCTTTGTAACAGGGAAGTAGGGGGACTGAGACTTGAACCCAGGCCCTTTGGC
TCCCACTGCATGGCATCCCTCTTGGGGAGGCTGAGGGTTGCTGTCTTAGTGGCTCCA
GACCTAAGCATGACCAGGTGTGAGAACACTAGTTGGGGCCGGGGCTGCCCTAGAACCCC
- 18804 AACTGGATCCATTTACTTGCATCTCATGCCAGCCTGGTCTACCCAGATGAAATTAACCC
AGAGATGAGAGCAAAGCTGCTCAGCAGCAGAGACTCTGAAGGCTTGGCGGTACCACTGTG
GGGCACTGGCATTGGAAGACTGCATACTCCATGCAGCCCAAGAGTCTGAGCTACTGTGG
TGTGGGGATGAGCTGCCAGCACCAATGCAGGCTCTGGCTCCTGGGCCACTAGTAATAC
CAAGGTACCCCTTATGCTGGAACCTGAAGCCCTGGCTGAGCCCAAGGGTCTCTAGGA
[C, T]
GACAGTTGGCAGCAGAGAGGTGCTTGGTAGAGCACAACCTTTACTAAGCCAAGGGTGTGG
CAGCAGAGAGGCCCTGTCTTACACCAGCAGAGCCATCCCTGTGCCGGATGTCTAGAGAGT
GTCCCTAGCGGGTGACCTCAGGACACACGGCCTTGGCCAGCAGGGAGATCCTAGCCAGC
CGTGTAGACCTGAGGTCCCATCAGTGTGCTCCTTTCTGACCCCTGAGCACCCAGAA
AGCTGTGACCTGATGTCTGTTGCTCCCATGTTCCAGGCCAAGCCACCATCACACCAACA
- 19084 GAGCCCCAGGGTCTCTAGGACGACAGTTGGCAGCAGAGAGGTGCTTGGTAGAGCACAAAC
TTTACTAAGCCAAGGGTGTGGCAGCAGAGAGGCCCTGTCTTACACCAGCAGAGCCATCCC
TGTGCCGGATGTCTAGAGAGTGTCCCTAGCGGGTGACCTCAGGACACACGGCCTTGCCC
AGCAGGGAGATCCTAGCCAGCCGTGTAGACCTGAGGTCCCATCAGTGTGCTCCTTTTC
TGACCCCTGAGCACCCAGAAAGCTGTGACCTGATGTCTTGGTGTCCCCATGTTCCAGGC
[C, T]
AAGCCACCATCACACCAACACTTGGCCCTCACACTCTCCAAGGCTGTTTCACATCCAGCAC
TGGCTTCAGGAATGAGCTCCTATTCCATCAACCCCTTCCCTCCTATGATTATGTCTCATG
GCCCCGGGAAGGGCTCTCAGGAGGGAGGGCTCTCCAGGACAATACTTGGCCCTTGCCC
ACCCCTTCAAACCAACAGTGGCTGGAACCTGGAATGTGTGAATGGAATATTACAGCATACCT
TGAGGCCTTAGTCTATGCACAGTGGCCCCAGTTATCCCCCTCCACAGCTGAGCTCCCC
- 19669 CACAGCTGAGCTCCCTTTACACCTCTCCAAGAACCTCCTCTCCTCCCTGCCTCCTCAT
GCCAACGCCACCTTAGGGGAGGCCCTGCAGGACACCTGGACAATGGACACTGGTCCCAA
GGGGGCCCATCCAGGATGGGGGTGCCATCCTGGGCTGTCTTCTTCTTGGCCTAGCCATG
CTTGCTGCTAACCCAGGGTCTCCTGGATCCCTAATCCTGCACCTCCAACCTCCAGGGAAC
ACAAGGACCCATTCTGCCCTGACTAGCCCTGTCTGCCAGGGTTCATACTACTCCTGTC
[G, A, C, T]
TCTCCCTGAGCCACCTTGGTGTGGGGTGGGCATCCCAACACCATCGAAGGCAGCTCCA
GGCTGAGGTGGAAGGAGGAAGACTTGGGAAGCATGTGAGGGAGCCCTGTTCCACCTTGC
CAGGCTCCGAAGCTCCTTATGGCCTTCCCCAGGTGACCTGGAGCAGCCAGTCTCCAG
GTGCTGGGCACCTGCCGAGACCCCTAGCCTCTCTACAGAGACTTTTCCCTAGTACAT
TCTGGGATGGAAGAACAGGAGAGGGAAGAGGAGGAGGCTTTCTCCAGGCCATA
- 20397 AAAAGTCCTGGGGCTGCGGGCTAGAGAGCAAGAAAAACGAGAAGGCTGCCCTCAAGGTG
CTGCAGGAGGATACCGGAAGCTGGGGCCCTTGTCTTCCGCGAGAGATCAACGTGCTGATC
TGCTTCTTCTGCTGGTGCATCTGTGGTTCTCCCGAGACCCCGGCTTCATGCCCGGCTGG

FIGURE 30

CTGACTGTTGCCTGGGTGGAGGGTGAGACAAAGTAAGTCTTGGATTCAATAGAAATCGCT
GGCTTAGGGCCAGGCGCGTTGGCTCACACATGTAATCCCAGCACTTTGGGAGGCTGAGGT
[G, C]
GGTGGGTCACTTGAAGTCAAGGAGATCGAGACCATCTGGCCAACATGGTGAAACCCCTGTC
TCTACTAAAAATACAGAAAATTAGCGAGGCATGGTGGCACATGCCTGTAGTCCCAGCTAC
TTGGGAGACTGAGGCAGGAGAATCACTTGAACCCAGGAGGCAGAGGTGTCAGTGAGCCCA
GATCGTGCCACTGCACTCCAGCCTGGGCAACAGAGAGAGACTCCGTCTCAAAAAAGAGA
AAGAAAGACACCCTGGCTTAGTGCATAGTGCCTAAATGCTGCTGGTCTCGGCTACAGG

21575 AGGTCTCTGAAATACACTTATGGAGAATGCACGCTGAGAGGGGGAAGTAACTGCTTAGGA
TCACCCAAAGTTGGTGGTCAAGAGTGTGGGCATCTTGATTCTAGCCAGGATTCAGTCTC
CCATACCACTCTTATTTTTTATTTTTTTGAGACAGAGTCTCACACTGTCACCCAGGCTG
GAGTGCAATGGCATGATCTCAACTCACTGCAACCTCCACCTCCAGCAATTCTCCTGCCT
CAGCCTGCCGAGTAGCTGGGATTACAGGCGCCCGCCAGCATGTCTGGCTAATTTTTTGTA
[T, C]
TTTTAGTAGAGACGGGGTTTCACTATGTTGGCCAGGCTGGTCTTGAACCTCGACCTCGT
GATCCGCCCCCTCAGCCTCCCAAAGTGTGGGATTACAAGTGTGAGCCACTGCACCTGG
CCACCACTCTTGACCTTGACTTTTAAGGCTGTGAGCCTGTTCTTTGCATAGAAGCATTT
GGACACAGAAGTCCCGAGTTGTGATGGGTTTGTGTAGTGACTGTCTCTGTGCGCAGATGA
GCTGTGCTTTTCCCACCTAGGTATGTCTCCGATGCCACTGTGGCCATCTTTGTGGCCAC

23363 TGTGTGTTTTCTGTGCCTGCATCCTTCGTATAACCGCACATTCCTTGAGGACATGGACTCT
GTCTTGTCTATCTAGGAACCTACACCACACACAGGGCCTGGAAGACAGAAAGTAACCTTT
GAGGATTGCAAGGAATGAGTGAATGAGTGACCGTGGTTAGCCAAAGAGAGGCAGAGGCAC
TGTCAGTTACCTCTGGGGCTTGATCACAATAATCTCTGCTTTGATTGTCTGAGGAAA
TCTTTCTTTTCCAATCCTTGTCAATATTGTTTGCTACTACTTTTGGTCTTCTACTGGCTA
[C, T]
TTAACATGGTAGCTACTTCAAAATTTTTCTTTAGCTAAGTATGTAGCAGCGTAGGAGGTG
AGGAACATGTTGGAAAACACACAAAATATAACTTTCTTTACCTCCTTCTTTCCCTCCTG
GGGAAGAAATGAGCCAGAGGGAGGATGAGCTAGCTTGTCTGTCTGTCTTCCAAACCAAC
CATCTACTTACCCAAAGTATCCAGGAGTGAATAGACAGACTTGGTCTAGTTATTGCTGTT
TCTTCAATATCTAGGACACAGCCTGGTGCTTAGTGGGTGCTAAGTTTTTGGGAGGTGAA

23413 CATGGACTCTGTCTTGTCTATCTAGGAACCTACACCACACACAGGGCCTGGAAGACAGAA
AGTAACCTTTGAGCGATTGCAGGAATGAGTGAATGAGTGACCGTGGTTAGCCAAGAGAGG
CAGAGGACACTGTCTAGTTACCTCTGGGGCTTGATCACAATAATCTCTGCTTTGATTGTG
CTGAGGGAATCTTTCTTTCCAATCCTTGTCAATATTGTTTGCTACTACTTTTGGTCTCTT
CTACTGGCTACTTAAATGTTAGTACTTCAAAATTTTTCTTTAGCTAAGTATGTAGCAG
[T, C]
GTAGGAGGTGAGGAACATGTTGGAAAACACACAAAATATAACTTTCTTTACCTCCTTCT
TTCCCTCCTGGGGAAGAAATGAGCCAGAGGGAGGGATGAGCTAGCTTGTCTGTCTGTCTC
TCCAACCAACCATCTACCTACCCAAGTATCCAGGAGTGAATAGACAGACTTGGTCTAGT
TATTGCTGTTTCTCAATATCTAGGACACAGCCTGGTGCTTAGTGGGTGCTAAGTTTTTG
CGGAGGTGAACAAATCCATCCATCTAGTCACCTCTCCATCCATCATCCATGCATCTATTC

23945 AAGTTTTTGCGGAGGTGAACAAATCCATCCATCTAGTCACCTCTCCATCCATCATCCATG
CATCTATTCTATGTCTTCTCATCCATCCATCCGTCCTCCATCCGTCCTCCATCCATCCATC
CATCCATCCATCCATCCATCCATCCATCTGTCCATCCATCCACCCATCTATCTATCCACC
CATCTATCTATCCATCCATCCATCCATCCATCCATCTATCCATCCATCCATCCATCCATC
CATCCATCCATCCATCTATCCATCCATCCATCCATCCATCCATCCATCCATCCATCCATC
[T, G]
GCCAGGCACAGAGATTACAGAGGAGATTGAGATACGGTCCCTGTTCTGTTGGCAGACTTCAC
AGACTAGGGAGGGGCACATATATGAAAGGGCATTTCAGGAAGTAGCACACAGCAGAGGGA
AAAATGTGAGGTATTTAGCTGAGGAGAAGTAGAAGATGAGGCTGGTAAGGCTACCAAGAAG
CCACTTCTGAGGGCCCCAAGATAGAGGGGTGTGGACTTGATCGTGAATGCAGTAGACA
GCCACTGAAGGACTGAGGCCAGGGGTGAGTTGGTCAGATCTGCACATGAGGAAATCACT

24483 ACAGCCACTGAAGGACTGAGGCCAGGGGGTGAAGTGGTCAGATCTGCACATGAGGAAATC
ACTCTGATGTCTGGAGTGGGGGCTGGGCTGGGCAGGGCTGGAGGAGAACTAGCTGAGA
CTCTGCAGCCTTCCATCTCACTCAGGCTCAGAACTTTGGACTCTGTGGACATTTCTCTCCT
CCTTTGGCCCCCAGCTCAGCACAGTCTCCAGCTTTACTTCGGACTCAGACTATTCTCTGCT
CAGCCTTCTGTGACTTCTCTGTTCTCCCTGAAACAGGAGTGTCTGCCAGGGCTCTGT
[C, A]
CTTGGCCTCTCCTCTTTTTACACTTCAATCTCTCTCCCTGGACATCTCTTCTCAGCCAAA
GCCCTAAATCTAAACCTTCAATTTCTGGTTGAAATCATTCTCTCTGAGCTTCCAAACTGT
GGAGCACTGAAGAGGAGGAGATGGATGTGAGACATTTGGGTGACTTGGTGACTGACTGGG
TATAAGGAAGGAGGGGAACAGAGACCAGGAGCATGACTCCAGCCTGCTGGGCTGGATGG
CTGGTGGATGGTGAGTCCATTACCAAACTGGGAGGCCAGAGAGAGAAGCAGATTCTGG

24643 CTCTGTGGACATTTCTCTCTCTTTGGCCCCCAGCTCAGCACAGTCTCCAGCTTTACTTC
GGACTCAGACTATCTCTCTCAGCCTTCGTTGCTGACTTCTCTGTTCTCCTGAAACAGG
AGTGTCTGCCAGGCTCTGTCTTGGCCTCTCTCTTTTTACACTTCAATTTCTCTCTCTCTG
ACAATCTCTCTCAGCCCCAAGCCCTAAATCTAAACCTTCAATTTCTGGTTGAAATCATT
CTCTGAGCTTCCAAAACCTGTGGAGCACTGAAGAGGAGGAGATGGATGTGAGACATTTGG
[A, G]
TGACTTGGTGACTGACTGGGTATAAGGAAGGAGGGGAACAGAGACCGGCAGCATGACTCC

FIGURE 3P

- CAGCCTGCTGGGCTGGATGGCTGGTGGATGGTGAATTCACCAAACCTGGGAGGCCCA
GAGAGAGAAGCAGATTCTGGGCTATGGAGGATGAATGCAGGGTGGAGCATGTTGAGTCTG
TTGTGCTCTTGGGACATCTGGATGGACATTTCCAGAAGGCATATGGGTATGTAATCCAC
ATAGTAGGCCAGCTGGCTGGAAATACAGATTTAGGAGACAGCAGAGTGAGGACGGGGATG
- 25329 TGAAGGAACTGCAAGCTGGGAGCGCTGAGATGACTGCCCTCCTGGTGTCTCCAGGCCTC
GGGGCTGTCCGTGTGGATGGGGAAGCAGATGGAGCCCTTGACCGCAGTCCCCCGGCAGC
CATCACCTTGATCTTGTCTTGTCTCGTTGCCGTGTTCACTGAGTGCACAAGCAACGTGGC
CACCACCACCTTGTCTGCCCATCTTTGCCCTCCATGGTAAGTAACCTGACAGTGGGGAG
GAGCCCTTCCATTTACAGGAACACATGGCCATATTGTGGGTCCCTGACGAGGCAGCAAT
[G, A]
TCCAGGCCAGACTCAGACCAGGCTTTGGAGACCAGGTCTGACTGTGACGTGGATTGTG
GACCCTGGATGCCTCTGCCCTGAGGCCTCCACTGCTTTGCCACTCCTCTTTGTACCCCT
CCTGCTGACCAAAGCACCACCATGGACCAAGTGCTCAAATTTATTTTATAAATCTAATT
GGATTATTTTCAAGCTGGGAGACAGGACTTGGGCTAAGGAGGAGCAGGCCAGTGGCGT
GGTCTCTGAGCATGTAGCACAGGTGTGCAGGAGGACTGCAGACTGGGAGCACCCTGGCT
- 25421 AGCCCTTGACCGCAGTGCCCCGGCAGCCATCACCTTGATCTTGTCTTGTCTCGTTGCCG
TGTTCACTGAGTGCACAAGCAACGTGGCCACCACCACCTTGTTCCTGCCCATCTTTGCCT
CCATGGTAAGTAACCTGACAGTGGGGAGGAGCCCTTCCATTTACAGGAACACATGGCCA
TATTGTGGGTCCCTGACGAGGCAGCAATGTCCAGGCCAGACTCAGACCAGGCTTTGGAGA
CCCAGGTCTGACTGTGACGTGGATTGTGGACCCTGGATGCCTCTGCCCTGAGGCCTC
[G, A]
CTGCTTTGCCACTCCTCTTTGTACCCCTCCTGCTGACCAAAGCACCACCATGGACCAAG
TGCTCAAATTTATTTTATAAATCTAATTGGATTATTTTCAAGCTGGGAGACAGGACTT
GGCTAAGGAGGAGCAGGCCAGTGGCGTGGTCTCTGAGCATGTAGCACAGGTGTGCAGGA
GGACTGCAGACTGGGAGCACCCTGGCTGGAAACCCAGGAAGAGGCCCTTGGAGGAGTGG
GGACTTGGGAGTAGGTAGGAAGGAGAGAGAATTCTGGGAGATGGAGCAGCACAAGGAA
- 25797 TATAAATCTAATTGGATTATTTTCAAGCTGGGAGACAGGACTTGGGCTAAGGAGGAGC
AGGCCAGTGCCGTGCTCTGAGCATGTAGCACAGGTGTGCAGGAGGACTGCAGACTGGG
AGCACCCTGGCTGGAAACCCAGGAAGAGGCCCTTGGAGGAGTGGGGACTTGGGAGTAGG
TAGGAAGGGAGAGAGAATTCTGGGAAGATGGAGCAGCACAGGAAGGCAATGTTGCACA
TGACTGAGGACTCCTGGAAGCCTGGCTTGGTGGAGCAGGGATAAGGGATCCTGGGAGT
[T, G]
GAGAGAGGTAGCTGTGGTGTGGGAAAAGCTGCTGAGTGCCAGGCTAAGGCATTCTGTT
CTATGGACTAGCATGTTTTTTAGTTGGGAGTTAGAAGAAAGCAGAGCTTATAGGAAATC
AGTGGCTATGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTATGCAATTCCTTCTGTC
ATCCATTGCAAAGACGTACCAGCTTCAGGGTAGTATGGAAAGATCCCTGGTCTCGCAGTC
AGAAGACCCGAGTTCAAGATGTGGGATCTCTGAACATGGCCCTTCAGTTCCTTCTCCGA
- 25926 GGCTGGAAACCCAGGAAGAGGCCCTTGGAGGAGTGGGGACTTGGGAGTAGGTAGGAAGGG
AGAGAGAATTCTGGGAAGATGGAGCAGCACAAGGAAGGCAATGGTGACATGACTGAGG
ACTCCTGGAAAGCCTGGCTTGGTGGAGCAGGGATAAGGGATCCTGGGGAGTGGAGAGAG
TAGCTGTCCGTTGTGGGAAAAGCTGCTGAGTGCCAGGCTAAGGCATTCTGTTCTATGGAC
TAGCATGTTTTTTAGTTGGGAGTTAGAGAAAGCAGAGCTTATAGGAAATCAGTGGCTA
[C, T]
GGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTATGCAATTCCTTCTGTCATCCATTGC
AAAGACGTACCAGCTTCAGGGTAGTATGGAAAGATCCCTGGTCTCGCAGTCAGAAGACCC
GAGTTCAAGATGTGGGATCTCTGAACATGGCCCTTCAGTTCTTTCTCCGAGAGCTGTGC
TGATGGCCAAGTAAGATGAGGGCTATGAAAGCCTCTGTAGACTGCAAAATGAGCATGGG
AGAGGCTGTCATTATCTGGAATTGGGAGACAGATTTACAGAGGGCCTGAACACAGGATT
- 27289 AACAGGGGGTTGCTGAAATAAATTATCATATATTCATATAATATGACATTATCAGGCCA
TTAAAAATCACAGTTTCAAAGAGTAATAAAATGGGAACATGCTCATAGTATAGTTTTTA
AAATTGCAGATGGTATATGGCTAAAAATGTCTAATAATGCAAAGATGTATACAGACCTTA
ATCCTCTAGCCTCCTCCCTAGAGATGACCTCTGTAAATTTCTCAAATATTTTCTGGATA
TTTTACACACTCACACACTTTTTTTGAGACAGAGTTTCACTCTTGTCAACCCAGGCTGGAG
[T, C]
GCAATGGTGTGATCTTGGCTCACTGCAACCTCCACCTCCCGGGTTCAAGAGATTCTCCTG
CCTCAGCCTCCCGAGTAGCTGGGATTACAGGTGCCTGCCACCTTGCCTGGCTAATTTTTT
GTATTTTATAGAGACGGGGTTTACCACATTGGTCAGGCTGGTCTCAAATCCTGACC
TCAGGTGATCCGCCTGCCTTGGCCTCCCAAAGTGCTGGGATTACAGGCGTGAGCCACTGC
GCCCCGCCATTATCTTAATTTTTTAAAAAATCTAACCATGAAGCCTTGGTTATCTTGGAG
- 27591 CAATGGTGTGATCTTGGCTCACTGCAACCTCCACCTCCCGGGTTCAAGAGATTCTCTGC
CTCAGCCTCCCGAGTAGCTGGGATTACAGGTGCCTGCCACCTTGCCTGGCTAATTTTTT
TATTTTTTAGTAGAGACGGGGTTTACCACATTGGTCAGGCTGGTCTCAAATCCTGACCT
CAGGTGATCCGCCTGCCTTGGCCTCCCAAAGTGCTGGGATTACAGGCGTGAGCCACTGCG
CCCGGCCATTATCTTAATTTTTTAAAAAATCTAACCATGAAGCCTTGGTTATCTTGGAGA
[G, T]
CTTTCCTGATTAGCACAAAAAGAAAAAATCCAATTCTTTACAGCTGCATACTATTCC
ATTATTGTATGTGTATATTTTATTTAACCATCCTGCTATTAGTGACCATTGAGTTGGC
TTCCTGTGTTTTGCCGTTACATGGTTGCAACAAACATGTTTGCATGTGTCTGCCCTCATG
TCGATGATACATGATTGATTGATAGATTTTAGGAATTACATCATTCATTATACACTCA
GCAAATATTTAATGAGTGCTACTCTCTGATAGGTGCTGTTGGATGTGGCTAAATTTTAA

FIGURE 3Q

- 28245 CATGTATAAAACACACATCCACAGATTTATATGCGGGAGAGAAGATGTGGTCCCTGG
CCTCTAGGCTCTCTCAGTCTGTGGCAAGACAGACAGACATGTGCACGCGGCACTGTAAGG
TTGAGCACAGTCTAAGTACTCAGCATGGTCTCTGGCACATAGTAGGTGCCCAAGAAATAC
ATGTGGAATGAATTGAGGGGGTAAGGCCTTCTAGGGCAGGTGGCCTCTGACCTCAGCCTT
CAGTGTTCCTGAGGTGGAATTATCTGCCAGAGACGTGGCAAAGGGAGAGGAACCAAGAC
[T, A]
GAGGCACAGAGGTTCAACGCTACCCGGCACATTGAGAGAATCCTTTTCAGAATCACGTCC
CCAAGAGCTTCTGTGTTCTGTACGGTGTATGTCAGTGTCTGTTTTCCGCACTCTCGCTC
CATCGGCTCAATCCGCTGTACATCATGCTGCCCTGTACCTGAGTGCCTCCTTTGCCCTT
CATGTTGCCCTGTGGCCACCCCTCCAAATGCCATCGTGTTCACCTATGGGCACCTCAAGGT
TGCTGACATGGTAACACAGCTGTTTTATTACTCCCGTCGGACTATAACGCTGTTGTCA
- 29337 CAGCTCTGGGCTGGGAAAGAGGCAAGTGTGAGCCCAAGAGGCCAGAAATGTACCTGGG
ACCAATCGGGTGTCTGTTATCTCAGAGCCTCTGCTGGGTATCTCAGGGACTCCATGAGCA
TTTTCAAAAAAAGGTGGGTCCCAGAAACCATGGACTGCAAACTTGACTCCAATCCCCAGT
AGTAAAAATATCTACAACAGGGTAGTGAAGCGATGGTTAGTGACCATGAGGGAAGCTTGCA
GAGCAGGCATCAGAAAGAGCCTGAGGAGGTCCACAGGGAAGCTGGCACGCTCTGTAGGA
[G, A, T]
AGTTAAGGCACTGGGGTGAAGCAATGAACCTGGACTCACGGAACACTGGGCTCTGTGACCG
TTTCCCTGAATGGCCTAAGCTGTTGCCCTCTGTCACTTCTCTGAGGTCAATTTCCAAATG
CGCAGGGCATAGAGAACCCATCCACTCTGCCTACTTCCAGGGATGCCTTGAGCACTGAC
GGATACCTGGGGACATGAAGTCGACTGTCTGGGGTGGGACACCCAGCCAGGGGAC
AGAGCATGGCACAGGGACATCAGAGGCCAGTGAGCCGACCTTTGTCTCTCTCTGTGAGA
- 29460 TCAAAAAAAGGTGGGTCCCAGAAACCATGGACTGCAAACTTGACTCCAATCCCCAGT
AAAATATCTACAACAGGGTAGTGAAGCGATGGTTAGTGACCATGAGGGAAGCTTGAGAG
CAGGCATCAGAAAGAGCCTGAGGAGGTCCACAGGGAAGCTGGCACGCTCTGTAGGATAG
TTAAGGCACTGGGGTGAAGCAATGAACCTGGACTCACGGAACACTGGGCTCTGTGACCGTT
TCCCTGAATGGCCTAAGCTGTTGCCCTCTGTCACTTCTCTGAGGTCAATTTCCAAATGCG
[G, A, C]
ACGGGCATAGAGAACCCATCCACTCTGCCTACTTCCAGGGATGCCTTGAGCACTGAGGA
TACCTGGGGGACATGAAGTCGCACTGTCTGGGGTGGGACACCCAGCCAGGGACAGA
GCATGGGCACAGGGACATCGAGGCCAGTGAGCCGACCTTTGTCTCTCTGTGAGAGCA
CTAGTCCCAGCAGGCCTCAGGGTGTGACTCTGTCTCTTTCCAGGTGAAAACAGGAGT
CATATGAACATAATTGGAGTCTTCTGTGTGTTTTGGCTGTCAACACCTGGGGACGGGC
- 29994 CAGGAGTCATAATGAACATAATTGGAGTCTTCTGTGTGTTTTGGCTGTCAACACCTGGG
GACGGGCCATATTTGACTTGATCATTTCCCTGACTGGGCTAATGTGACACATATTGAGA
CTTAGGAAGAGCCACAAGACCACACACAGCCCTTACCTCTCAGGACTACCGAACCT
TCTGGCACACCTTGTAACAGAGTTTGGGGTTCACACCCCAAATGACCCAACGATGTCCA
CACACCACCAAACCCAGCCAATGGGCCACCTCTCTCCAGCCAGATGCAGAGATGG
[A, T]
CATGGGCAGCTGGAGGGTAGGCTCAGAAATGAAGGGAACCCCTCAGTGGGCTGTGGACC
CATCTTTCCCAAGCCTTGCCATTATCTCTGTGAGGGAGGCCAGGTAGCCGAGGGATCAGG
ATGAGGCTGTGTACCCGCTCTGCCTCAAGCATCCCCACACAGGGCTCTGGTTTTAC
TCGCTTCGTCTAGATAGTTAAATGGGAATCGGATCCCTGGTTGAGAGCTAAGACAAC
CACCTACCAAGTCCCATGTCCTTCCAGCTCACCTTGAGCAGCCTCAGATCATCTCTGTCT
- 30207 CACCCCAAAATGACCCACGATGTCCACACACCACCAAAACCCAGCCAATGGGCCACCTC
TTCTCCAAAGCCAGATGCAGAGATGGTCATGGGCAGCTGGAGGGTAGGCTCAGAAATGA
AGGGAACCCCTCAGTGGGCTGCTGGACCCATCTTTCCCAAGCCTTGCCATTATCTCTGTG
AGGGAGGCCAGGTAGCCGAGGGATCAGGATGCAGGCTGCTGTACCCGCTCTGCCTCAAGC
ATCCCCACACAGGGCTCTGGTTTTCTACTCGCTTCGTCTAGATAGTTTAAATGGGAATC
[G, A]
GATCCCCGTTGAGAGCTAAGACAACCACTACCAAGTGCCTATGTCCCTTCCAGCTCAC
CTTGAGCAGCCTCAGATCATCTCTGTCACTCTGGAAGGGACACCCAGCCAGGGACGGAA
TGCTTGGTCTTGAGCAACCTCCCACTGCTGGAGTGCGAGTGGGAATCAGAGCCTCTGAA
GCCTCTGGGAACCTCTCTGTGGCCACCACCAAGGATGAGGAATCTGAGTTGCCAATCT
CAGGACGACACCTGGCTTGCCACCACAGTGACACCAGGCCAACCTACGCCCTTCATCA
- 30497 AATGGGAATCGGATCCCTGGTTGAGAGCTAAGACAACCACTACCAAGTGCCTATGTCCC
TTCCAGCTCACCTTGAGCAGCTCAGATCATCTCTGTCACTCTGGAAGGGACACCCAGC
CAGGGACGGAAATGCCCTGGTCTTGAGCAACCTCCCACTGCTGGAGTGCGAGTGGGAATCAG
AGCCTCCTGAAGCCTCTGGGAACCTCTCTGTGGCCACCACCAAGGATGAGGAATCTGA
GTTGCCAATCTCAGGACGACACCTGGCTTGCCACCACAGTGACACCAGGCCAACCTAC
[T, G]
CCCTTCATCACTTGGTTCTGTTTTAATCGACTGGCCCCCTGTCCCACTCTCCAGTGAGC
CTCTTCAACTCCTTGGTCCCTGTGTCTGGGTCAACATTTGCCGAGACGCTTGGCTG
GCACCTCTGGGGTCCCCCTTTTCTCCAGGCAGGTCACTTTTCTGGGAGATGCTTCCC
CTGCCATCCCCAAATAGCTAGGATCACACTCCAAGTATGGGCAGTGATGGCGCTCTGGGG
GCCACAGTGGGCTATCTAGGTCTCTCCCTCACCTGAGGCCAGAGTGACACAGCTGTTAA
- 30738 TTGCCAACTTCAGGACGACACCTGGCTTGCCACCCACAGTGACACCAGGCCAACCTACG
CCCTTCATCACTTGGTTCTGTTTTAATCGACTGGCCCCCTGTCCCACTCTCCAGTGAGC
CTCTTCAACTCCTTGGTCCCTGTGTCTGGGTCAACATTTGCCGAGACGCTTGGCTG

FIGURE 3R

GCACCCCTCTGGGGTCCCCCTTTTCTCCAGGCAGGTCATCTTTCTGGGAGATGCTTCCC
CTGCCATCCCCAAATAGCTAGGATCACACTCCAAGTATGGGCAGTGATGGCGCTCTGGGG
[G, A]
CCACAGTGGGCTATCTAGGTCCTCCCTCACCTGAGGCCAGAGTGGACACAGCTGTTAAT
TTCCACTGGCTATGCCACTTCAGAGTCTTTCATGCCAGCGTTTGAGCTCCTCTGGGTAAA
ATCTTCCCTTTGTTGACTGGCCTTCACAGCCATGGCTGGTGACAAACAGAGGATCGTTGAG
ATTGAGCAGCGCTTGGTGATCTCTCAGCAAACAACCCCTGCCCGTGGGCCAATCTACTTTG
AAGTTACTCGGACAAAGACCCCAAAGTGGGGCAACAACCTCCAGAGAGGCTGTGGGAATCT

30758 CCTGGCTTGCCACCCACAGTGCACCACAGGCCAACCTACGCCCTTCATCACTTGGTTCTG
TTTTAATCGACTGGCCCCCTGTCCCACCTCTCCAGTGAGCCTCCTTCAACTCCTTGGTCC
CCTGTTGTCTGGGTCAACATTTGCCGAGACGCTTGGCTGGCACCCCTCTGGGGTCCCCCT
TTTCTCCAGGCAGGTCATCTTTTCTGGGAGATGCTTCCCCTGCCATCCCCAAATAGCTA
GGATCACACTCCAAGTATGGGCAGTGATGGCGCTCTGGGGGCCACAGTGGGCTATCTAGG
[T, C]
CCTCCCTCACCTGAGGCCAGAGTGGACACAGCTGTTAATTTCCACTGGCTATGCCACTT
CAGAGTCTTTCATGCCAGCGTTTGAGCTCCTCTGGGTAAAATCTTCCCTTTGTTGACTGG
CCTTCACAGCCATGGCTGGTGACAAACAGAGGATCGTTGAGATTGAGCAGCGCTTGGTGAT
CTCTCAGCAAACAACCCCTGCCCGTGGGCCAATCTACTTGAAGTTACTCGGACAAAGACC
CCAAAGTGGGGCAACAACCTCCAGAGAGGCTGTGGGAATCTTCAGAAGCCCCCTGTAAAG

31045 TGGGCTATCTAGGTCCTCCCTCACCTGAGGCCAGAGTGGACACAGCTGTTAATTTCCAC
TGGCTATGCCACTTCAGAGTCTTTCATGCCAGCGTTTGAGCTCCTCTGGGTAAAATCTTC
CCTTTGTTGACTGGCCTTCACAGCCATGGCTGGTGACAAACAGAGGATCGTTGAGATTGAG
CAGCGCTTGGTGATCTCTCAGCAAACAACCCCTGCCCGTGGGCCAATCTACTTGAAGTTA
CTCGGACAAAGACCCCAAAGTGGGGCAACAACCTCCAGAGAGGCTGTGGGAATCTTCAGAA
[G, -]
CCCCCTGTAAGAGACAGACATGAGAGACAAGCATCTTCTTTCCCCGCAAGTCCATTTT
ATTTCTTCTTGTGCTGCTCTGGAAGAGAGGAGTAGCAAAGAGATGAGCTCCTGGATGG
CATTTTCCAGGGCAGGAGAAAAGTATGAGAGCCTCAGGAAACCCCATCAAGGACCGAGTAT
GTGCTGCTGCTTCTTGGGTGGGACGATTCTTGACCACACTGTCCAGCTCTTGCTCTCATT
AATGCTCTGCTCCCGGAAAGCTCCACTGTGCTGCTGACTTGTCTCTGGTTCCTGCA

32591 AGTGATGGCCTCCCCAACCAACCCCTTTGCTTCTTAGGGCAAGGACCACCCTGTCTCAT
TGATCACTGTCTGAGCCTATCTCAGGGGAGGCAAAAGAGAGGACCTGTATTTCAGAGAT
CTTCCCTTGGCATGACTTGCTTTTGGCCACTTACCTTTCCCTACAAGCTCTATGAGGCCA
AGGCCCTTCATGGTTAGTGTAAGGAGCAGTGGGCATGGAGTTGGAAGATCTGGGTTGGAA
CGGTAAGTCCCACTAAGTGTGATTCTGAACACTTAAGTATAGCCATACATGCTCT
[C, T]
TTATTTGCTTTTGATGGCAAATAAGAGAAGGCCAGCAACAGTGGCTTAAACCAGAAGG
TCAATTAATGTTTACTTTTCAGGAAGTCTGTAGGTAGATGGTTGCTGGCATTTGGCCCAAC
AGCTCATTTTCAGCCTCCAAGGACTTGCCTCCATAGTCCACTCTGTCTCTTAAAGCCTT
CACACTTTTACCCCATGCTTGACCCCGAGGCTACATACACAGT

Bac accession number: AC034305

Chromosome: 17

FIGURE 3S

SEQUENCE LISTING

<110> PE CORPORATION (NY)

<120> ISOLATED HUMAN TRANSPORTER PROTEINS,
NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS,
AND USES THEREOF

<130> CL000662PCT

<140> TO BE ASSIGNED

<141> 2001-12-05

<150> 09/729,094

<151> 2000-12-05

<150> 60/211,220

<151> 2000-06-13

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 2223

<212> DNA

<213> Homo sapiens

<400> 1

```

gtctccctcc cgcgcgatgg cctcggcgct gagctatgtc tccaagttca agtccttcgt 60
gatcttggtc gtcaccccg ccttgctgct gccactcgtc attctgatgc ccgccaagtt 120
tgtcaggtgt gcctacgtca tcctcctcat ggccatttac tggtagcacag aagtcattccc 180
tctggctgtc acctctctca tgcctgtctt gcttttccca ctcttcaga ttctggactc 240
caggcaggtg tgtgtccagt acatgaagga caccaacatg ctgttcctgg gcggcctcat 300
cgtggccgtg gctgtggagc gctggaacct gcacaagagg atcgccctgc gcacgctcct 360
ctgggtgggg gccaaagcctg cagcgctgat gctgggcttc atgggcgtca cagccctcct 420
gtccatgtgg atcagtaaca tggcaaccac ggccatgatg gtgcccctcg tggaggccat 480
attgcagcag atggaagcca caagcgagc caccgaggcc ggcctggagc tggtaggaca 540
gggcaaggcc aaggagctgc caggagagtc agtgattttt gaaggcccca ctctggggca 600
gcaggaagac caagagcgga agaggttgtg taaggccatg accctgtgca tctgctacgc 660
ggccagcatc gggggcaccg ccaccctgac cgggacggga cccaacgtgg tgctcctggg 720
ccagatgaac gagttgtttc ctgacagcaa ggacctcggt aactttgtct cctggtttgc 780
atttgccctt cccaacatgc tggtagtgct gctgttcgcc tggctgtggc tccagtttgt 840
ttacatgaga ttcaatttta aaaagtctcg gggctgcggg ctagagagca agaaaaacga 900
gaaggctgcc ctcaaggtgc tgcaggagga gtaccggaag ctggggccct tgtccttcgc 960
ggagatcaac gtgctgatct gcttcttcct gctgtcatc ctgtggttct cccgagacct 1020
cggcttcatg cccggctggc tgactgttgc ctgggtggag ggtgagacaa agtatgtctc 1080
cgatgccact gtggccatct ttgtggccac cctgctattc attgtgcttt cacagaagcc 1140
caagttaaac ttccgcagcc agactgagga agaaaaggaaa actccatttt atccccctcc 1200
cctgctggat tggaaagtaa cccaggagaa agtgccctgg ggcacgtgct tgctactagg 1260
gggcggtatt gctctggcta aaggatccga ggcctcgggg ctgtccctgt ggatggggaa 1320
gcagatggag cccttgacag cagtgcctcc ggcagccatc accttgatct tgtccttgct 1380
cgttgccgtg ttactagagt gcacaagcaa cgtggccacc accaccttgt tctgccccat 1440
ctttgcctcc atgtctcgct ccatcgccct caatccgctg tacatcatgc tgccctgtac 1500
cctgagtgcc tcttttgcct tcatgttgcc tgtggccacc cctccaaatg ccatcggtgt 1560
cacctatggg cacctcaagg ttgctgacat ggtgaaaaca ggagtcataa tgaacataat 1620
tggagtcttc tgttgttttt tggctgtcaa cacctgggga cgggccatat ttgacttggg 1680
tcatttccct gactgggcta atgtgacaca tattgagact taggaagagc cacaagacca 1740
cacacacagc ccttaccctc ctacaggacta ccgaaccttc tggcacacct tgtacagagt 1800
tttggggttc acaccccaaa atgacccaac gatgtccaca caccaccaa acccagccaa 1860
tgggcccact ctccctccaa gccagatgc agagatggtc atgggcagct ggagggtagg 1920
ctcagaaatg aagggaaccc ctcatgtggc tgctggaccc atctttccca agccttgcca 1980
ttatctctgt gagggaggcc aggtagccga gggatcagga tgcaggctgc tgtacccgct 2040
ctgcctcaag catccccac acagggtctt ggttttcaact cgcttcgtcc tagatagttt 2100
aaatgggaat cagatccctt ggttgagagc taagacaacc acctaccagt gcccatgtcc 2160
cttcagctc accttgagca gcctcagatc atctctgtca ctctggaagg gacacccag 2220
cca
2223

```

<210> 2
 <211> 568
 <212> PRT
 <213> Homo sapiens

<400> 2

```

Met Ala Ser Ala Leu Ser Tyr Val Ser Lys Phe Lys Ser Phe Val Ile
 1          5          10          15
Leu Phe Val Thr Pro Leu Leu Leu Leu Pro Leu Val Ile Leu Met Pro
          20          25          30
Ala Lys Phe Val Arg Cys Ala Tyr Val Ile Ile Leu Met Ala Ile Tyr
          35          40          45
Trp Cys Thr Glu Val Ile Pro Leu Ala Val Thr Ser Leu Met Pro Val
          50          55          60
Leu Leu Phe Pro Leu Phe Gln Ile Leu Asp Ser Arg Gln Val Cys Val
65          70          75          80
Gln Tyr Met Lys Asp Thr Asn Met Leu Phe Leu Gly Gly Leu Ile Val
          85          90          95
Ala Val Ala Val Glu Arg Trp Asn Leu His Lys Arg Ile Ala Leu Arg
          100          105          110
Thr Leu Leu Trp Val Gly Ala Lys Pro Ala Arg Leu Met Leu Gly Phe
          115          120          125
Met Gly Val Thr Ala Leu Leu Ser Met Trp Ile Ser Asn Met Ala Thr
          130          135          140
Thr Ala Met Met Val Pro Ile Val Glu Ala Ile Leu Gln Gln Met Glu
145          150          155          160
Ala Thr Ser Ala Ala Thr Glu Ala Gly Leu Glu Leu Val Asp Lys Gly
          165          170          175
Lys Ala Lys Glu Leu Pro Gly Ser Gln Val Ile Phe Glu Gly Pro Thr
          180          185          190
Leu Gly Gln Gln Glu Asp Gln Glu Arg Lys Arg Leu Cys Lys Ala Met
          195          200          205
Thr Leu Cys Ile Cys Tyr Ala Ala Ser Ile Gly Gly Thr Ala Thr Leu
210          215          220
Thr Gly Thr Gly Pro Asn Val Val Leu Leu Gly Gln Met Asn Glu Leu
225          230          235          240
Phe Pro Asp Ser Lys Asp Leu Val Asn Phe Ala Ser Trp Phe Ala Phe
          245          250          255
Ala Phe Pro Asn Met Leu Val Met Leu Leu Phe Ala Trp Leu Trp Leu
          260          265          270
Gln Phe Val Tyr Met Arg Phe Asn Phe Lys Lys Ser Trp Gly Cys Gly
          275          280          285
Leu Glu Ser Lys Lys Asn Glu Lys Ala Ala Leu Lys Val Leu Gln Glu
290          295          300
Glu Tyr Arg Lys Leu Gly Pro Leu Ser Phe Ala Glu Ile Asn Val Leu
305          310          315          320
Ile Cys Phe Phe Leu Val Ile Leu Trp Phe Ser Arg Asp Pro Gly
          325          330          335
Phe Met Pro Gly Trp Leu Thr Val Ala Trp Val Glu Gly Glu Thr Lys
          340          345          350
Tyr Val Ser Asp Ala Thr Val Ala Ile Phe Val Ala Thr Leu Leu Phe
          355          360          365
Ile Val Leu Ser Gln Lys Pro Lys Phe Asn Phe Arg Ser Gln Thr Glu
370          375          380
Glu Glu Arg Lys Thr Pro Phe Tyr Pro Pro Pro Leu Leu Asp Trp Lys
385          390          395          400
Val Thr Gln Glu Lys Val Pro Trp Gly Ile Val Leu Leu Leu Gly Gly
          405          410          415
Gly Phe Ala Leu Ala Lys Gly Ser Glu Ala Ser Gly Leu Ser Val Trp
          420          425          430
Met Gly Lys Gln Met Glu Pro Leu His Ala Val Pro Pro Ala Ala Ile
          435          440          445
Thr Leu Ile Leu Ser Leu Leu Val Ala Val Phe Thr Glu Cys Thr Ser
450          455          460
Asn Val Ala Thr Thr Thr Leu Phe Leu Pro Ile Phe Ala Ser Met Ser
465          470          475          480
Arg Ser Ile Gly Leu Asn Pro Leu Tyr Ile Met Leu Pro Cys Thr Leu

```

<400> 3						
ttcaaccatt	gtggaagaca	ctgtggcgat	tcctcaagga	tctagaacca	gaaatatcat	60
ttgaccacgc	aatttttatta	ctgggtatat	acccaaagga	ttataaatca	tgctgtcata	120
agacacatg	cacatacttt	acaatatgcaa	agacttaaaa	ccaacccaaa	tgctccatcaa	180
tgtagatagt	gataaagaaa	atgtggcaca	tacataccat	ggaatactat	gcagccattata	240
aaaataatga	ggtcattgtcc	tttgcaggga	catggatgaa	gcttgaagcc	atcattctca	300
gcaaactaac	acaggaacag	aaaaccaaac	accacatgtt	ctcagtcata	agtgaggagt	360
gaaccagtga	aacgcattga	cacaggggagg	ggaacatcac	acacgggggc	ctgtcagggg	420
gttgaggggc	aaggggaggg	agagcattag	gacaaatacc	taatgcatgt	gggtctctaaa	480
acctaataat	ccggttgata	gctgcagcaa	accacatagg	cacatgtata	ccatgtgaac	540
aaacctgcac	attctgcaca	tgtatcccag	aacttaagtt	aaaattaaaa	aaaaagaaaa	600
gaaaaaagaa	ctgaagtgtg	ttacttgctc	tcattcatgc	atcccggaga	aaaaggtttg	660
agtgcacatc	ctggattagg	tactagaaaa	ggcactagct	ggacaggtgg	tgatgaataa	720
acacagacat	aaatagaaat	tacatcataa	taatgtgtca	tatattttaa	aatagctaca	780
agatatctta	aatgttctca	ccacaagaaa	atgacaaaata	ttggggccag	acgcggtggc	840
tcacgcctgt	aatcccagca	ctttggggaga	ccgaggttggg	cgtgactacct	gaggtcagga	900
gttcgagacc	agcctggcta	acatggtgaa	accccatctt	tactaaaaat	gcaaaaaatt	960
agccgggctg	ggtggtgcac	acctgtaatc	ccagctactt	gggaggctga	agcaggagat	1020
ttgcttgaac	ctaggtggca	gaggttgacg	tgagccgaga	tcgtgccact	gcactccagc	1080
ctgggtgcaca	ggagcacaca	tctgtctcaa	acaaaacaaa	aaaaaacaaa	aacaagagaa	1140
atgataaata	ttcgagtgat	aaatatgtct	attagcctga	tttgaacaca	ccacaattat	1200
acacacattg	aaaaatcaca	tggtaccccg	taaataataga	caatgatttg	tcaattaaaa	1260
atgaaataac	acttaaaaaa	taaaaaagta	aaaagtaaaa	attacaccaa	taaatataag	1320
aggtacaaat	tgtgtctaagt	gccctgggga	cacaggaaaag	gcgggaaaac	ccagggctat	1380
atgatcatgag	gttacaagg	gaaaaggaca	ggagggaagg	aattgcagca	ggggcttggg	1440
agaatgcatg	tccttgggtg	caggtcacag	gaagggaaga	atgagcttga	ttcaggtagt	1500
gttgaaatct	cgggccgaga	cacgtccagt	ctgcggaagg	ctggacatct	gggactctgg	1560
catcatggct	gggttgaaag	cagaggatgg	taatcactaa	ggagccgctg	gtggttaggc	1620
ccacgacatg	gatgagactc	cccaaaaagga	agctgcagaa	tgagaggcag	gcagaggaga	1680
ggaaaagaga	aaatcacaga	ggtggggatg	tctttgtcat	cgtgtgtctc	cagtgcccaa	1740
aacagggcct	cgcgggagaag	aggtgtctcg	cacctgtctg	ttgccttggc	ggctgaatga	1800
atacatgggc	gactgtctca	gtgtgcctt	agtttgtctc	cttctctctc	agagctccgt	1860
ttcctctctg	cctgggtcgg	gcgggcagct	gcggtcgtcg	aggctcgtgt	gggcccctcc	1920
aagacgcgtg	tccgcactct	cccgcggggc	gtctgcgggg	tgcagcgtcc	actggagcgc	1980
gacagccctc	gggacagag	aggacagttg	cctcgtcttc	ctgtgcgatc	gcccaggagc	2040
tccggggcgg	agagtgcgag	cggggaaaag	gggtcctgca	cctagatggt	gcgggcagtg	2100
gcgaggaagc	caggggggac	cggggaagca	ggcccgcggt	gcggagggcg	cggggcgttg	2160
ggggacacct	ctcggagaga	caccggaggg	gcggaagtaa	ggagatgaa	aggagaggga	2220
gatcggggag	atagacctga	gagaccgaga	ggcctgcaga	gagtttcatc	cgggaccctt	2280
cagagcccgag	gaaagacaga	atgcggacgc	gggagggcgc	cttactgccea	agcgggcagc	2340
accagtgaac	aaaacacgcc	ccgcttggca	gccccgggac	gcacctctgc	ctcggcagcg	2400
caggagaggc	ttggacagcg	cgagatgcta	gggccagggc	tgcccctaga	gggctggccc	2460
gaagcgtttg	agtccaaaag	cgctcccac	cgccgcgggg	tggcagaatt	gggggcaggc	2520
gcgtccccaa	gaccccgagg	ggtggcccgc	ccccaggggc	gcggggaggc	gccccctgtc	2580
ggggcgggagt	tgctaccggc	ccctcccaca	tccccgggga	ctgtggcccc	ttcttaagcc	2640
cgcgggcgct	ctagctgccc	ctcactcgtc	tgcgccgcca	gtctccctcc	cgcgcatagg	2700
cctcggcgct	gagctatgtc	tccaagttca	agtccttcgt	gatcttgttc	gtcacccgcg	2760
tcttctgtct	gccactcgtc	attctgatgc	ccgccaaagt	cagttgcata	tccaggcagc	2820
ccttcggaca	cccgcgctgc	tgtgcccact	aacgggcacc	gaccccgga	gccctgagct	2880
ggagcgcacg	gatttcgcgg	tgagcacagc	ttctcccaag	cgccgcagct	cagaagctcc	2940

aggtgccgga	tgggaggtgc	cctgtaaaga	atctgagggg	catggcgacc	ccagggcgca	3000
ccacccttgg	ggtttacaga	tcccagggcg	caagagccgt	ccaggcaagc	acggaacact	3060
cgaaagtgagc	acagatctca	gccacacaga	tcccagcctt	aggctcagcc	cctggctccg	3120
aatcgaatct	cccacagtgc	ataaccctgt	ttccccccaa	aatgccacct	gcgccaacag	3180
ggaacctggg	agcttgccctt	tccctctctc	tctcctgtct	tttcccttcg	ccaaagaaga	3240
cttcaagctg	taggtggctt	ctgccgtcag	gagggaccta	caggaaaaaa	atcatcacc	3300
acgtggatcc	tgcgctgtct	ttgccactct	ctggcccttc	cttgggcctt	agtgtctcta	3360
tctatgatcc	acattccttc	caacctggag	agccacatct	gattcataat	cctgctccag	3420
gctgcaggca	ggttggggtt	ggcctgcttt	gcctcctgcc	tgctggggct	gtagcaggag	3480
gcgggacaca	ttcccagagc	tcgcagcctt	gggtggcagg	acctggagtt	gcagggaagc	3540
ttcctcccag	gccctagtct	cctaattgctt	ctgtgaggga	gagagagaat	gaatggcctt	3600
ggccgcaggg	tgggcgcagg	ctccactggg	ctgtgcacag	ccagtttggc	ggaggcccaa	3660
gccttttgaa	gccttttgtg	gctgctggct	gctccttcc	cgttccctct	ttcagccctt	3720
tcactctcag	cccagacagg	aaacctccag	ctccccacct	cccccccca	ggcaggtttg	3780
ggaacacagag	gagctcttca	ggggatgtct	tgggggggag	ctctagagga	agggagtga	3840
ctgggggtgtc	aggaagccaa	cctgcaagag	aactggactt	ccaccattct	gtcatctgtg	3900
tgacttttagc	caagtgcctg	tgtctctctg	gccttgggtg	tctcatctgt	acaatggggc	3960
taggtgggtt	ggttctgaca	ctttaaggct	ttgggaatca	aaagggatta	acgttagttc	4020
ctaggatggg	ggaggggag	actgggagga	gcccttgggt	gggcctagca	caggcccttg	4080
atgggtcaag	gacagcagat	ccaactgtgg	aggctgtgca	gttgccacac	caactgtggg	4140
cagccacact	tcctctgtag	caagaagtct	gggggtgtta	ttgtccaggg	gaagtagcca	4200
ggcaggagag	ctgcgatttc	agctctgtct	gtagggagtg	atgttccctg	gaatgattat	4260
agtagcttgg	ctgaccttcc	tgccacagga	gacccactc	acacacacac	acatacacag	4320
tctctctctc	acacacacac	actcacaac	acacacagac	acacacacaa	acacacagac	4380
acacacaaaa	cacacgcaca	aacacaaaca	cacacaaaaa	cacacgcaca	aacacacacc	4440
caaacacaca	aacacagaca	cacaaacaca	cacacaaaca	cacacacaaa	acacacaaaa	4500
aacacaaaaca	cacacataca	catacacaca	cacacacaca	cacacctga	actggaaacc	4560
ctaactcagt	gtgtgtgtat	gtgtgtgtat	gtgtgtgtgt	gtaagagaga	gagagagaga	4620
gattaagctg	tcctttgagt	gaggaccagg	gaggggaaga	agagaaccca	gggagagtc	4680
ttccaaaggc	tgccttcacg	agctttcctt	ctggcggggg	tgggtgagga	ccctggacct	4740
tgtcttcttg	ttttttccct	ttctgcctgt	tttggtcacc	ctgcccccac	cctccatggc	4800
cgccccattg	tgcaaggaaa	cccagagggg	acacagcacg	ggcaggggcag	ctgggaagct	4860
ggtgagaagc	tgggaggacc	ttggcagcct	gagcaacaca	gtccttgcca	ggaggtagct	4920
ccaggggcac	gccaccctct	gccaacaccc	aggcctctct	cctcacggac	tgtctccagt	4980
tttctgtctc	ccacctggat	tccctcctgg	cctcatctct	gtccactct	ctctatcctt	5040
cctctgggtc	tttttttaat	tgaaaaaaaaa	tttaatgaaa	taaattgatag	atttcttgta	5100
tcacttattt	tattaaaatg	taaaagggtt	cttttttgca	aatctgtaag	atataaagta	5160
aaaataaaag	tacactcaaa	tcccataagt	tattcacatt	ttgatgaaca	tctttccaga	5220
tgaatctctc	tctctcttcc	cagacacaca	cacacacagt	cacacacagt	aggttttgcc	5280
tgcatttttt	cattaagtgg	tgtgtcagga	cacctgcct	tgttaatgtt	gaacttttct	5340
aacatccgct	tcccacctg	ccctctccct	tgacactgtg	gaggcattct	agactagggg	5400
ggtcaagcct	gttgacttca	gggatgaggc	acctcctggg	cttctaata	gtggcgcgga	5460
ggtgaggggg	cagttaacct	tgtgtctcgt	cctctttcct	agtgggtctg	cttgactcct	5520
ccaggaacgc	acagtgtaca	ttgggtgacg	acgccacct	ctgcttctaa	gtttagagaa	5580
tcaaaagtta	ccgaggactt	tgtgcgccat	atgggaagaa	tgagcactct	taaatccacg	5640
atttgcagat	gaagacatga	aacaagaggg	gacagggacc	aggattggga	gcaggaggag	5700
taatttatga	gcgacattgt	ttagaattgc	tatcacttga	tgatagtaag	aagcaaaact	5760
atttttagct	aatattattg	ttttaaaatt	ctctccaatg	cgccctctca	tgtgtgcce	5820
ctggaggcat	cattctgatg	gctgcgccag	ggtacacccc	gacaccaagc	cccaaggga	5880
ttagtggctg	ccaaaggcca	gacagtggct	gacagtgggc	gccaatcata	tctgtctggg	5940
gtcaaagcct	gggcttccag	tcacgctgct	gttccgcctt	tagttcagcg	gctgtcaacc	6000
agcattgacc	ctttctcctg	ccctcaccct	gccccacaa	aggggaactt	tggcaaaagta	6060
cagagacatt	ttttgctgtc	caacctggag	acagtcttac	tggcatctca	taggtggagg	6120
ccagcggtgc	tctaaacacc	ctgcagtgtca	cagctcccac	aacaaagcat	catttagccc	6180
aaaatgtcag	tgtgccgagg	ctgagggacc	ctgcctccca	gtaggggagt	gccctgggtt	6240
gctcgtggga	tgtgaaaaa	agattttatt	ttttgtggct	gataacacaa	ccctgacaaa	6300
gaattttcaa	gtcttctctg	actgttttgt	gcaaataata	catacgctct	tctgggtgat	6360
gagaagcagg	gattgtgtac	aggtgcatct	gttcttcagc	agcattgtca	gagttaaact	6420
cagatgaatg	ctattgattc	ctttaataaa	catttgcaaa	gatggccggg	cacagtggct	6480
catgcctgta	atcccagcac	tttgggaggc	cgagacaggt	ggatcacgag	gccaggagat	6540
caagaccatc	ctggccaaca	tggtgaaatc	ccctctctac	taaaaaata	aaaattagcc	6600
gggcgtgggt	gcgcttgtct	gtagtcccag	ctactcagga	gactgaggca	ggagaatcgc	6660
ttgaacctgg	gaggtagagg	ctgcagtgtg	ccaagattgc	accactgcac	tccagcctgg	6720
ggacagagca	aggctctgtc	tcaaaaataa	gtaagtaagt	aaataaataa	ataaataaat	6780
aaataagcaa	gaatttgcaa	agatatccta	agtgttgggc	ctgttctgga	tgtgtaggac	6840
ggtgatctac	aaatacagca	ggttcttgaa	taatgttgat	tcattcaata	tcatttcatt	6900
ataatgttga	tgaggggagg	aaaaaaaaag	aaggatccct	tgagccagg	agatggagg	6960
tacagtgagc	tgtgaccgtg	ccacttcact	cccacctggg	caacagagcc	agaccctgtc	7020

```

tcaaaaaaa aaaaaaagaa ataaaagagc gagagagaaa gaaaagaaaa tgattactgg 7080
ctggggccac tgtctgtgtg gagegtgcac attaccctcg tgtccacatg gcttttcttt 7140
ggctagtatg gtttccttcc acatcccaaa cccgtgcacg ttaggtgaat tggagtgctt 7200
gtatgggtccc tgtctgagtg agcgtgggcg tgcgtgtcag tgtgcattct gcaatgggat 7260
ggcatcttgt ccagggctgg tttccacctt gtaccctgag ctgccgggac aggatctggg 7320
cacccaagac cctgacctgc tgtaactggg taaataatta tctaacttgt tttcaatggt 7380
tcttaagtat atgtatagct cacattccct tcagtgttta atattggaag tgttttgggtc 7440
tttatttaag atctccgtga tgtttttgtg accagaaata tgctgtagaa atttaactgt 7500
tgtttctagc aatttgcccta tgggaatatt ggcttatggt gtttcgctta cgcattgcaa 7560
tttccaaaaa ccaatcaatg atgttaagtg aggactcact gtactgtttg tgctttcgag 7620
tcacgcactg gttgtggtgg tagaaggaca gttgaggaaa cagtgacaac tccatagtct 7680
aatggctggg gagggtactc agaggaaggg cacaaccag actatagaag aggcgcaggg 7740
agacatctaa gaaggaactc tgaggttggg cgcggtggct cagcctgta atccagcac 7800
tttagggtagc tgaggtgggc ggatcatgag gtcaggaggt cgagatcagc ctggccaatg 7860
tggcaaaacc ccgtctctac taaaaataca aaaattagca gggcgtggtg gcaggtgcct 7920
gcaatcccaa ctccggaggt tgaggcagga gaatcgctt aacccgggag gtggaggttg 7980
cagtgaagct agattgtgcc attgcactcc agcctgggca acaggatcga aactctgtca 8040
cacacacaca cacaaaaaat actgatgaaa cataaaacaa cctagggagg tggctagttt 8100
tatcacataa ttattattac ttttatttca atagctttag gggtacacgt agttttcgtt 8160
tacatgaatg aattggatag tggtaagtc tgagatttta atccctccct cctatcccac 8220
cctgtctgct tctaagtctc cagtatccat tgcaccacgc tatatacctc tggataacca 8280
tagcttagct ccactttata agggagaaca tgcactatth ggctttccat tgctgagtca 8340
cttctcttag aacaatggcc tcttagggcg caagagcgac actccatctc aaaaataaaa 8400
taataataaa accaaaaaaa ccagggtattt tattcttctt ctcttctctc tcttctctct 8460
ttcttttctt cttcttccat ccccttctct ctcttctctt tatccctctc tctctcttct 8520
cctcttctct cttcttctct ctttcttctc ttgtctctat cttgatcttt tcttttgaga 8580
ggcagctaata ccaagggttg agaagatgaa agaactgtcc tagaaccaca cagctgggaa 8640
ggaggggaggc agggaggagg ggtgggaatg gggcaggagt cctttgcgaa tagatccctg 8700
gcctgaccgg ggaaagctgt gctgaccagg gctggggaac aagatgactt tgaggggaat 8760
ccctctgaga tcagcactgt gctttgacaa tccatgccag ccgccgtccg gagtgttctg 8820
ggggtgggga gaggaggcg gcaacacgct gaggcctcag gactgtctct tcagtttgtc 8880
aggtgtgcct acgtcatcat cctcatggcc atttactggt gcacagaagt catccctctg 8940
gctgtcacct ctctcatgcc tgtcttgctt ttccactct tccagattct ggactccagg 9000
caggtgagca gaccaagggt atcctggtga ctttctggtt ctccctctct ctttctctct 9060
agtcccaact gtgagtcgca caggccctgg ggtgaccgga aaacctcat ttgtggattc 9120
tccctggcag ggagacacca ctcgagcctg catccccact ccaagctgtc cctgaagtca 9180
gcatctgggg actgggtggc tctagtgtgt ggcaaggagc agtcctgatg aggccttctg 9240
gccacgctcc aggtgtgtgt ccagtacatg aaggacacca acatgctgtt cctgggaggc 9300
ctcatcgtgg ccgtggctgt ggagcgctgg aacctgcaca agaggatcgc cctgcgcacg 9360
ctcctctggg tgggggccaa gcctgcacgg taattacgcc ttctctctct tgccagctgg 9420
ctctgcata gccccagggc tggaaagggg tggaggatgg cacagaccag gccatccact 9480
ggtgaaggct ggccatgggc ttacctggac ttggctgggt ggggtgcagt tatagcttta 9540
gtgggagaga ccagatgcat gcgtggtggt ggcacatggt gaggcagcag aagtaagggt 9600
cctcgaatcc agaggaggtg ggtcagcaag agtccttgca ggcttggaag gctttctggg 9660
ggaggcagct agctgcagg ttcacccggg aacaaattgg atagaggctg gatcaagctg 9720
tgtctgata gataaggga gcaggccaga agtggtctca ctaccagct catggggaag 9780
cagaagggtc ctctctccaa gctggagcat ctattcccac tgcaaagaag cttcttatct 9840
tccccgatat cactcagtac ccagcttct ctctccattt ccaggatctc tctgccaat 9900
ctagctagcc atttccagct aagccatgga gtcaatataa tcataatcat aaccataatc 9960
aatcatgata ataatgggta tattgagtg ctacaaggcc ccaggcatga taccaggagc
10020
ttatgaattg cctcatthaa ttcttaccac aactctgaga gctgagtatt cttactgccc
10080
acattctgtg gatgagggat tggaggcaga gagggataaa gtgatttgct catggacaca
10140
cggggattgg acccagcttc tctgataagg cctgtgtcct ctctaatacag aaactcaggg
10200
catatcttcc ttttgagaca atgtgtcccc tcaatgatgg cacgtccttg gccagccat
10260
caggagtcag ctgctggtca gtttacggta aattcctctt gaggcgcccc tgtgtcaggg
10320
gctgtgctag accctgagca cacacagacg ctagcctgcc ctgcaggcaa ccatgccccg
10380
gggctccctt ggtgcctcag attcccctga gcaggggact ttgtggctct ctgatctgtt
10440
acacatcctg gcattgattc cttccagtaa ttgcttttagc atcaaatcaa aagccatcat
10500

```

attttctaga aatgagagac cccaggaaaag tggacctcag ggcctcaga attcttctgc
 10560
 ttggctccct tgagtggcca gcttgggtgg gaggccactc cagtgggttt cattctgcag
 10620
 catgctggag agcttccact tccaaacca agttcacaca tgcttctgta tccttctctgc
 10680
 caccctgctc ctctgagtat ggtctccggt tgtccaaggc actgcctgtc ctgggagtca
 10740
 cctgtatgtg aggcaaccctt ggtgccttga gatatcatgt agaagccttg gttcttctca
 10800
 gacaactcca ttcatgcaaa ctctccccct cctcctagcc tgggtcccgg gctttgtttt
 10860
 tttttgggtc cataatgtct gcctgtgtgg acagcagctt gggccctggg gcagaacagc
 10920
 tcctaggtcc cttcttcagg ctctacccc tgcccctgct cctaccccca ggtgaattag
 10980
 gagccctgag gaggagcctg gctgcagcga ggcccacaga ctgagagtag ctgagctcct
 11040
 tctgtcccta gccttggaac gctggggcat gtagagccac agagcagagt caggccctgc
 11100
 cctgctcaca gccaaggag agagcagaca tggaaacagc tgctttgaac ccagcacagc
 11160
 gatgattaga gtatggggaa ggattgagaa gggtcaggcc agccccacct ggtgcacaca
 11220
 ctgagagcgt ggtcccagag gagggatgtt gtttgagcag gctctgaagg accatgagga
 11280
 gtcttctga tagacagcag agaaggagc aggggttaca agcaaaggga gtgtttcttc
 11340
 tgaatactgt ttgtgtgata gctccactgc agcatggagg ggtcaaagtg tatgtgcggg
 11400
 gcggagggga gatggcaggg ttggagtggc agccgggaga atagtcacac tttcccaagc
 11460
 tccctccca gctcacccta cccctactct gcttagccct tctgaacttc tgagaggtgc
 11520
 aacagagttt gggggtgggt gggaatttcc tagccagaag tgggaagctg gggctgcctg
 11580
 cacatagggg tattccagca caccctaggg caagctcata ttgagttggc accatctgga
 11640
 tgcctgggct tcccctgcta gatgggtggg cagggggtgct ccttagaacc acgactggat
 11700
 ctgaggcctc ttggtaaccc cagaagcaag cagagtagac atcagtcagc ggtgtgggag
 11760
 aggcaggagg gagagaggaa tggaggaaagc aaagaaggga aggaggaggg gaggggaggc
 11820
 tctaaaaccg tcatccctat tccaatatct gatcttgaat tggcctcaac acctgtgcat
 11880
 ccctgcaggg gtggaccag tcccagttg ctcccaggg agtacggggg tggggtggg
 11940
 attctctggc tttctccct gcccctctc tgcaggctga tgctgggctt catgggcgtc
 12000
 acagccctcc tgtccatgtg gatcagtaac acggcaacca cggccatgat ggtgcccac
 12060
 gtggaggcca tattgcagca gatggaagcc acaagcgag ccaccgaggc cggcctggag
 12120
 ctggtggaca agggcaaggc caaggagctg ccagggtgagc ccctggccag ggcactgcca
 12180
 ggccacaaca gcagccttcc cctccctctg ctggcaaatg ctttggccac ctcttctcc
 12240
 ctgtctgctt cccggagccc tcccttaaac acgcatagag aaaaaaaaaat agaaaatact
 12300
 gttgtcctaa gttttaggag gggattattg cacacaactt agatccttta atagagcttt
 12360
 gaacaaagtc tcaccctcag ttcccatcag ttgcagaaat cagtgtgttc acctgattat
 12420
 tcatttgggc atctttcgag cacttaggga tgcccctcac tccttgctac tcctgctcat
 12480
 cctcaaggag gccttttctg acctcctcga gcagctcaaa tccttccact ctctgctccc
 12540

ataggctctgg ggcttggcgt cccatgcttg cttccctgct aggtgcgaag ctcaggggaag
12600
acgagtcagc atctaccttg ccgtctgccg tgttccctta ccatccccag cccagtgcag
12660
tagagtcagg gtctgtggct gacggcctga ttgccagacc ctgggcaagg tcctggggct
12720
tacagagagg aatcggggcac atccctgccca gcaactctta tggagcccag tggggcagct
12780
aaatcagcag agctgggatt tcccaatcct caggtcagca gcagagtcag gacctggggc
12840
tgggtgggca gccccatga ctggctcagc taacagcgct gtgcccacca caggagtcga
12900
agtgatTTTT gaaggcccca ctctggggca gcaggaagac caagagcgga agaggttggtg
12960
taaggccatg accctgtgca tctgtacgc ggccagcatc gggggcaccg ccaccctgac
13020
cgggacggga cccaacgtgg tgctcctggg ccagatgaac gagtgaagtcc ttggtcgac
13080
cttctgggga caacgaagtg ggtaccgggg ctggagggac ctgcccacct ctctctgctc
13140
ctctgcagag tcctggaaaag cctcggggca gccagacctg gcctggggagc ctggcagggg
13200
tggaaagatg tggccccatc tagcctctgt gtccctggcac ccctgtgccc acacagaagc
13260
cttagagagg ataggagct gatgtcaggg gagctaactc cccagtctgc tttctgctat
13320
gatgcaagac ccaccacctc ccctggggtc agggactctg gctcagagag ggagtgtgga
13380
ttgaactctg agctaaagtc atggcagatg acaatgtact tccagacgct gggtccttgg
13440
ttgaaacttg tagaaaatag acacctctaa aagactcccc agcactccct ttgctcactg
13500
cttttggtgg ctaatggtga tggccccatg gcatccgagg tctacagatg gtatgaaggg
13560
ctgggggttg gtcattcact gcttactgc ttcgttatag tccccttggt aggtatcagg
13620
tgaacctagg gatggtttgg aactttctag ccttgggcac aaagggatgc aggccatgag
13680
gaccccaaga gggagagaaa cctggggcct gccgcggggt agtcatggtc tgttgagggt
13740
ggcaagatgc ctggggcttc caggcatgtc tgggtacataa atgtactaat tgaggtatgt
13800
actaattgca gtgggcaggc aaaaaataa ggtgatgccca tcctttgcag acaggagcct
13860
ggacaggggt ggggagggca gtgggcgcag gagctgggag tgggaaagga caggctctgga
13920
gcctggctgg gcagaaactg gaggttcaac aaccggttg ttttaatttc gggagtgttt
13980
tctgtaatga tatccttaca gttctccagt aactttcttt gggaagagca gcccgctctg
14040
gctgagtggt gaaagctctg cgctgcttt gacactcttg agctaaaggg ggcgccctg
14100
gggctagcag agccccgggg atgggaggcg gggcctgtgg tggaaagtac cctcctccag
14160
cctccgctct gggaagcttt tgagatttcc tttgctaagt ggggggaccg ttctttgcag
14220
aaaccacag agcgagattg ctgaggtctc tgcagatccc caaagatgac agccaaatta
14280
catgcatgtg tataaaagggt gtatttttct ttttttctt tttgagacaa gtctcgctct
14340
gtcgcccagg ctggagtgcg gtggcgcat gttggctcac tgcaacctct gcctcctggg
14400
ttcaagcgat tctcccgctt cagcctccct attagctggg attacaggcg cccgccacca
14460
tgccctgtta atttttgtat ttttagtgga gacgggggtt caccatgttg gccaggccag
14520
tcttaagctc ctgacctgtg gcccacctg cctcggcctc ccagagtcct ggaattacag
14580

gcgtgagccc ctgcgcccgg ccacaaagtt gtatTTTTtct ggagggatgg gccataactt
 14640
 ccatgagact cttagcaagg cctggacaca cagaagagtc agtgggtcat ttctcggcct
 14700
 tgtcttgtgc tgtggccatg ttctgaggct cccactcgat taggggacaa tgcttgccaa
 14760
 tggacttggg ggctagacct caggaggatg tggcctccac acaggcgcg cctctcagggc
 14820
 ccagctgctg ctccgtcccc acgcacaggg ccaggctggc tcccacagct cagcatctga
 14880
 ggtggggggc ggtgtcttct tgtaggttgt ttcttgacag caaggacctc gtgaactttg
 14940
 ctctctgggt tgcatTTGc tttcccaaca tgctgggtgat gctgctgttc gcctggctgt
 15000
 ggctccagtt tgtttacatg agattcaagt aagtttgagc tgctcacagc ctaattatgc
 15060
 ctaattatgc ctcaaagctg cagaagagcc ctgagactca ataggcaggt ttacaaagtc
 15120
 ctctgtgtct ggccctgatc tttctccagc cctgtctcct gctagtctgc cctctgttc
 15180
 ctctgaaccc aggctgctca ctgagctttg tgcacacgtg gtcccccttc cctggaatgc
 15240
 cattctctac ctccacacct cctcagcctt caaggctagt tcaaagtctg ctccctgac
 15300
 ttttccccac cccattcca tctctgagcg gcccctgggc atatcacagg cctgtccttt
 15360
 agtatctgca tttggcttcc ggtgactttg aattcctcca gaaccactct gatgctgggc
 15420
 accccgcaca gctcccagca caggggaggaa gagcaggcag gttaaagcaa ttaaagataa
 15480
 gctgggtccc acgtgccagt tcgacattgc tggacaagct tcctctttgc cgtgtgggtc
 15540
 catcaggcca ggtcaccgca aacctgtgac ttagctctga gctgagcgca tacgctctgt
 15600
 gcctcaatgc acggggagtt taagtcgagt aaaaccagca gtgattatga ccaaattcat
 15660
 ccaaaccag acatttactg aatacctctg gtgttccag cagtgtacag gtcctagaaa
 15720
 gtttaccttc ctgttcctag cacacaggca agttcatcag gggtcacctt tgatggcagc
 15780
 cagactttgg acagaaacca tgacctgtgg ctgacaaata gtaaaaaaa agttattgtt
 15840
 tttctaaaac acacaaattt atctgtggtg caaagggtgat caggccacac caggatagaa
 15900
 agtactcagc tctgagttaa gtgcctgtgc tctgtgcctc catccacagg aagttcgagc
 15960
 caagtcaaac cagggaatt tgtgaccaga ggaagagac tgcagagctc agaggcaaaa
 16020
 gtgcccacgg aaacctgtga ttttgtgggg aaaataggga attttcctaa gttttctct
 16080
 gaaggaggaa ctgttttgaa aactccatt aaaaagttgc tatacaggcc gggcgcgatg
 16140
 gctcacacct gtaatcccaa catttttga ggccgagggt ggcagatcgc ctgaggtcag
 16200
 gagtttgtaa ccagcctggc caacatggtg aaaccccgct tctactaaaa atacaaaaat
 16260
 tagccgggcg tggtagccca cgctgaaat cccagcactt tgggaggcca aggaggcg
 16320
 atcgctgag gtcaggagct cgagaccagc ctggccaaca tggtgaaacc ccatctctac
 16380
 taaaaataca aaagttagct gggcatggtg gcacatgcct gtaacccag ctacttgga
 16440
 ggctgaggca ggagaattgc ttgaagccgg gaggtagagg ttgcagtaag ccaagatcat
 16500
 gccactgcac tccagcctgg gcgacagagc aagactctgt ctcaaaacaa aaaaaaagt
 16560
 tgctatacat attcaaaaca atcataataa tgatagtaag aatgacaata ttaatgatca
 16620

ttgccaaac cccactctgt cctgcccatg gacggggcag gggaaactgt ttgcatggct
 16680
 gcctggccac ccagcctggc tttgacagta gctctctttg ccctgcctct tgaatctgca
 16740
 ccagggccaa agtcctgttc atttgttcac atccgtcgaa caggtctctc aggagatggg
 16800
 cctgaacctg ctgcagggtga gcatctgtgt ctctcatgg ggcaacagga ataataatga
 16860
 ccaacattta ttgagtgtc atcatgtgcc agacatgatt tcgagcgctc ttttccttcc
 16920
 tttattttat tttattttat tttattttat tttattttat tttattttat tttattttat
 16980
 tttatttttg agacagtgtc ttgctctgtc acccatgctg gagtgcagtg gtatgatctc
 17040
 ggctcgctgc aacctccacc acctgggttc aagcaattcc ccctgcctca gcctcccaag
 17100
 tagctggaat tacaggcacc caccaccacc atgcctggct aatttttgta ttttttagta
 17160
 gagatggggg tttgccatgt tggccaggct ggtcttgaac tcctaaccctc cggatgatccg
 17220
 cctccttggt cctcccaaag tgctggcggt acagatgtga gccacctcgc ctggcccaag
 17280
 cactcttaaa cttaattaat tttcacaaca acctgcgagg tcagcactat tattattatt
 17340
 cccaatttac agacaaggca actgaggcat ggagagggtga tgtggtcaac acagagcttt
 17400
 gtaacaggga agtaggggga ctgagacttg aaccaggcc ctttggtcc cactgcatgg
 17460
 catccctct tggggaggct gagggttgc gtccttagtt gcctccagac ctaagcatga
 17520
 ccagggtgca gaaacactag ttggggccgg ggctgcccta gaacccaag gcctactgag
 17580
 aaagaggagg gagatagcat ggcccgagg ccgcaaggcc accatcagct tcttgtctgg
 17640
 ccagaggcag atgtcaggcc cctggagact cacagccaga acctgaagct gagtccacc
 17700
 agcctggcac ggccttcac agcttttgc gactggcggg ggagcctgag agtgtctgca
 17760
 gcaggggggt tctgagcatg ctggtggtgg ggtgcgtggc tgcagtccag tcccaccct
 17820
 tccccttccc gacgggccac tctagtttg acgcatgcag tgtggtggc cggggtagct
 17880
 cacggcagct ttgttttggc tccagatctg gaaggtagag gacagctttt acattcggtt
 17940
 tgagtgggtg gaacagtgtc ctggcccagg ccacgtcctg ccacaaacta agacctggtg
 18000
 gtccctgcct gcctttgtgg cctcatggac ctcccacct gaggccaggg agcacctgtc
 18060
 tcagcggcag gaggcagctc cactgtcagc tgttgcctc actagagttc ctcatctgaa
 18120
 cgatcctgga gaacgaggtt aagttcttgg cctctagcct aatccagaac aactatcttg
 18180
 ctgaagagcc tagtgagcc tcctaggcta tatctagcca aaggggccag accccacccc
 18240
 aggaccacca agaactacat gggatattat tactgggttat acctaactgt cccaaccagg
 18300
 cttacctct gtaatagcca tgagggttct ttgggacccc tgccagggca gaggcattgca
 18360
 aagctcaaga atctctcccc tcttgttggc tctgcaacat attcagtcca agttcaccat
 18420
 ggtgcatcat ggtgaaggct gttctgctgc aggaggactc tgtggtcccc acccctgacc
 18480
 ctgacctagg cccctcacag gccaaactgga tccatttact tgcattctcat gccagcctgg
 18540
 tcatcaccag atgaaattaa ccagagatg agagcaaagc tgctcagcac gagagactct
 18600
 gaaggcttg cgtaccact gtggggcact ggcatggaa gactgcatac tccatgcagc
 18660

cccagagtct gcagctactg tgggtgtggg gatgagctgc cagcaccaaa tgcaggtctt
18720
ggctcctggg ccactagtaa taccaaggtc accccttatg ctggaaacct gaagcccctg
18780
gctgagcccc aggggtctcta ggacgacagt tggcagcaga gaggtgcttg gtagagcaca
18840
aactttacta agccaagggt gtggcagcag agaggccctg tcttacacca gcagagccat
18900
ccctgtgccg gatgtctaga gagtgtccct agcgggtgac cctcaggaca cacggccttg
18960
cccagcagg agatcctagc cagccgtgta gacctgaggt cccatcagtg ttgcctcctt
19020
ttctgacccc tgagcacccc agaaagctgt gacctgatgt cctgggtgtcc ccatgttcca
19080
ggccaagcca ccatcacacc aacacttggc cctcacactc tccaaggctg ttcacatcca
19140
gcactggcctt caggaatgag ctccatttcc atcaaccctt tccctcctat gattatgtct
19200
catggccccc gggaagggtt ctcacgaggg agggctctcc aggacaatac tcttggcctt
19260
gccacccctt tcaaaccaac agtggctgga actggaatgt gtgaatggaa tattcagcat
19320
accttgaggc cttagtctta tgcacagtgg cccagttat cccccctcca cagctgagct
19380
cccctttaca cctcctccaa gaacctctc tctcctctgc ctcctcatgc caacgccacc
19440
ttaggggagg ccctgcagga caccctggac aatggacact ggtcccaagg gggcccatcc
19500
aggatggggg tgccatcctg ggctgtcttc cttcttgccc tagccatgct tgctgctaac
19560
cccaggttct cctggatccc taatcctgca cctccaactc cagggaacac aaggacccat
19620
tctgcccttg actagccctg tctgccaggg ttcatactca ctccctgcat ctccctgagc
19680
caccttggtg atgggggttg gcatcccaac accatcgaag gcagctccag gctgaggtgg
19740
aaggaggaag acttggaag catgtgaggg agccctgttc ccaccttgcg caggctccga
19800
agtccttat ggcttcccc caggtgaccc tggagcagcc agtctccagg tgtctgggca
19860
cctgccgaga ccctctagcc tctctacaga gactttttcc ctagtacatt ctgggatgga
19920
agaacaggag agggaaagag gcaggaaggg cttttctcca ggcccatag caggcgagga
19980
cagcattatg tgtctttttg ctacattctg ctgtagaaca tttaggctcc atctgaccag
20040
cacctgagcc aaccagtctg ccctgccctt ctctcatctt tgcattctcc agttttaaaa
20100
agtcctgggg ctgcgggcta gagagcaaga aaaacgagaa ggctgccctc aagggtgctgc
20160
aggaggagta ccggaagctg gggcccttgt ccttcgcgga gatcaacgtg ctgatctgct
20220
tcttctgct ggtcatcctg tggttctccc gagaccccg gctcatgccc ggctggctga
20280
ctgttgccctg ggtggagggt gagacaaagt aagtcttgga ttcaatagaa atcgctggct
20340
tagggccagg cgcgttggtt cacacatgta atcccagcac tttgggaggc tgagggtgggt
20400
gggtcacttg aggtcaggag atcgagacca tcctggccaa catggtgaaa ccctgtctct
20460
actaaaaata cagaaaatta gcgaggcatg gtggcacatg cctgtagtcc cagctacttg
20520
ggagactgag gcaggagaat cacttgaacc caggaggcag aggttgcaat gagcccagat
20580
cgtgccactg cactccagcc tgggcaacag agagagactc cgtctcaaaa aaagagaaa
20640
aaagacacca ctggcttagt gcactagtgc ctaaatgctg ctggtctcgg ctacagggtg
20700

caagaggaat gtgggccagg cactcatgct tggccaagac ttttcctctt ttgggagctg
 20760
 ggtttcagag agcactctgt tggtttcatg actcattttt gttttctgac caagctccac
 20820
 aataagaccc taatgtgttc ctgtggatc ctctcctccc tgagtaggct gaggagaaaa
 20880
 tccttgcca ggcagggtg ccagagctgt gatgagagag atttcttggg ctaggagtag
 20940
 ggttcccaga gctctagttt ccaaactctt gctctgccat cttccctttc tcatcttcac
 21000
 atctggtcaa atccctccaa aggcacacat ctaggagct tcatagacag agacttgcca
 21060
 aagggggtac atgtagtttc tctcctggct aagacgttgt cagaatggaa gaaaggatga
 21120
 gaaacatgta catcctagaa aaggcagaag atgtgggcag ggagatgctg gtatgatggc
 21180
 catttcgttt tgaaggctcg cttaggctcag caccaaagtc ttcattggtca ccctggtgaa
 21240
 ccagacaga attctagaga acctgggtcaa gaagaggtcc tgaaatacac ttatggagaa
 21300
 tgcacgctga gagggggaag taaactgctt aggatcacc aaagtgtgtg gtcaagagtg
 21360
 tgggcatctt gatttctagc caggattcag tctcccatc cactcttatt tttttatttt
 21420
 tttgagacag agtctcacac tgtcaccag gctggagtgc aatggcatga tctcaactca
 21480
 ctgcaacctc cacctcccag caattctcct gcctcagcct gccgagtagc tgggattaca
 21540
 ggcgcccgc agcatgtctg gctaattttt tgtattttta gtagagacgg ggtttacta
 21600
 tgttgccag gctggtcttg aactcctgac ctctgatcc gcccgctca gcctccaaa
 21660
 gtgctgggat tacaagtgtg agccactgca cctggccacc actcttgacc ttgactttta
 21720
 aggtgtgag cctgtttctt tgcatagaag catttgga cagaactgcc ggagtgtga
 21780
 tgggtttgtt gactgactgt ctctgtcgca gatgagctgt gcttttcccc acctaggtat
 21840
 gtctccgatg ccactgtggc catctttgtg gccaccctgc tattcattgt gccttcacag
 21900
 aagcccaagt ttaacttccg cagccagact gaggaaggta agtctcctgt tctgatcgcc
 21960
 cagtcacag gactggagcc ctggaaccaa agggtcacta tgggatgcct tgggccctag
 22020
 agggagaaaa tcccatcata tccaagagga ttggctacaa aagcctggga aacagtggct
 22080
 ttcaagccac cggtggtatt atttagtgca aaatatcttt tttgcttttt aacatttgaa
 22140
 ttttaacattt gaaattttat ttatattaca acaggaacag aaaatgtttc aaattttcca
 22200
 taactgat ttccattcag cacaattttt tgttttctct cttcctccca gtctttgcta
 22260
 atatgcctgt atattacatt ataataca caacagttt gaatcctatt tgtttgtgt
 22320
 ttcttctacc acttttgatt gatattacac tataaacatt tccactatt gctacagtct
 22380
 tcaaatatat tttctcta atatggcatta tattgcgttg aggggttgta atcattctcc
 22440
 tgttatagaa cattttggct gttttgaatt ttttattttc ataaattaat gttttcttgc
 22500
 atatagcttt tcctttgagg gtattttttc tttaggataa acttctagga gtaatatgtc
 22560
 tgggtgata gaatacaag tcttaatggc ccttaaaatg tatggccaaa ttgcttttca
 22620
 aaaaggctac accaatttac gatgctattg gcagtgtgtg taatagtttg atcatatcct
 22680
 caccagcaat gtatatatta ttgtaaactt tagctaattt ataagtagga gatgtacct
 22740

cattgtcctt attagcttta ttcccccttg attagatttc ttttgtcttc taatgtgtct
 22800
 ctgggtgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtttctt tttcttccca
 22860
 gaaaggaaaa ctccatttta tccccctccc ctgctggatt ggaaggtaac ccaggagaaa
 22920
 gtgccctggg gcatcgtgct gctactaggg ggcggatttg ctctggctaa aggatccgag
 22980
 gtaactttctc cagccacagg ctgcccagag ccctcttctt cgtcaagagg gtggcgtttc
 23040
 tccacccttc catccctggg ctgtgtgtgt tctgtgcctg catccttcgt ataaccgcac
 23100
 attccttgag gacatggact ctgtcttctc atctaggaac tctacaccac acacagggcc
 23160
 tggaagacag aaagtaacct ttgagcgatt gcaggaatga gtgaatgagt gaccgtgggt
 23220
 agccaagaga ggcagaggac actgtcagtt accctctggg gcttgatcac aataatctct
 23280
 gctttgattt gtctgaggga aatctttctt tccaatcctt gtcaatattg tttgtacta
 23340
 cttttgttcc ttctactggc tacttaacat ggtagctact tcaaaatttt tctttagcta
 23400
 agtatgtagc agcgtaggag gtgaggaaca tgttgaaaaa cacacaaaaa tataactttc
 23460
 ttacctcct tctttccctc ctggggaaga aatgagccag agggagggat gagctagctt
 23520
 gctgctgctg tctccaacc aaccatctac ctaccaagt atccaggagt gtaatagaca
 23580
 gacttggctc agttattgct gtttcttcaa tatctaggac acagcctggg gcctagtggg
 23640
 tgctaagttt ttgcgagggt gaacaaatcc atccatctag tcacctctcc atccatcac
 23700
 catgcatcta ttcatgtctt catccatcca tccgtcgcgc catccgtcca tccatccatc
 23760
 catccatcca tccatccatc catccatcca tctgtccatc catccacca tctatctatc
 23820
 caccatcta tctatccatc catccatcca cccatctatc catccatcca tccatccatc
 23880
 catccatcca tccatccatc tatccatcca tccatccaac attcctatta tgtccctagt
 23940
 gttatgccag gcacagagat tacagaggag attgagatac ggtccctgtt cgtggcagac
 24000
 ttcacagact agggaggggc acatatatga aagggcattt cagggaagtag cacacgagca
 24060
 agggaaaaat gtgaggtatt tagctgagga gaagtagaag atgaggctgg taaggctacc
 24120
 agaagccact tctctgaggg cccaagata gaggggtgtg gacttgatcg tgaatgcagt
 24180
 agacagccac tgaaggactg aggccagggg gtgagttggg cagatctgca catgaggaaa
 24240
 tcactctgat gtctggagtg ggggcctggg ctgggcaggg ctggaggag aactagctga
 24300
 gactctgcag ccttccatct cactcaggct cagaactttg gactctgtgg acattctctc
 24360
 ctcctttggc cccagctca gcacagtctc cagctttact tcggactcag actattctgt
 24420
 ctcagccttc gttgctgact tctctgttct ccctgaaaca ggagtgtctg cccaggctct
 24480
 gtccttggcc tctctctttt ttacacttca ttctctccct ggacaatctc ttctcagccc
 24540
 aaagccctaa atctaaacct tcaatttctg gttgaaatca ttctcctgag cttccaaaa
 24600
 tgtggagcac tgaagaggag gagatggatg tgagacattt gggtgacttg gtgactgact
 24660
 gggataagg aaggagggga acagagaccg gcagcatgac tccagcctg ctgggctgga
 24720
 tggctgggtg atggtgagtc cattcaccaa actgggaggc ccagagagag aagcagattc
 24780

tgggctatgg aggatgaatg caggggtggag catgttgagt ctgttggtct cttgggacat
 24840
 ctggatggac atttccagaa ggcatatggg tatgtaaatc cacatagtag gccagctggc
 24900
 tggaaataca gatttaggag acagcagagt gaggacgggg atgaaaatgg tgggaatgga
 24960
 tgaggtcacc tatgaggtgt agagagagag ggtcgggagg ggaggatggg ccaagcttgc
 25020
 cctggccctg aaggaactgc aagctgggag cgctgagatg actgccctcc tggtgcttcc
 25080
 caggcctcgg ggctgtccgt gtggatgggg aagcagatgg agcccttgca cgcagtggcc
 25140
 ccggcagcca tcaccttgat cttgtccttg ctcggtgccc tgttactga gtgcacaagc
 25200
 aacgtggcca ccaccacctt gttcctgccc atctttgcct ccatggtaag taacctgaca
 25260
 gtggggagga gcccttccat ttcacaggaa cacatggcca tatttggggt ccctgacgag
 25320
 gcagcaatgt ccaggccaga ctacagaccag gctttggaga cccaggctctg actgtgacgt
 25380
 ggatttgtgg accctggatg cctctgcccc tgaggcctcc actgctttgc cactcctctt
 25440
 tgtaccctc ctgctgacca aagcaccaac catggacca gtgctcaa attttttata
 25500
 aatctaattg gattatTTTT caagctgggg agacaggact tgggctaagg aggagcaggc
 25560
 cagtgccgtg gtctctgagc atgtagcaca ggtgtgcagg aggactgcag actgggagca
 25620
 ccaactggctg gaaacccag gaagaggcct tggaggagtg gggacttggg agtaggtagg
 25680
 aaggagaga gaattctggg aagatggagc agcacaagga aaggcaatgg tgcacatgac
 25740
 tgaggactcc tggaagcctg gcttggtgag cacagggata agggatcctg gggagtggag
 25800
 agaggtagct gtcggttgtg ggaaaagctg ctgagtcca ggctaaggca ttctgttcta
 25860
 tggactagca tgttttttag ttgggagtta gaagaaagca gagcttatag gaaaatcagt
 25920
 ggctatgggt tttttttttt tttttttttt tttttttttt atgcatttcc ttctgtcatc
 25980
 cattgcaaag acgtaccagc ttcagggtag tatggaaaga tccctggtct cgcagtcaga
 26040
 agaccgaggt tcaagatgtg ggatctctga acatggccct tcagttcttt cttccgagag
 26100
 ctgtgctgat ggccaagtaa gatgagggt atgaaaagcc tctgtagact gcaaaatgag
 26160
 catgggagag gctgtcatta ttctggaatt gggagacaga tttacagagg gcctgaacac
 26220
 aggattgaag gtggtgaatt tccattcggc tgcctgggcg tctgcatgta taaaaagcaa
 26280
 acctaagtgg tttttttctc ctccaagtga agatgaaagt gttaaaaata gcaaggaggt
 26340
 gaaagtgttc aaaatagcaa agtggcctgt ctctcttct cctaagcaga ctgtccaaac
 26400
 agacgcccag tagaaggagc accttttgat actgggcacg tggtggtgat gcctcctctc
 26460
 tcctagcaca ggccgtggct ttgtcatctc cagccctaac tgggagcacc gagggtcca
 26520
 accaggcaaa tgcaggccct aacgggctct ttgaaaacgg gcttttctag aaccaggaac
 26580
 ctcaagtaaa aactcccca gctacctcta aggccatca cactcctgtc tcacgcccac
 26640
 ctatgagaaa ggaaaagggt atggtcattg agctgggctg cagaggagtg tgagggtgag
 26700
 acaccatgag gtaccacag ccaggaaaac gaggatggtc ggggagacgc gccagcgaag
 26760
 agctgggccc ctgcgtggga cccctcagtg gttcccaggg ggcgtgggac ttgcgcagtc
 26820

ctttcagagg gctgtttacc aacaggaacc gtaacattaa acctgctcag accccttgac
 26880
 tcagcaatatt catgtctggg aatataatctt aggaaaaataa tcagagatgc ctaccaacat
 26940
 atgtgatgat gatgtatgac agaattatta tacaatatata tccatagtaa caggggggttt
 27000
 gctgaaataa attatcatat attcatataa tatgacatta tcaggccatt aaaaatcaca
 27060
 gtttcaaaga gtaataaaat gggaacatgc tcatagtata gtttttttaa attgcagatg
 27120
 gtatatggct aaaaatgtct aataatgcaa agatgtatac agaccttaat cctctagcct
 27180
 cctccctaga gatgacctct gtaattttct caaatatttt tctggatatt ttacacactc
 27240
 acacactttt ttgagacag agtttcactc ttgtcaccca ggctggagtg caatgggtgtg
 27300
 atcttggctc actgcaacct ccacctcccg ggttcaagag attctcctgc ctcagcctcc
 27360
 cgagtagctg ggattacagg tgcctgccac cttgcctggc taattttttg tatttttagt
 27420
 agagacgggg ttccaccaca ttggtcaggc tgggtctcaa ctctgacct caggtgatcc
 27480
 gcctgccttg gcctcccaa gtgctgggat tacaggcgtg agccactgcg cccggccatt
 27540
 catcttaatt tttaaaaaat ctaaccatga agccttgggt atcttggaga gctttcctga
 27600
 ttgacacaaa aagaaaaaaa aatccaattc ttacagctg catactattc cattatttgt
 27660
 atgtgtcata ttttatttaa ccatcctgct attagtacc attgagttgg cttcctgtgt
 27720
 tttgccgtta catgggttga acaaacatgt ttgcatgtgt ctgccctcat gtgcatgata
 27780
 catgattgat ttgatagatt ttaggaatta catcattcat tcatacactc agcaaatatt
 27840
 taatgagtgc ctactctctg ataggtgctg ttggatgtgg ctaaaattta aagtgtagaa
 27900
 tttaaaaggt ggctacaaa ttccatgtgc aaaatgacct cagcatgta taaaaacaca
 27960
 cacatccaca gatttatatg cgggagagaa gatgtggtcc ctggcctcta ggctctctca
 28020
 gtctgtggca agacagacag acatgtgcac gcggcactgt aaggttgagc acagtctaag
 28080
 tactcagcat ggtctctggc acatagtagg tgccaagaa atacatgtcg aatgaattga
 28140
 gggggtaagg ccttctaggg caggtggcct ctgacctcag ccttcagtgt tccgtaggtg
 28200
 gaattatctg ccagagacgt ggcaaaagg agaggaacca agactgaggc acagagggtc
 28260
 aaacgtaccc ggcacattca gagaatcctt ttcagaatca cgtccccaag agcttctgtg
 28320
 ttctgtacgg tgatgttga gtgctgtttt tccgcagtct cgctccatcg gcctcaatcc
 28380
 gctgtacatc atgctgccct gtacctgag tgcctccttt gccttcatgt tgctgtggc
 28440
 caccctcca aatgccatcg tgttcaccta tgggcacctc aaggttgctg acatggtaac
 28500
 acagctgttt ttatttactc ccgtcggact ataacgtgt tgtcataagg gatgccccat
 28560
 ttatgaatga cagagtttca aaacgatgtc atgtgacttg ggaatgccac ggaacatcca
 28620
 gacctgtagc cattgttgac atttataatg cagcttttct tctttttctg agatgatctc
 28680
 aagcctcaca cactgttctt tctctgaggt gggttataga ctctcccacc tggagaagcc
 28740
 tgtgcaggca ccaggggag ccttggaagg ggtgaagggt gggctgaggg actcatatgg
 28800
 ccaaggatga acttgacaaa ttagcaagaa ccatgaagat aggcagggca ggcttaggca
 28860

gcagggggat gctaatagaca gtcacagaga tttgtagggg tgcctgaaga ggtagaagca
 28920
 gggagagggg gagagagagc actgcctggg agtagatgat gccttggaac caaatgtagt
 28980
 cagaggaaga actcttcatt agctctgtca cctttgctgg gagaagggca gctttgcagc
 29040
 tctgggctgg gaaagaggca agtgtttgag cccaagaggc cagaaatgta cctgggacca
 29100
 atcgggtgtt cgttatctca gagcctctgc tgggtatctc agggactcca tgagcatttt
 29160
 caaaaaaaaa ggtgggtccc agaaacatg gactgcaaac ttgactccaa tccccagta
 29220
 aaatatctac aacagggtag tgaagcgatg gttagtacc atgagggaag cttgcagagc
 29280
 aggcacaga aagagcctga ggaggtccac agggaaagctg gcacgtcctt gtaggatagt
 29340
 taaggcactg gggtagcaa tgaacctgga ctacaggaac actgggctct gtgaccgttt
 29400
 ccctgaatgg cctaagctgt tgcctcctgt cacttctctg aggtcatttt ccaaatgcgc
 29460
 acgggcatag agaaccatc cactctgcct acttcccagg gatgccttga gactgagga
 29520
 tacctggggg acatgaagtc gactgtcctt gggggtcggg acacccagc caggagacaga
 29580
 gcatggcaca gggacatcga ggcccagtga gccgaccctt tgcctcctc tctgagagca
 29640
 ctatgtccca gcaggcctca ggggtgctgac tctgtctctt ttccagggtga aaacaggagt
 29700
 cataatgaac ataattggag tcttctgtgt gtttttggct gtcaacacct ggggacgggc
 29760
 catatttgac ttggatcatt tccctgactg ggctaagtgt acacatattg agacttagga
 29820
 agagccacaa gaccacacac acagccctta ccctcctcag gactaccgaa ctttctggca
 29880
 caccttgtac agagttttgg ggttcacacc ccaaatgac ccaacgatgt ccacacacca
 29940
 ccaaaacca gccaatgggc cacctcttcc tccaagccca gatgcagaga tggatcatggg
 30000
 cagctggagg gtaggctcag aaatgaaggg aacccctcag tgggctgctg gacccatctt
 30060
 tcccaagcct tgccattatc tctgtgaggg aggccaggta gccgagggat caggatgcag
 30120
 gctgctgtac ccgctctgcc tcaagcatcc cccacacagg gctctggttt tcaactcgtt
 30180
 cgtcctagat agtttaaag ggaatcgat cccctgggtg agagctaaga caaccaccta
 30240
 ccagtgccca tgtcccttcc agctcacctt gacgagcctc agatcatctc tgtcactctg
 30300
 gaaggacac ccagccagg gacggaatgc ctggtcttga gcaacctccc actgctggag
 30360
 tgcgagtggg aatcagagcc tcctgaagcc tctgggaact cctcctgttg ccaccacaa
 30420
 aggatgagga atctgagttg ccaactcag gacgacacct ggcttgccac ccacagtga
 30480
 ccacaggcca acctacgcc ttcatcactt ggttctgttt taatcgactg gccccctgtc
 30540
 ccacctctcc agtgagcctc cttcaactcc ttgggtccct gttgtctggg tcaacatttg
 30600
 ccgagacgcc ttggctggca ccctctgggg tccccctttt ctcccaggca ggtcatcttt
 30660
 tctgggagat gcttcccctg ccaccccaa atagctagga tcacactcca agtatgggca
 30720
 gtgatggcgc tctgggggcc acagtgggct atctaggtcc tccctcacct gaggccaga
 30780
 gtggacacag ctgttaattt cactggcta tgccacttca gagtctttca tgccagcgtt
 30840
 tgagctcctc tgggtaaaat cttccctttg ttgactggcc ttcacagcca tggctggtga
 30900

caacagagga tcgttgagat tgagcagcgc ttggtgatct ctcagcaaac aacccttgcc
 30960
 cgtgggccaa tctacttgaa gttactcgga caaagacccc aaagtggggc aacaactcca
 31020
 gagaggctgt gggaatcttc agaagccccc ctgtaagaga cagacatgag agacaagcat
 31080
 cttctttccc ccgcaagtcc attttatttc cttcttggtc tgctctggaa gagaggcagt
 31140
 agcaaagaga tgagctcctg gatggcattt tccagggcag gagaaagtat gagagcctca
 31200
 ggaaacccca tcaaggaccg agtatgtgtc tggttccttg ggtgggacga ttcctgacca
 31260
 cactgtccag ctcttgctct cattaaatgc tctgtctccc gcgaaagct ccactgtgct
 31320
 gctgacttgt ctctggtttt ctgcagtgtg gggagcccag ggaggtggat gaatgaacag
 31380
 ttagttacgc cctgccacc tgctgggtgc caggccttcc tgtccctgtt gaatccacta
 31440
 gttatctcgt tcgatctttg cagcaaccct gtgagatagg aagggtgtat tatcttgctt
 31500
 tgtctttcaa aaaaaagcg aggtcaggg aggccaaagg aagtgtccaa agtcacacat
 31560
 caagttactg gcagttacag ttccaaccaa gagcttccaa ctccataccc cctgctcctt
 31620
 ctgctagcca tgaagggtt tggccttata gggctttagt ggaaaggtga gtggccaaga
 31680
 gcaagtccat gccaaaggaa gatctccaaa catgagtccc tgtctgttgc ctcccctgag
 31740
 ataggcacag gacaagtgat caatgagaca ggggtgtcct tgccctaaga agcaaagtgt
 31800
 ttggttgggg agggaaagtag ggaaaggct gccacctccc cccaccaagg tacaactgct
 31860
 gacttccttc ctcccagcc ctctatcact gccctctgtg ccgctgccgt tgactggcct
 31920
 gccccaccag actgagggt ctgactgcc accgagtcta gtgtcagcat tatggctgac
 31980
 ccagagcagg ctatacagtt agtatgatgg ataaataaat gattggtcag tgcagtcaat
 32040
 taggtgcaag ctgttgtag taggcaagg caatgaagg catccaagg gggcattgaa
 32100
 ggatgagtag aatggccagg ggtaatggg gaggaactgg tgggtgggtg gaggactctt
 32160
 ccagacacca tgtggttag ggctgacaaa aagctgggtg gagggcttcc agagtgccaa
 32220
 gctcccacct gaagaggctg accagaggcc aatcctaacc aactctaggt gttggctgga
 32280
 gttgactaa agtgtatggc ctcccacc aaacctttg cttcttaggg caaggaccac
 32340
 cctgtctcat tgatcactgt cctgagccta tctcagggga ggcaaaagag agggacctgt
 32400
 attcagagat cttcccttg catgacttgc ttttggccac ttaccttcc ctacaagctc
 32460
 tatgaggcca aggcccttca tggtagtgt aaggagcagt gggcatggag ttggaagatc
 32520
 tgggttgga cggtactgc cactaactcg atgtgtgatt ctgaacactt aacttagcca
 32580
 tacatgctct cttatttct tttgatggca aataagagaa gggccagcaa acagtggctt
 32640
 aaaccagaag gtcaattaat gtttactttt caggaagtct gtaggtagat ggttgctggc
 32700
 attggcccaa cagctcattt cagcctccaa ggacttgcgc tccatagtcc actctgtcat
 32760
 cttaaagcct tcacactttt acccccatgc ttgaccccca ggctacatac acagct
 32816

<210> 4

<211> 619

<212> PRT

<213> *Xenopus laevis*

<400> 4

```

Met Val Ser Ile Gly Lys Trp Ile Leu Ala Asn Arg Asn Tyr Phe Ile
 1          5          10          15
Ile Phe Leu Val Pro Leu Phe Leu Leu Pro Leu Pro Leu Val Val Pro
 20          25          30
Thr Lys Glu Ala Ser Cys Gly Phe Val Ile Ile Val Met Ala Leu Phe
 35          40          45
Trp Cys Thr Glu Ala Leu Pro Leu Ala Val Thr Ala Leu Phe Pro Val
 50          55          60
Leu Leu Phe Pro Met Met Gly Ile Met Asp Ser Thr Ala Val Cys Ser
 65          70          75          80
Gln Tyr Leu Lys Asp Thr Asn Met Leu Phe Ile Gly Gly Leu Leu Val
 85          90          95
Ala Ile Ser Val Glu Lys Trp Asn Leu His Lys Arg Ile Ala Leu Arg
 100         105         110
Val Leu Leu Ile Val Gly Val Lys Pro Ala Leu Leu Leu Gly Phe
 115         120         125
Met Val Val Thr Ala Phe Leu Ser Met Trp Ile Ser Asn Thr Ala Thr
 130         135         140
Thr Ala Met Met Ile Pro Ile Ala Gln Ala Val Met Glu Gln Leu His
 145         150         155         160
Ser Ser Glu Gly Lys Val Asp Glu Arg Val Glu Gly Asn Ser Asn Thr
 165         170         175
Gln Lys Asn Val Asn Gly Met Glu Asn Asp Met Tyr Glu Ser Val Met
 180         185         190
Pro Ser Gly Lys Met Ala Leu Ala Ile Asp Asn Thr Tyr Ala Thr Glu
 195         200         205
Asn Glu Gly Phe Glu Ile Gln Glu Lys Ser Thr Lys Asp Pro Glu Pro
 210         215         220
Ser Lys Gln Glu Lys Gln Ser Ile Gly Pro Ile Val Ile Glu Pro Glu
 225         230         235         240
Asp Glu Lys Gln Thr Glu Glu Lys Gln Lys Glu Lys His Leu Lys Ile
 245         250         255
Cys Lys Gly Met Ser Leu Cys Val Cys Tyr Ser Ala Ser Ile Gly Gly
 260         265         270
Ile Ala Thr Leu Thr Gly Thr Thr Pro Asn Leu Val Met Lys Gly Gln
 275         280         285
Met Asp Glu Leu Phe Pro Glu Asn Asn Asn Ile Ile Asn Phe Ala Ser
 290         295         300
Trp Phe Gly Phe Ala Phe Pro Thr Met Leu Val Leu Leu Ala Leu Ser
 305         310         315         320
Trp Leu Trp Leu Gln Phe Ile Tyr Leu Gly Val Asn Phe Lys Lys Asn
 325         330         335
Phe Gly Cys Gly Gly Asn Ala Glu Gln Lys Glu Lys Glu Lys Arg Ala
 340         345         350
Phe Arg Val Ile Ser Gly Glu His Lys Lys Leu Gly Ser Met Thr Phe
 355         360         365
Ala Glu Ile Ser Val Leu Val Leu Phe Ile Leu Leu Val Leu Leu Trp
 370         375         380
Phe Thr Arg Glu Pro Gly Phe Met Pro Gly Trp Ala Thr Ile Ser Phe
 385         390         395         400
Asn Lys Gly Gly Lys Glu Met Val Thr Asp Ala Thr Val Ala Ile Phe
 405         410         415
Val Ser Leu Met Met Phe Phe Phe Pro Ser Glu Leu Pro Ser Phe Lys
 420         425         430
Tyr Gln Asp Thr Asp Lys Pro Gly Met Lys Pro Lys Leu Arg Val Pro
 435         440         445
Pro Ala Leu Leu Asp Trp Lys Thr Val Asn Glu Lys Met Pro Trp Asn
 450         455         460
Ile Val Ile Leu Leu Gly Gly Gly Phe Ala Leu Ala Lys Gly Ser Glu
 465         470         475         480
Glu Ser Gly Leu Ser Leu Trp Leu Gly Glu Lys Leu Thr Pro Leu Gln
 485         490         495
Ser Ile Pro Pro Ala Ala Ile Ala Leu Ile Leu Cys Leu Leu Val Ala
 500         505         510
Thr Phe Thr Glu Cys Thr Ser Asn Val Ala Thr Thr Thr Leu Phe Leu

```

	515		520		525										
Pro	Ile	Leu	Ala	Ser	Met	Ala	Lys	Ala	Ile	Gln	Leu	Asn	Pro	Leu	Tyr
	530					535					540				
Ile	Met	Leu	Pro	Cys	Thr	Leu	Ser	Ala	Ser	Leu	Ala	Phe	Met	Leu	Pro
	545					550					555				560
Val	Ala	Thr	Pro	Pro	Asn	Ala	Ile	Ala	Phe	Ser	Tyr	Gly	Gln	Leu	Lys
				565						570					575
Val	Ile	Asp	Met	Ala	Lys	Ala	Gly	Leu	Leu	Leu	Asn	Ile	Leu	Gly	Val
			580					585					590		
Leu	Thr	Ile	Thr	Leu	Ala	Ile	Asn	Ser	Trp	Gly	Phe	Tyr	Met	Phe	Asn
		595					600					605			
Leu	Gly	Thr	Phe	Pro	Ser	Trp	Ala	Asn	Ala	Thr					
	610					615									

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/046407 A3

(51) International Patent Classification⁷: C12N 15/12,
15/11, 15/63, 1/21, 5/10, C07K 14/705, 16/28, A01K
67/027, C12Q 1/68, G01N 33/68

(21) International Application Number: PCT/US01/45661

(22) International Filing Date: 5 December 2001 (05.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/729,094 5 December 2000 (05.12.2000) US

(71) Applicant: PE CORPORATION (NY) [US/US]; 761
Main Avenue, Norwalk, CT 06859 (US).

(72) Inventors: CHATURVEDI, Kabir; c/o Celera Genomics
Corporation, 45 West Gude Drive C2-4 #21, Rockville,
MD 20850 (US). WEI, Ming-Hui; c/o Celera Genomics,
45 West Gude Drive C2-4#21, Rockville, MD 20850
(US). KETCHUM, Karen, A.; c/o Celera Genomics, 45
West Gude Drive C2-4#21, Rockville, MD 20850 (US).
DIFRANCESCO, Valentina; c/o Celera Genomics, 45
West Gude Drive C2-4#21, Rockville, MD 20850 (US).
BEASLEY, Ellen, M.; c/o Celera Genomics, 45 West
Gude Drive C2-4#21, Rockville, MD 20850 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report:
13 March 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/046407 A3

(54) Title: ISOLATED HUMAN TRANSPORTER PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN TRANSPORTER PROTEINS, AND USES THEREOF

(57) Abstract: The present invention provides amino acid sequences of peptides that are encoded by genes within the human genome, the transporter peptides of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the transporter peptides, and methods of identifying modulators of the transporter peptides.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/45661

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/11 C12N15/63 C12N1/21 C12N5/10
C07K14/705 C07K16/28 A01K67/027 C12Q1/68 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A01K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, GENSEQ, SEQUENCE SEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
L	US 2002/019028 A1 (BEASLEY ELLEN M ET AL) 14 February 2002 (2002-02-14) L:priority figures 1,2	1-23
L	--- DATABASE GENSEQ 'Online! 11 January 2002 (2002-01-11) TANG YT ET AL: "Human NaDC-2 homologue-encoding cDNA, SEQ ID NO:1284" Database accession no. ABA09508 XP002222098 L document cited to provide information on the relevant sequence disclosed in WO 01 57188 the whole document	1-5,8-16
P,X	& WO 01 57188 A (HYSEQ INC ;LIU CHENGHUA (US); TANG Y TOM (US); DRMANAC RADOJE T (US)) 9 August 2001 (2001-08-09) --- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

27 November 2002

Date of mailing of the international search report

11/12/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Espen, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/45661

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
L	<p>DATABASE GENSEQ 'Online! 11 January 2002 (2002-01-11) TANG YT ET AL: "Human NaDC-2 homologue, SEQ ID NO:2634" Database accession no. ABB12264 XP002222099 L document cited to provide information on the relevant sequence disclosed in WO 01 57188 A the whole document</p>	1-5,8-16
P,X	<p>& WO 01 57188 A (HYSEQ INC ; LIU CHENGHUA (US); TANG Y TOM (US); DRMANAC RADOJE T (US)) 9 August 2001 (2001-08-09) ---</p>	
E	<p>WO 02 12340 A (INCYTE GENOMICS) 14 February 2002 (2002-02-14) SEQ ID NOs 25,55 ---</p>	1-5, 7-16, 19-23
P,X	<p>WO 01 57274 A (CHEN WENSHENG ; HANZEL DAVID K (US); PENN SHARRON G (US); RANK DAVI) 9 August 2001 (2001-08-09) SEQ ID NOs 15609,25711, and 26073 ---</p>	1-6,8-16
P,X	<p>DATABASE GENSEQ 'Online! 16 October 2001 (2001-10-16) TANG YT ET AL: "Human protein encoding cDNA sequence SEQ ID NO:448" Database accession no. AAH99613 XP002222100 PX document cited to provide information on the relevant sequence disclosed in WO 01 53455 the whole document</p>	1-5,8-16
P,X	<p>& WO 01 53455 A (HYSEQ INC ; LIU CHENGHUA (US); TANG Y TOM (US); DRMANAC RADOJE T (US)) 26 July 2001 (2001-07-26) ---</p>	
P,X	<p>DATABASE GENSEQ 'Online! 16 October 2001 (2001-10-16) TANG YT ET AL: "Human protein sequence SEQ ID NO:1187" Database accession no. AAM25672 XP002222101 PX document cited to provide information on the relevant sequence disclosed in WO 01 53455 A2 the whole document</p>	1-5,8-16
P,X	<p>& WO 01 53455 A (HYSQ INC; LIU CHENGHUA (US); TANG Y TOM (US); DRMANAC RODOJE T (US)) 26 July 2001 (2001-07-26) ---</p>	
	-/--	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/45661

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SEKINE T ET AL: "Expression cloning and characterization of a novel multispecific organic anion transporter" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 272, no. 30, 25 July 1997 (1997-07-25), pages 18526-18529, XP002097953 ISSN: 0021-9258 -----	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/45661

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 17, 18
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 17,18

Present claims 17 and 18 refer to an agent defined by reference to a desirable characteristic or property, namely that binds to any of the peptides of claim 2.

The claims cover all agents having this characteristic or property, whereas the application provides no support within the meaning of Article 6 PCT and no disclosure within the meaning of Article 5 PCT. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the agent as a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search impossible.

Consequently, no search has been carried out for claims 17 and 18.

Nucleic acid molecules relating to SEQ ID NO 3 having a length of 32816 nucleotides, were not searched, since it is not apparent what is the relationship between the human transporter proteins claimed and this lengthy nucleic acid sequence.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/45661

Patent document cited in search report		Publication date /	Patent family member(s)	Publication date
US 2002019028	A1	14-02-2002	AU 3944402 A WO 0246407 A2	18-06-2002 13-06-2002
WO 0157188	A	09-08-2001	AU 2591801 A AU 3128801 A AU 3297101 A AU 3300301 A AU 3329301 A AU 3484701 A AU 3484801 A AU 3486501 A AU 3494401 A AU 3665801 A AU 3666001 A AU 3666301 A AU 3672101 A AU 4314201 A EP 1254265 A1 EP 1254266 A1 EP 1254268 A1 EP 1254269 A1 EP 1254270 A1 WO 0153326 A1 WO 0155334 A2 WO 0155335 A2 WO 0157255 A1 WO 0157260 A1 WO 0157175 A2 WO 0157261 A1 WO 0157262 A1 WO 0157187 A2 WO 0157265 A1 WO 0157188 A2 WO 0157266 A1 WO 0157267 A1 WO 0157190 A2 US 2002128187 A1 US 2002137044 A1 US 2002127199 A1	31-07-2001 14-08-2001 07-08-2001 07-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 06-11-2002 06-11-2002 06-11-2002 06-11-2002 06-11-2002 26-07-2001 02-08-2001 02-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 09-08-2001 12-09-2002 26-09-2002 12-09-2002
WO 0212340	A		NONE	
WO 0157274	A	09-08-2001	AU 3087801 A AU 3087901 A AU 3088001 A AU 3088101 A AU 3088201 A AU 3088301 A AU 3275701 A AU 3275801 A AU 3275901 A AU 3276001 A AU 3311401 A AU 3658901 A AU 6343201 A AU 9295701 A GB 2361238 A ,B GB 2373500 A	14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 20-11-2001 14-08-2001 14-08-2001 14-08-2001 14-08-2001 11-12-2001 02-04-2002 17-10-2001 25-09-2002

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/45661

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0157274	A	GB 2374929 A	30-10-2002
		GB 2375539 A	20-11-2002
		GB 2375111 A	06-11-2002
		GB 2374872 A	30-10-2002
		WO 0157270 A2	09-08-2001
		WO 0157271 A2	09-08-2001
		WO 0157272 A2	09-08-2001
		WO 0157273 A2	09-08-2001
		WO 0186003 A2	15-11-2001
		WO 0157274 A2	09-08-2001
		WO 0157275 A2	09-08-2001
		WO 0157276 A2	09-08-2001
		WO 0157277 A2	09-08-2001
		WO 0157278 A2	09-08-2001
		WO 0157251 A2	09-08-2001
		WO 0157252 A2	09-08-2001
		WO 0192524 A2	06-12-2001
		US 2002081590 A1	27-06-2002
		US 2002048763 A1	25-04-2002
		US 2002048800 A1	25-04-2002
		US 2002123474 A1	05-09-2002
		GB 2360284 A , B	19-09-2001
		EP 1158049 A1	28-11-2001
		US 2002102252 A1	01-08-2002
		AU 9481201 A	08-04-2002
		WO 0226818 A2	04-04-2002
WO 0153455	A	26-07-2001	
		AU 2292401 A	31-07-2001
		AU 2591801 A	31-07-2001
		AU 2593601 A	31-07-2001
		AU 2595501 A	31-07-2001
		AU 2596501 A	31-07-2001
		AU 2598301 A	31-07-2001
		AU 2728401 A	31-07-2001
		AU 2734401 A	31-07-2001
		AU 2734801 A	31-07-2001
		AU 2738501 A	31-07-2001
		AU 3265701 A	31-07-2001
		EP 1242596 A1	25-09-2002
		EP 1240178 A2	18-09-2002
		EP 1242580 A1	25-09-2002
		EP 1242443 A1	25-09-2002
		EP 1250346 A2	23-10-2002
		EP 1254256 A2	06-11-2002
		EP 1248848 A1	16-10-2002
		WO 0153312 A1	26-07-2001
		WO 0153453 A2	26-07-2001
		WO 0153326 A1	26-07-2001
		WO 0153454 A2	26-07-2001
		WO 0153455 A2	26-07-2001
		WO 0153456 A2	26-07-2001
		WO 0153466 A1	26-07-2001
		WO 0152616 A2	26-07-2001
		WO 0153500 A1	26-07-2001
		WO 0153515 A1	26-07-2001
		WO 0153485 A1	26-07-2001
		US 6465620 B1	15-10-2002
		US 2002146692 A1	10-10-2002

Form PCT/ISA/210 (patent family annex) (July 1992)